

Systematics and diversity of Neotropical ants

Sistemática y diversidad de las hormigas neotropicales

FERNANDO FERNÁNDEZ¹; ROBERTO J. GUERRERO²;
ANDRÉS F. SÁNCHEZ-RESTREPO³

¹ Ph. D. Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Carrera 30 No. 45 - 03, Bogotá D.C. Colombia, ffernandezca@unal.edu.co, <https://orcid.org/0000-0002-6862-3592>. ² Ph. D. Grupo de Investigación Insectos Neotropicales, Programa de Biología, Facultad de Ciencias Básicas, Universidad del Magdalena, Carrera 32 # 22-08, Santa Marta, Colombia, rguerrero@unimagdalena.edu.co, <https://orcid.org/0000-0003-3244-2754>. ³ Ph. D. Fundación para el Estudio de Especies Invasivas (FuEDEI), Hurlingham, Buenos Aires, Argentina. y/o Consejo Nacional de Investigaciones Científicas y Técnicas (Conicet), Buenos Aires, Argentina, andres.sanchez@javeriana.edu.co, <https://orcid.org/0000-0001-7514-068X>.

Abstract: With a long history of more than 100 million years and about 14,000 described living species, ants are one of the most important and well-known groups of insects in the world. Ants are key elements in the structure and dynamics of terrestrial systems, especially in the tropics, as well as models in studies of evolution, ecology, and monitoring of disturbed ecosystems. For all this, it is essential to know their history and phylogeny, in order among other things, to have a solid base for their systematics and taxonomy. Within Aculeata ants appear to be the sister group of Apoidea, and the ancestor of the Formicidae may have inhabited the northern hemisphere during the early Cretaceous. Since then, and especially since the Eocene, ants have spread throughout the planet, occupying almost all environments from forests to deserts. The Neotropical region has 137 genera and around 3,100 species of ants. The Neotropics seem to have been the “cradle” and “museum” of the groups of ants, which would explain their great diversity and a high degree of endemisms. This review describes the current state of knowledge of ants in the Neotropical region from a systematic point of view, with a synopsis of all supraspecific taxa described to date. Critical genera, problems to be solved, and perspectives for the study of these insects are also presented.

Keywords: Evolutionary patterns, phylogenetic hypotheses, species delimitation, taxonomy, Hymenoptera.

Resumen: Con una larga historia de algo más de 100 millones de años y alrededor de 14.000 especies vivientes descritas, las hormigas son uno de los grupos de insectos más importantes y conocidos en el mundo. Las hormigas son elementos clave en la estructura y dinámica de los sistemas terrestres, en especial en los trópicos, así como herramienta en estudios de evolución y ecología, y monitoreo de ecosistemas perturbados. Por todo esto es indispensable conocer su historia y filogenia para, entre otras cosas, tener una sólida base de su sistemática y, finalmente, de su taxonomía. Dentro de Aculeata las hormigas parecen ser el grupo hermano de Apoidea y el ancestro de Formicidae pudo haber habitado el hemisferio norte a comienzos del Cretáceo. A partir de entonces, en especial del Eoceno, las hormigas se han diseminado por todo el planeta y ocupan casi todos los ambientes, desde bosques a desiertos. La región Neotropical comprende 137 géneros y alrededor de 3.100 especies. El Neotrópico parece haber sido “cuna” y “museo” de los grupos de hormigas, lo que explicaría su gran riqueza y alto grado de endemismos. Esta revisión muestra el estado actual de conocimiento de las hormigas en la región Neotropical desde el punto de vista sistemático, con una sinopsis de todos los taxones supraespecíficos descritos hasta la fecha. También se presentan géneros críticos, problemas por resolver y perspectivas para el estudio de estos insectos.

Palabras clave: Patrones evolutivos, hipótesis filogenéticas, delimitación de especies, taxonomía, Hymenoptera.

Introduction

Ants are one of the most important groups of insects in the world, with an elevated presence and influence on the structure and dynamics of terrestrial ecosystems, especially in the tropics and their low-altitude areas. Due to their abundance and varied biology, they are useful tools for conservation studies and for monitoring threatened areas. Knowledge about the evolutionary history, phylogenetic relationships, and the classification and identification of the species of Neotropical ants, provides biologists and naturalists updated information for the recognition of the genera and species of the region.

Corresponding author

Fernando Fernández. Ph. D. Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Carrera 30 No. 45 - 03, Bogotá D.C. Colombia, ffernandezca@unal.edu.co, <https://orcid.org/0000-0002-6862-3592>

Suggested citation

FERNÁNDEZ, F.; GUERRERO, R. J.; SÁNCHEZ-RESTREPO, A. F. 2021. Systematics and diversity of Neotropical ants. Revista Colombiana de Entomología 47 (1): e11082. <https://doi.org/10.25100/socolen.v47i1.11082>

Received: 17-Mar-2021

Accepted: 24-Mar-2021

Published: 15-Apr-2021

Revista Colombiana de Entomología

ISSN (Print): 0120-0488

ISSN (On Line): 2665-4385

<https://revistacolombianaentomologia.univalle.edu.co>

Open access

 creativecommons.org/licenses/by-nc-sa/4.0/deed.es

Publishers: Sociedad Colombiana de Entomología
SOCOLEN (Bogotá, D. C., Colombia)
<https://www.socolen.org.co>
Universidad del Valle (Cali, Colombia)
<https://www.univalle.edu.co>

© 2021 Sociedad Colombiana de Entomología -
SOCOLEN y Universidad del Valle - Univalle

Ants have long been considered a natural group within the Hymenoptera. The possession of the metapleural gland is one of the unique traits in ants, considered the morphological synapomorphy par excellence, even visible in some fossils of Cretaceous amber of almost 100 million years old (Wilson *et al.* 1967; Agosti *et al.* 1998; Barden and Grimaldi 2013, 2014; LaPolla *et al.* 2013; Barden 2017; Barden *et al.* 2017; Lattke and Melo 2020). The presence of a petiole, geniculated antennae, infrabuccal sac, deciduous wings in females, and perennial acellular colonies with workers (always apterous females), are additional and unique features within the Hymenoptera (Bolton 2003; Taylor 2007).

Boudinot (2015) offers more unique characters in ants, with the inclusion of new and comparative information on males. Boudinot *et al.* (2020) provide a diagnosis of the Formicoidea superfamily based especially on Mesozoic specimens. These authors propose the name Formicapoidina for Formicoidea plus Apoidea (sphecid wasps and bees), and they list 10 traits to define Formicoidea, among those: the elongation of the postgenal bridge; the mandibular condyle and the procoxae; “petiolization” of the first metasomal segment; prora (anteroventral process of the second external metasoma); and angled junction of Rs and M in the forewing. They recognize two clades, the basal Camelomecia clade and the total clade Formicidae, which in turn is divided into the clades Sphecomyrmes (with three Cretaceous subfamilies: Haidomyrmecinae, Zigrasimeciinae, and Sphecomyrmicinae) and Antennoclypeata (with Cretaceous Brownimeciinae and Crown clade Formicicidae). Borysenko (2017) compares plesiomorphic and autoapomorphic characters to separate “basal” ants (stem groups) from modern ants (crown groups) and performs a first statistical analysis of the scape lengths and flagellomeres of worker antennae to separate the basal (and extinct) groups of modern ants. This author uses criteria from the Phylum Code (PhyloCode) to define the “crown” group of ants such as Formicidae Stephens, 1829; the clade originated with the most recent common ancestor of *Martialis heureka* Rabeling and Verhaagh, 2008, and *Formica rufa* Linnaeus, 1761. This clade then corresponds to the family Formicidae Latreille, 1809, excluding the basal groups. Pan-Formicidae Borysenko, 2017, is defined as “the total clade composed of the crown clade FormicidaeP and all extinct species that share a more recent common ancestor with FormicidaeP rather than with any living species that is not a member of FormicidaeP”. Similarly, Borysenko (2017) defines all the subfamilies of Formicidae under the same criteria. One of the various consequences of the Borysenko (2017) study is that clades such as Armaniidae / Armaniinae could be included within Sphecomyrmicinae, and Cretaceous genera such as *Brownimecia* would remain in the main group of ants.

With the internal phylogeny of Aculeata using four nuclear genes, Pilgrim *et al.* (2008) confirm the monophyly of Aculeata, Chrysoidea, and Apoidea, but they establish the paraphyly of Vespoidea, proposing several superfamilies and suggesting a proximity between apoids and ants, a new and unexpected relationship. The same relationship was confirmed in the analysis using genomes (“phylogenomics”) by Johnson *et al.* (2013) where the relationship between apoids and ants is confirmed (Fig. 1). These relationships, if stable, are very important not only when looking for suitable outside groups, but also when establishing suitable scenarios in the first steps of the evolution of ants. An ancestor between ants and neighboring groups no longer consists of an

ectoparasitoid wasp (according to the phylogeny of Brothers 1999), but of a wasp that digs a nest in the ground that it supplies at once or gradually with prey, as occurs in Ampulicidae and Sphecidae, which are basal apoids. This scenario may make the recent proposal of the first soil-dwelling ants more plausible, subsequently originating lineages that would colonize the litter and trees (Lucky *et al.* 2013).

The recent publication of the phylogeny and history of Hymenoptera by Peters *et al.* (2017) offers a more stable panorama since several clades proposed by morphology and genes are confirmed. This includes the paraphyly of Aculeata and Chrysoidea, and Vespoidea as defined by Brothers (1999) and the sister group relationships between ants and Apoidea. This position implies that the ants must comprise the superfamily Formicoidea with a family (Formicidae) or two (depending on how the nature of the extinct taxon Armaniidae / Armaniinae is interpreted). On the other hand, the ancestor of this Jurassic clade suggests the origin of the ants towards the Upper Jurassic or the Jurassic-Cretaceous transition (Moreau and Bell 2013).

Branstetter *et al.* (2017) use ultra-conserved elements (UCE) to explore the internal relationships of Aculeata with an emphasis on ants and Apoidea. This study offers strong support for the relationship between ants and Apoidea and an Upper Jurassic or Lower Cretaceous antiquity, as in some previous studies. With this series of data, the position of ants within Aculeata appears to be clear. However, Camacho *et al.* (2018) draw attention to the incongruities in the trees that show internal relationships in Aculeata, especially between ants and apoids. These authors suggest the need for more data and different analytic dynamics, in addition to the standard ones, to improve our understanding of the relationships between ants and other aculeates.

Bolton (2003) provides a list of ant synapomorphies based on morphological traits; Boudinot (2015), in the first




Figure 1. Phylogenetic relationship within Hymenoptera Aculeata, after Johnson *et al.* (2013). Social lineages marked with red points at tips.

synthesis of ants from the point of view of male morphology, recapitulates and expands these synapomorphies. Naturally, several of these undergo changes such as secondary loss or modifications, or they do not apply (see Bolton (2003), and Boudinot (2015) for exceptions, and Taylor (2007)). Boudinot (2015), offers other attributes that help to characterize ants, even if they are not synapomorphies: antenna with 4 to 12 flagellomeres in the female or 5 to 13 in the male; bulbous neck (radicle) and scape with a common axis; epicnemius very small, not visible in situ; abdominal segment II with sternum and tergus equally sclerotized; variable wing venation, with extreme reductions, without closed cells; jugal lobe present or absent; abdominal sternum IX can be complex and have apical modifications (such as projections, teeth, and lobes).

Currently, 16 subfamilies of living ants are recognized (12 in the Neotropics: Heteroponerinae is included within the Ectatomminae: Feitosa and Prada-Achiardi 2019), all of them are monophyletic (Ward 2010, 2014; Borowiec *et al.* 2019). Table 1 lists the world's living and fossil ant subfamilies.

Phylogenies based on morphology

Keller (2011) performs the only modern study on the phylogeny of ants using external morphology of females with an emphasis on poneromorphs. This study is based on 139 characters (many of them new) and 105 terminals (taxa), covering most subfamilies and genera (except Martialinae). The main result is that the poneromorph ants form a paraphyletic group, as they include dorylomorphs, leptanilomorphs, and myrmicomorphs. Although this research agrees with previous studies questioning the monophyly of poneromorphs (Moreau *et al.* 2006; Brady *et al.* 2006), it does not agree for other groupings as suggested by gene-based phylogenies. Cantone and von Zuben (2019) use hindwing venation to explore the phylogeny of ants. These authors propose three types of wing venation morphology; the first one (I) is the most primitive; and II and III are derived by reduction. Part of the hypotheses of these authors suggest a high degree of convergence between subfamilies and unrelated genera.

Table 1. Fossil (*) and living taxa of the world, with distribution and habits. Subfamilies present in the Neotropics in bold. Boudinot *et al.* (2020) relate several Mesozoic genera as uncertain.

TAXON	DISTRIBUTION	NOTES
FORMICOIDEA		
Armaniidae*	Cretaceous, Northern Hemisphere.	Unknown biology.
FORMICIDAE		
BROWNIMECIINAE*	Cretaceous, USA.	Unknown biology. Specialized mandibles.
FORMICIINAE*	Eocene, Europa.	
HAIDOMYRMECINAE*	Cretaceous, Burma, northern Myanmar (Asia).	Unknown biology. Specialized mandibles.
SPHECOMYRMINAE*	Cretaceous, Northern Hemisphere.	Unknown biology.
ZIGRACIMECIINAE*	Cretaceous, Burma, Canada.	Unknown biology
MARTIALINAE	Brazil.	Underground, unknown biology; probably specialized predator.
LEPTANILLINAE	Old World.	Underground, specialized predator.
AGROECOMYRMECINAE	Neotropics.	Soil; unknown biology; probably specialized predator.
AMBLYOPONINAE	Worldwide.	Soil. Mostly predators.
APOMYRMINAE	Afrotropical.	Soil. Predators.
PARAPONERINAE	Neotropical.	Mainly on the ground, nest under tree stumps; omnivorous.
PONERINAE	Worldwide, mostly tropical.	Mainly on the ground, with not very large colonies; great diversity of social biology (with some species without queens) and feeding habits.
PROCERATIINAE	Worldwide.	Mainly on the ground, general to specialized predators.
ANEURETINAE	Sri Lanka.	Wood. Predator.
DOLICHODERINAE	Worldwide, mainly temperate.	Wide range of nesting habits, behavior and food sources. Includes relationships with other insects or plants.
DORYLINAE	Worldwide, mainly tropical.	Ground or ground surface. Include army and driver ants with apterous queens. Predators. Includes Cerapachyinae, Ecitoninae, Leptanilloidinae, Aenictinae, Aenictogitoninae as junior synonyms.
FORMICINAE	Worldwide, mainly temperate.	Wide range of nesting habits, behavior and food sources. Includes parasitisms and relationships with other insects or plants.
ECTATOMMINAE	Australia, Asia, New World.	Mainly on the ground, some arboreal. Nesting biology and food sources varied. Includes Heteroponerinae.
MYRMICINAE	Worldwide.	Wide range of nesting habits, behavior and food sources. Includes parasitisms and relationships with other insects or plants.
MYRMECIINAE	Australia.	Ground. Predators.
PSEUDOMYRMECINAE	Worldwide, mainly tropical.	Mainly arboreal, omnivorous, some associated with plants.

Gene-based phylogenies

Recent molecular analyses (Moreau *et al.* 2006; Brady *et al.* 2006; Rabeling *et al.* 2008; Schmidt 2013; Moreau and Bell 2013) coincide on several important points, such as the monophyly of the Formicidae, most subfamilies, and some suprafamily clades. If we take as reference the phylogeny of Moreau and Bell (2013), Leptanillinae is the sister group of the rest of the ants (and not Martialinae as in Rabeling *et al.* 2008), forming a trichotomy between Martialinae, the poneroid subfamilies (Amblyoponinae, Proceratiinae, Paraponerinae, Agroecomyrmecinae, and Ponerinae), and the formicoid group (the rest of the subfamilies). In addition to the support of the formicoid group, the study supports the dorylomorph subclade (*sensu* Bolton 2003), and the subclades Dolichoderinae + Aneuretinae, and Formicinae + (Ectatomminae *s.l.* + Myrmicinae). Another important difference in the proposal of Moreau and Bell (2013) and the previous ones is that the sister group of Myrmicinae is Ectatomminae (and not Formicinae), a topic already suggested by William Brown earlier in Brown (1954). Since support for the group made up of the poneroid subfamilies is low, Ward (2014) suggests these subfamilies are an unresolved polytomy.

Borowiec *et al.* (2019) explore the phylogeny of ant subfamilies based on the sequences of 11 nuclear genes and

exploring the heterogeneity in the composition of sequences between taxa (emphasis on non-formicomorphs), as well as the divergence times of the clades. These authors conclude that the ants arose towards the Albian / Aptian [Lower Cretaceous, about 124/104 million years ago] and that Martialinae + Leptanillinae are the sister group of the rest of the ants. A third important conclusion is the monophyly of the poneroid group with the relationships ((Apomyrminae + Amblyoponinae) + Proceratiinae) + ((Agroecomyrmecinae + Paraponerinae) + Ponerinae) (Fig. 2).

In addition to their study of ant phylogeny, Moreau and Bell (2013) explore the evolution of ants under the hypothesis of the tropics as a “museum” (lineage reservoir site) or “cradle” (site of origin of many groups) of the diversification of these insects. These authors establish an age of between 158 and 139 million years ago and 10 periods of significant changes in the diversification of ants. The authors conclude that the Neotropical region has been both a “cradle and a museum”.

Internal explorations in the subfamilies have been carried out in Amblyoponinae (Ward and Fisher 2016), Dolichoderinae (Ward *et al.* 2010; Boudinot *et al.* 2016), Dorylinae (Brady *et al.* 2014; Borowiec 2019), Formicinae (Blaimer *et al.* 2015), Myrmicinae (Ward *et al.* 2015), Ponerinae (Schmidt 2013), and Pseudomyrmecinae (Ward and Downie 2005).




Figure 2. Phylogenetic relationship of the ant subfamilies, based on the 11 nuclear genes, modified from Borowiec *et al.* (2019). Photographs taken from AntWeb (2021), full credits in the acknowledgements section.

Classification

Table 1 lists the subfamilies of fossil and living ants according to the most recent proposals. Currently, the Neotropical region comprises 12 subfamilies, 137 genera, and about 3,100 described species (Table 2). Figure 3 shows the diversity of ant genera and species in the world by subfamily, and for the Neotropical region to January of 2021. Figure 4 shows the comparison of ant genus and species counts for the Neotropical region, discriminated by country.

Ant diversity and hyperdiverse insect families

Insects are considered the most diverse group of eukaryotes on the planet, with more than a million species described in some 1,200 families and 28 living orders. Certainly, this diversity is concentrated in the holometabolous orders, and within these in several subgroups, such as Parasitoida in Hymenoptera, Polyphaga in Coleoptera, Pupipara in Diptera, or Macrolepidoptera in Lepidoptera. Furthermore, of the more than 1,000 living insect families, only 19 can be considered hyperdiverse

with an arbitrary estimate of 10,000 or more species (Fig. 5). Except for Miridae (Hemiptera), the rest belong to the four hyperdiverse insect orders, namely, Hymenoptera, Coleoptera, Diptera, and Lepidoptera. Formicidae occupies the 13th position with its almost 15,000 described species (Fig. 5). It probably does not rise much more on this scale, because in tremendously diverse groups such as moths there are not as many taxonomists working on their descriptions as there are in ants. If some estimates suggest more than 20,000 species of ants in the world, surely these must be more numerous in groups than the poorly known as Geometridae, Noctuidae, Chrysomelidae, or Staphylinidae. Another issue is reasonable identification capacity at least to the genus level. In this sense and at least for America, only for the families Formicidae and Braconidae are there tools (such as identification manuals, keys, and catalogs) that allow the user to be secure (or at least acceptable) in identifications of subfamilies and genera. In the case of ants, the sources of information for arriving at a species names are greater, with more than 70 % of the known species being named with some security.

Table 2. Synopsis of the subfamilies, tribes and genera of extant Neotropical ants. Number of species refer only to Neotropics. In front of each subfamily the number of genera is provided.

SUBFAMILY AGROECOMYRMECINAE (1)	SUBFAMILY ECTATOMMINAE (5)
<i>Tatuidris</i> Brown and Kempf, 1968 - 1, Central America to Brazil	Tribe Ectatommini
SUBFAMILY AMBLYOPONINAE (2)	<i>Ectatomma</i> F. Smith, 1858 - 14, Neotropical
<i>Fulakora</i> Mann, 1919 - 12, Neotropical	<i>Gnamptogenys</i> Roger, 1863 - 90, Neotropical
<i>Prionopelta</i> Mayr, 1866 - 5, Neotropical	<i>Typhlomyrmex</i> Mayr, 1862 - 7, Neotropical
SUBFAMILY DOLICHODERINAE (12)	Tribe Heteroponerini
Tribe Bothryomyrmecini	<i>Acanthoponera</i> Mayr, 1882 - 4, Neotropical
<i>Bothriomyrmex</i> Emery, 1869 - 1, Honduras, Costa Rica	<i>Heteroponera</i> Mayr, 1877 - 13, Neotropical
Tribe Dolichoderini	SUBFAMILY FORMICINAE (14)
<i>Dolichoderus</i> Lund, 1831 - 54, Neotropical	Tribe Camponotini
Tribe Leptomyrmecini	<i>Camponotus</i> Mayr, 1861 - 450, Neotropical
<i>Anillidris</i> Santschi, 1936 - 1, Argentina, Brazil	<i>Colobopsis</i> Mayr, 1861 - 8, Mexico and Bahamas to Costa Rica
<i>Azteca</i> Forel, 1878 - 70, Neotropical	Tribe Gigantiopini
<i>Dorymyrmex</i> Mayr, 1866 - 41, neotropical	<i>Gigantiops</i> Roger, 1863 - 1, South America
<i>Forelius</i> Emery, 1888 - 18, neotropical	Tribe Formicini
<i>Gracilidris</i> Wild and Cuezzo, 2006 - 1, Colombia, Brazil, Paraguay	<i>Formica</i> L., 1758 - 1, Mexico to Guatemala
<i>Leptomyrmex</i> Mayr, 1862 - 1, Brazil	Tribe Lasiini
<i>Linepithema</i> Mayr, 1866 - 19, Neotropical	<i>Nylanderia</i> Emery, 1906 - 23, Neotropical
Tribe Tapinomini	<i>Paratrechina</i> Motschulsky, 1893 - 1, Neotropical
<i>Liometopum</i> Mayr, 1861 - 1, Mexico to Guatemala	<i>Zatania</i> LaPolla, Kallal and Brady, 2012 - 5, Central America and West Indies
<i>Tapinoma</i> Foerster, 1850 - 20, Neotropical	Tribe Melophorini
<i>Technomyrmex</i> Mayr, 1870 - 2, Costa Rica, Panama, Colombia	<i>Lasiophanes</i> Emery, 1895 - 5, Chile, Argentina
SUBFAMILY DORYLINE (12)	Tribe Myrmelachistini
<i>Acanthostichus</i> Mayr, 1887 - 19, Neotropical	<i>Brachymyrmex</i> Mayr, 1868 - 53, Neotropical
<i>Cheliomyrmex</i> Mayr, 1870 - 4, Neotropical	<i>Myrmelachista</i> Roger, 1863 - 47, Neotropical
<i>Cylindromyrmex</i> Mayr, 1870 - 10, Neotropical	Tribe Plagiolepidini
<i>Eciton</i> Latreille, 1804 - 12, Neotropical	<i>Acropyga</i> Roger, 1862 - 27, Neotropical
<i>Labidus</i> Jurine, 1807 - 7, Neotropical	<i>Anoplolepis</i> Santschi, 1914* - 1, Introduced in Mexico and Chile
<i>Leptanilloides</i> Mann, 1923 - 18, Neotropical	<i>Plagiolepis</i> Mayr, 1861* - 1, Introduced in Mexico, Bahamas and West Indies
<i>Neocerapachys</i> Borowiec, 2016 - 2, Panama and South America	<i>Prenolepis</i> Mayr, 1861 - 1, Mexico, Jamaica, Panama
<i>Nomamyrmex</i> Borgmeier, 1936 - 2, Neotropical	
<i>Neivamyrmex</i> Borgmeier, 1955 - 103, Neotropical	
<i>Ooceraea</i> Roger, 1862* - 1, introduced in West Indies	
<i>Sphinctomyrmex</i> Mayr, 1866 - 3, Brazil	
<i>Syscia</i> Roger, 1861 - 34, Central America, Colombia, Brazil	
	SUBFAMILY MARTIALINAE (1)
	<i>Martialis</i> Rabeling and Verhaag, 2008 - 1, Brazil

continue...

... continue

SUBFAMILY MYRMICINAE (67)

Tribe Pogonomyrmecini (3)

Hylomyrma Forel, 1912 - 14, Neotropical

Patagonomyrmex Johnson and Moreau, 2016 - 3, Argentina and Chile

Pogonomyrmex Mayr, 1868 - 31, Neotropical

Tribe Stenammini (2)

Aphaenogaster Mayr, 1863 - 4, Central America and Colombia

Stenamma Westwood, 1839 - 40, Central America to Ecuador

Tribe Crematogastrini (9)

Cardiocondyla Emery, 1869 - 2, Neotropical

Carebara Westwood, 1840 - 30, Neotropical

Crematogaster Lund, 1831- 80, Neotropical

Nesomyrmex Wheeler, 1910 - 86, Neotropical

Perissomyrmex M.R. Smith, 1947 - 1, Central America

Temnothorax Mayr, 1861 - 55, Central America, West Indies, Northern South America

Tetramorium Mayr, 1855 - 4, Neotropical

Trichomyrmex Mayr, 1865 - 1, Neotropical

Xenomyrmex Forel, 1884 - 4, Neotropical

Tribe Attini (41)

Acanthognathus Mayr, 1887 - 6, Neotropical

Acromyrmex Mayr, 1865 - 25, Neotropical

Allomerus Mayr, 1878 - 8, Neotropical

Amoimyrnex Cristiano Cristiano, Cardoso and Sandoval-Gómez, 2020 - 3, southern South America

Apterostigma Mayr, 1865 - 40, Neotropical

Atta Fabricius, 1804 -14, Neotropical

Basiceros Schulz, 1906 - 7, Neotropical

Blepharidatta Wheeler, 1915 - 5, South America

Cephalotes Latreille, 1802 -160, Neotropical

Cyatta Sosa-Calvo, Schultz, Brandão, Klingenberg, Feitosa, Rabeling, Bacci, Lopes and Vasconcelos, 2013 - 1, Bolivia, Brazil

Cyphomyrmex Mayr, 1862 - 21, Neotropical

Daceton Perty, 1833 - 2, Neotropical

Diaphoromyrma Fernández, Delabie and Nascimento, 2009 - 1, Brazil

Eurhopalothrix Brown and Kempf, 1961 - 11, Neotropical

Kalathomyrmex Klingenberg and Brandão, 2009 -1, South America

Lachnomyrmex Wheeler, 1910 - 16, Neotropical

Lenomyrmex Fernández and Palacio, 1999 - 6, Costa Rica to Ecuador

Mycetagroicus Brandão and Mayhé-Nunes, 2001 - 4, Brazil, Bolivia

Mycetarotes Emery, 1913 - 4, Colombia, Venezuela, Ecuador, Brazil, Paraguay

Mycetophylax Emery, 1913 - 22, Neotropical

Mycetosoritis Wheeler, 1907 - 6, Mexico to Nicaragua

Mycoceropurus Forel, 1893 - 4, Neotropical

Mycetomoellerius Solomon, Rabeling, Sosa-Calvo and Schultz, 2019 - 29, Neotropical

Myrmicocrypta F. Smith, 1860 - 24, Neotropical

Ochetomyrmex Mayr, 1877 - 2, South America

Octostruma Forel, 1912 - 34, Neotropical

Paramycetophylax Kusnezov, 1956 - 1, Brazil and Argentina

Paratrachymyrnex Solomon, Rabeling, Sosa-Calvo and Schultz, 2019 - 9, neotropical

Phalacromyrmex Kempf, 1960 - 1, Brazil

Pheidole Westwood, 1839 - 463, Neotropical

Procryptocerus Emery, 1887 - 40, Neotropical

Protalaridris Brown, 1980 - 7, Colombia, Venezuela, Ecuador

Pseudoatta Gallardo, 1916 - 1, Argentina

Rhopalothrix Mayr, 1870 - 8, Neotropical

Sericomyrmex Mayr, 1865 - 19, Neotropical

Strumigenys F. Smith, 1860 -186, Neotropical

Talaridris Weber, 1941 - 1, Colombia, Trinidad, Brazil, Guiana

Trachymyrnex Forel, 1893 - 6, Mexico to Guatemala

Tranopelta Mayr, 1866 - 2, Neotropical

Wasmannia Forel, 1893 - 10, Neotropical

Xerolitor Sosa-Calvo, Schultz, Ješovnik, Dahan and Rabeling, 2018 - 1, Bolivia, Brazil

Tribe Solenopsidini (12)

Adelomyrmex Emery, 1897 - 16, Neotropical

Bariamyrmex Lattke, 1990 - 1, Costa Rica, Venezuela

Cryptomyrmex Fernández, 2003 - 2, Guyana, Brazil, Paraguay

Kempfidris Fernández, Lattke and Feitosa, 2014 - 1, South America

Megalomyrmex Forel, 1885 - 38, Neotropical

Monomorium Mayr, 1855 - 22, Neotropical

Oxyepoecus Santschi, 1926 -16, South America

Rogeria Emery, 1915 - 21, Neotropical

Solenopsis Westwood, 1840 - 115, Neotropical

Stegomyrmex Emery, 1912 - 2, Panama and South America

Syllophopsis Santschi, 1915* - 1, introduced in Puerto Rico, Lesser Antilles

Tropidomyrmex Silva, Feitosa, Brandão and Diniz, 2009 - 1, Brazil

SUBFAMILY PARAPONERINAE (1)

Paraponera F. Smith, 1858 - 1, Neotropical

SUBFAMILY PONERINAE (17)

Tribe Ponerini

Anochetus Mayr, 1861- 25, Neotropical

Belonopelta Mayr, 1870 - 2, Mexico to Colombia

Centromyrmex Mayr, 1866 - 3, Neotropical

Cryptopone Emery, 1893 - 5, Neotropical

Dinoponera Roger, 1861 - 9, South America

Hypoponera Santschi, 1938 - 35, Neotropical

Leptogenys Roger, 1861- 80, Neotropical

Mayaponera Schmidt and Shattuck, 2014 - 6, Neotropical

Neoponera Emery, 1901 - 76, Neotropical

Odontomachus Latreille, 1804 - 24, Neotropical

Ponera Latreille, 1804 - 2 Mexico to Guatemala

Pachycondyla F. Smith, 1858 - 17, Neotropical

Pseudoponera Emery, 1920 - 4, Neotropical

Rasopone Schmidt and Shattuck, 2014 - 25, Neotropical

Simopelta Mann, 1922 - 21, Neotropical

Thaumatomyrmex Mayr, 1887 - 12, Neotropical

Tribe Platythyreini

Platythyrea Roger, 1863 - 8, Neotropical

SUBFAMILY PROCRERATIINAE (3)

Tribe Proceratiini

Discothyrea Roger, 1863 - 10, Neotropical

Proceratium Roger, 1863 - 16, Neotropical

Tribe Probolomyrmecini

Probolomyrmex Mayr, 1901 - 8, Neotropical

SUBFAMILY PSEUDOMYRMECINAE (2)

Myrcidris Ward, 1990 - 2, Colombia, Guyana, Brazil

Pseudomyrmex Lund, 1831 - 150, Neotropical




Figure 3. A. Number of ant genera in Neotropical region compared to the rest of the world. B. Number of ant species in Neotropical region compared to the rest of the world. Data updated to January 2021 (Bolton 2021).

Systematics

The Neotropical region is probably the richest realm for ants (Fisher 2010). To date, it comprises 137 genera and around 3,100 species, 38 % of the genera, and 39 % of the world's species (Fig. 3). It is also the region with the highest number of endemics (Fernández 2003a). Part of the explanation is that the Neotropics has been the “cradle and museum” of the myrmecofauna (Moreau and Bell 2013); it is region that generates clades and preserves them. Within the Neotropical region, South America has had a complex geological and

climatic history throughout the ages, especially since the Cenozoic; the subcontinent is a sum of cratons, each of them with a history that speaks of diversity of ecosystems and landscapes. On the other hand, episodes of connection and isolation with other continental masses have enriched the groups of ants that dominate South American landscapes, especially since the mid-Cenozoic (LaPolla *et al.* 2013). If we add to this the existence of about nine genera yet to be described (R. Feitosa and J Sosa-Calvo pers. comm.; FF personal observation), along with many species, in the near future the Neotropics will reaffirm itself as the metropolis of the ants.




Figure 4. Comparison of ant genera (A) and species (B) count in the Neotropical region (discriminated by country). Brazil has the highest diversity with ~80 % of the genera and ~40 % of the species present in the Neotropical region, followed by Colombia (~73 % genera and ~29 % species) and Mexico (~70 % genera and ~28 % species).

The panorama for the taxonomy of ants is relatively acceptable, especially when compared to other megadiverse groups such as the Scarabaeidae, Staphylinidae, Tachinidae, Noctuidae, or Chrysomelidae. Virtually all genera of ants can be identified from workers, and this scenario is valid even for males (Boudinot 2019). Likewise, there are keys, monographs, or synopses of most genera, leaving only a few orphaned groups (Table 3). This vision can be summarized with the key groups like this:

1) *Hypoponera*. Small and homogeneous ants, very difficult to delimit species. Shawn Dash's unpublished doctoral thesis suggests (Dash 2011), however, that the genus is workable with good use of morphometry and other tools. Currently, Amanda Dias and Rodrigo Feitosa (Federal University of Paraná) are developing a project on the systematics and phylogeny of the Neotropical *Hypoponera*.

2) *Camponotus*. After *Pheidole* this is the most diverse genus in the world and in the Neotropics, with about 450 species in this region. As with *Pheidole*, the taxonomy of *Camponotus* is based on major workers, making it difficult to put a species name on the many smaller workers commonly collected in the field. It is a genus that has occupied about 30 years of work of William P. Mackay (University of Texas at El Paso), who recently published a review of the fauna of Colombia (Mackay 2019b). Mackay (2019a) provides a synopsis of the subgenera of *Camponotus* and review the subgenus *Camponotus*.

3) *Pheidole*. Probably the largest and most complex genus of ants in the world has more than 700 species in the Neotropics. Wilson's (2003) review and Longino (2009a) update are two great steps for clarifying the group's taxonomy. However, secure identification is highly dependent on major workers (as in *Camponotus*), which creates a problem with the numerous museum exhibits that only have minor, with no direct link to major workers. Longino (2009b) presents an interactive key that can allow the identification of species with minor workers. Regional synopses are currently underway for Central America (Longino, J. T.), Colombia (Guerrero, R. J. and colleagues), and Brazil (Feitosa, R. and colleagues).

4) *Crematogaster*. Very common ants and difficult to identify due to the problems separating intra- from interspecific variation. The internal phylogeny of the group recently explored (Pedraza, L. and colleagues) maintains some species groups, divides others, and creates some new groups. These groups of species can promote regional or local reviews that can clarify the taxonomy of the group. A review of the groups associated with *limata* is underway.

5) *Solenopsis*. Reviews by Trager (1991), Pacheco and Mackay (2013), and Pitts *et al.* (2018) collectively cover the entire *Solenopsis* fauna of the New World. However, most species belong to this bothersome group and species identification without females is very difficult. The differences between workers are very subtle with the drawback of intra-specific or even intranidal variation.

6) *Carebara*. This is a complex genus, comprising monomorphic, dimorphic, polymorphic, and intercaste species. Monomorphic species (*Carebara* in the old sense) are probably dimorphic since the soldiers have not been able to be associated with field collections. The identification of species with only minor workers (*Carebara lignata* group) is practically impossible due to traditional morphology. One reason is that these workers are very small and many external morphological attributes have been removed; *C. minuta* could be the smallest ant in the world, at just 0.96 mm total length.

7) *Azteca*. This is perhaps the greatest of all challenges since identification with only workers is practically impossible. Except for specific groups (such as those associated with *Cecropia*) it is necessary to have a series of nests with the queen included.

Some genera orphaned in the past will soon have their taxonomy resolved; for example, for *Acromyrmex* (Christian Rabeling and colleagues, Arizona State University), *Anochetus* (Itanna Fernandes and colleagues, INPA, Brazil), *Aphaenogaster* (Jack Longino, The University of Utah), *Atta* (Corina Barrera, Universidad Estadual Paulista - UNESP), *Basiceros* (Beto Brandão and Rodrigo Feitosa, Brazil), *Hypoponera* (Amanda Dias and Rodrigo Feitosa, Federal




Figure 5. The most hyperdiverse insect families in the world places ants (red silhouette) occupying the 13th position. Free silhouettes images obtained from <http://phylopic.org/>.

University of Paraná), *Myrmelachista* (Rodolfo Probst, The University of Utah), *Myrmecocrypta* (Jeffrey Sossa-Calvo and Ted Schulz, Smithsonian), *Nylanderia* (John LaPolla, Towson University), *Syscia* (Jack Longino, The University of Utah) and *Tapinoma* (Roberto Guerrero, Universidad del Magdalena).

A synopsis of subfamilies, tribes and genera of extant Neotropical ants is presented in Table 2, and the list of the Neotropical ant genera that have been reviewed in the last 25 years, including keys for the Neotropical region or at least for some countries in the region, is presented in Table 3.

Species delimitation

In groups of ants with an elevated number of species (and a generally wide distribution), morphology alone is insufficient to know how many species there are, how to separate them, and how to provide identification keys for non-specialist users. This implies that researchers must use various tools and multiple evidence to find out how diverse the group under consideration is. A clear advantage for this strategy is to have a rich knowledge of evolution, speciation and population dynamics. This information, together with biology and distributional data, offers an overview of the evolutionary biogeography of the group of ants studied. As happens in other taxa, one of these approaches involves the use of genetic information to establish the boundaries between species. This approach to species delimitation tries to determine how species and their populations are grouped in a broader sense than its classification (Carstens *et al.* 2013).

With the development of better analytical methods, increased computational capacity and the proliferation of genomic information, these genetic tools have become increasingly useful. It is important to keep in mind that the primary goal of these methodologies, based on molecular markers, is to identify independent evolutionary lineages and validate them as biodiversity units (Queiroz 2007; Carstens *et al.* 2013). It is from these units that species are discriminated using independent contrasts based on ecological, geographic, physiological, or morphological data (Leavitt *et al.* 2015; Prebus 2020). The main difficulty with this approach is the variability in the resulting delimitations, which, if not contrasted with other types of evidence (or if not available), it is not clear where the boundary should be placed between different species. For example, a *Solenopsis* study in Ecuador (Delsinne *et al.* 2012) shows high variability in some of the species but the marker used, COI (Cytochrome Oxidase Subunit I), was insufficient to resolve the limits of these ants on its own. Furthermore, fine molecular techniques and the abuse of the cryptic species concept can skyrocket the number of species in any group of insects. An additional problem of a practical nature is the near impossibility of identifying species whose delimitation is purely based on genetic markers. A user would have to repeat the same laboratory and analytic protocols to differentiate species, without the guarantee of obtaining the same results, precisely because of the genetic variation between the populations.

At the very least, the examples shown above indicate that in some cases, the separation of species by morphology is conservative, and at least the species richness of a group is not

Table 3. Ant genera with synopsis or keys for Neotropics, or Neotropical countries, in the last 25 years. Monotypic genera omitted.

Subfamily and genera	Reference
SUBFAMILY AMBLYOPONINAE	
<i>Fulakora</i>	Lattke (1991), Lacau and Delabie (2002), Arias-Penna (2008a)
<i>Prionopelta</i>	Ladino and Feitosa (2020)
SUBFAMILY DORYLINAE	
<i>Acanthostichus</i>	Mackay (1996)
<i>Cylindromyrmex</i>	De Andrade (1998)
<i>Eciton</i>	Palacio (1999, 2019)
<i>Labidus</i>	Palacio (1999, 2019)
<i>Leptanilloides</i>	Brandão <i>et al.</i> (1999), Donoso <i>et al.</i> (2006), Borowiec and Longino (2011)
<i>Neocerapachys</i>	Brown (1975)
<i>Nomamyrmex</i>	Palacio (1999, 2019)
<i>Neivamyrmex</i>	Palacio (1999, 2019)
<i>Sphinctomyrmex</i>	Feitosa <i>et al.</i> (2012)
<i>Syscia</i>	Brown (1975), Longino and Branstetter (2021)
SUBFAMILY DOLICHODERINAE	
<i>Dolichoderus</i>	Ortiz and Fernández (2011)
<i>Dorymyrmex</i>	Cuezzo and Guerrero (2012)
<i>Forelius</i>	Cuezzo (2000), Guerrero and Fernández (2008), Guerrero (2021)
<i>Linepithema</i>	Wild (2004, 2007), Escárraga and Guerrero (2016)
<i>Tapinoma</i>	Escárraga <i>et al.</i> (2021), García-Avendaño and Guerrero (2018)
<i>Technomyrmex</i>	Bolton (2007), Fernández and Guerrero (2008)
SUBFAMILY ECTATOMMINAE	
<i>Ectatomma</i>	Fernández (1991), Arias-Penna (2008b)
<i>Gnamptogenys</i>	Lattke <i>et al.</i> (2007, 2008)
<i>Acanthoponera</i>	Arias-Penna and Fernández (2008)
<i>Heteroponera</i>	Arias-Penna and Fernández (2008)
SUBFAMILY FORMICINAE	
<i>Acropyga</i>	LaPolla (2004)
<i>Brachymyrmex</i>	Ortiz and Fernández (2014), Ortiz-Sepulveda <i>et al.</i> (2019)
<i>Myrmelachista</i>	Longino (2006a)
<i>Zatania</i>	LaPolla <i>et al.</i> (2012)
SUBFAMILY MYRMICINAE	
<i>Pogonomyrmex</i>	Fernández and Palacio (1997), Lattke (2006), Cuezzo and Claver (2009)
<i>Patagonomyrmex</i>	Johnson and Moreau (2016)
<i>Aphaenogaster</i>	Mackay and Mackay (2002), Longino and Cover (2004)
<i>Stenamma</i>	Fernández <i>et al.</i> (1996), Branstetter (2013)
<i>Cardiocondyla</i>	Mackay (1995)
<i>Carebara</i>	Fernández (2004a), Fischer <i>et al.</i> (2014)
<i>Crematogaster</i>	Longino (2003)
<i>Acanthognathus</i>	Galvis and Fernández (2009)
<i>Acromyrmex</i>	Fernández <i>et al.</i> (2015)
<i>Allomerus</i>	Fernández (2007a)
<i>Amoimyrmex</i>	Cristiano <i>et al.</i> (2020)
<i>Apterostigma</i>	Lattke (1997)
<i>Blepharidatta</i>	Brandão <i>et al.</i> (2015)
<i>Atta</i>	Fernández <i>et al.</i> (2015)
<i>Basiceros</i>	Probst <i>et al.</i> (2019)
<i>Mycetomoellerius</i>	Mayhé-Nunes and Brandão (2002, 2005, 2007)
<i>Cephalotes</i>	De Andrade and Baroni Urbani (1999), Sandoval-Gómez and Sánchez-Restrepo (2019)
<i>Cyphomyrmex</i>	Snelling and Longino (1992), Mackay and Mackay (2002)
<i>Lachnomyrmex</i>	Feitosa and Brandão (2008)
<i>Lenomyrmex</i>	Fernández and Palacio (1999), Fernández (2001), Delsinne and Fernández (2012)
<i>Mycetophylax</i>	Klingenberg and Brandão (2009), Sosa-Calvo <i>et al.</i> (2017)
<i>Mycocepurus</i>	Mackay <i>et al.</i> (2004)
<i>Myrmicocrypta</i>	Sosa-Calvo and Schultz (2010)
<i>Ochetomyrmex</i>	Fernández (2003b)
<i>Octostruma</i>	Palacio (1997), Longino (2013)

continue...

...continue

Subfamily and genera	Reference
<i>Pheidole</i>	Wilson (2003), Longino (2009a)
<i>Procryptocerus</i>	Longino and Snelling (2002)
<i>Protalaridris</i>	Lattke <i>et al.</i> (2018)
<i>Rhopalothrix</i>	Longino and Boudinot (2013)
<i>Sericomyrmex</i>	Ješovník and Schultz (2017)
<i>Strumigenys</i>	Bolton (2000), Longino (2006b)
<i>Tranopelta</i>	Fernández 2003b
<i>Wasmannia</i>	Longino and Fernández (2007), Cuezzo <i>et al.</i> (2015)
<i>Adelomyrmex</i>	Fernández and Mackay (2003), Fernández (2003c), Longino (2012)
<i>Cryptomyrmex</i>	Fernández (2004b)
<i>Megalomyrmex</i>	Brandão (1990, 2003), Longino (2010), Boudinot <i>et al.</i> (2013)
<i>Monomorium</i>	Fernández (2007b)
<i>Oxyepoecus</i>	Albuquerque and Brandão (2004, 2009), Delsinne <i>et al.</i> (2012)
<i>Rogeria</i>	Kugler (1994), LaPolla and Sosa-Calvo (2006)
<i>Solenopsis</i>	Trager (1991), Pacheco and Mackay (2013), Pitts <i>et al.</i> (2018)
<i>Stegomyrmex</i>	Feitosa <i>et al.</i> (2008)
SUBFAMILY PONERINAE	
<i>Anochetus</i>	González-Campero and Elizalde (2008), Zabala (2008)
<i>Belonopelta</i>	Fernández (2008)
<i>Centromyrmex</i>	Fernández (2008)
<i>Cryptopone</i>	Mackay and Mackay (2010)
<i>Dinoponera</i>	Fernández (2008), Lenhart <i>et al.</i> (2013)
<i>Leptogenys</i>	Lattke (2011)
<i>Neoponera</i>	Wild (2002, 2005), Mackay and Mackay (2010)
<i>Odontomachus</i>	Rodríguez (2008)
<i>Pachycondyla (sensu lato)</i>	Wild (2002), Mackay and Mackay (2010)
<i>Pseudoponera</i>	Mackay and Mackay (2010)
<i>Rasopone</i>	Mackay and Mackay (2010), Longino and Branstetter (2020)
<i>Simopelta</i>	Mackay and Mackay (2008), Fernandes <i>et al.</i> (2015)
<i>Thaumatomyrmex</i>	Fernández (2008)
<i>Platythyrea</i>	De Andrade (2004), Fernández (2008)
SUBFAMILY PROCRATIINAE	
<i>Discothyrea</i>	Sosa-Calvo and Longino (2008), Chaul (2020)
<i>Probolomyrmex</i>	Oliveira and Feitosa (2019)
<i>Proceratium</i>	Baroni Urbani and De Andrade (2003), Sosa-Calvo and Longino (2008), Fernandes <i>et al.</i> (2019)
SUBFAMILY PSEUDOMYRMECINAE	
<i>Pseudomyrmex</i>	Ward (1989, 1999, 2017)

being artificially maximized (see also Andersen *et al.* 2016). Of course, the opposite can be expected, accepting broad species criteria where morphological and genetic diversity is considered as simply being variants of a polymorphic species. After all, in insects, it has long been accepted that there is great phenotypic variation due to historical and ecological conditions of the various lineages or populations of many widely distributed species.

The ant landscape in the Neotropics is mixed. For 70 % - 75 % of the genera there are reviews, monographs, synopses, or taxonomic notes that allow reasonable sorting and identification of species. The AntWeb (2021) digital portal also offers high-quality photographs of most types of collections usually inaccessible to Latin American biologists. This digital page as well as that of the AntCat (Bolton 2021) also includes the Barry Bolton catalog and access to many digitized documents with original descriptions of species from Linnaeus. If the monotypic genera and those currently being reviewed (*Hypoponera*, *Nylanderia*, *Crematogaster*, *Myrmicocrypta*, and *Tapinoma*) are added to this information, the panorama is encouraging.

The nature or complexity of each group is what defines what types of tools should be used to delimit ant species. In genera as diverse as *Strumigenys* or *Pheidole*, morphology and/or morphometry may suffice to separate and name specimens. It is more a matter of precision and patience to name a specimen or establish if it is a new taxon. In genera such as *Solenopsis* or *Azteca* it is clear that many data sources must be used: biochemical (as cuticle chemistry), genetic (several markers), and natural history (ecology, behavior).

It is important to remember that the definition of a species represents the conclusion of a scientific investigation based on the knowledge and tools available at the time of the study. This definition may change with the development of new techniques, even in groups that are currently considered well-known. The promises of barcoding lagged far behind in the promise of replacing traditional taxonomy. Only the use of different criteria and tools can give a reasonable idea for the separation of evolutionary lineages and consequently the proposition and recognition of species. The integration of various information resources together with progress in data analysis will allow a better approach to the diversity of ants.

New tools and techniques useful for delimiting species continue to appear. For example, Ješovník *et al.* (2017) used genomic data based on ultra-conserved elements (UCE) to reconstruct evolutionary histories and infer species limits in *Sericomyrmex*. Near infrared spectroscopy allows the analysis of hydrocarbons and other chemical compounds present on the surface of ants, facilitating the correct identification of cryptic species (Kinzner *et al.* 2015). The recent use of X-ray microtomography in the taxonomy of ants (Sarnat *et al.* 2016; Fischer *et al.* 2016; Hita García *et al.* 2017) is very promising, although some challenges remain to be achieved (Hita García *et al.* 2017). This technique allows detailed evaluations of the external and internal morphology of the specimen without causing damage. Volumetric models of the specimen are obtained that can be visualized and manipulated in three dimensions with resolution from micrometer to nanometer. Virtual reconstructions can be rotated, sectioned, and measured to gain a comprehensive understanding of the anatomy and morphology of the studied specimen (Hita García *et al.* 2017).

Recently, Prebus (2020) used an integrated approach for the delimitation of species within the *salvini* group of the genus *Tennothorax*; this approach included molecular data of the UCE type, sequences of the mitochondrial genome, morphometric data, and geographical distributions analyzed under novel strategies that allow discrimination between possible population structure and reproductive isolation in the possible resulting species (species delimitation hypothesis). The Prebus (2020) results are based on a methodological scheme very rich in procedures and extremely useful for those genera with difficulties in the delimitation of the species (*e.g.*, *Hypoponera*). But again, it is crucial to keep in mind that these innovative methods do not replace, but rather complement, other traditional methods and procedures (Schlick-Steiner *et al.* 2010). Let us remember that “the taxonomist’s nightmare is the evolutionary biologist’s delight” (Dejaco *et al.* 2016).

Perspectives and challenges

Whitfield and Kjer (2008) point out that it is possible that, due to ancient rapid radiation from many insect lineages, the morphological and/or molecular information has been erased or misrepresented in such a way that we lose clues for reconstructing the phylogenies of basal groups (stem groups). If ants originated long ago (perhaps towards the Jurassic-Cretaceous transition) in a short period of explosive diversification, clarifying the relationships of their basal groups will be difficult. If so, this would explain the difficulty in finding good support for the group closest to the root and sister group of the rest of the ants (Martialinae? Leptanilinae? The clade formed by both subfamilies? See Borowiec *et al.* 2019). However, the recent development of phylogenomics has increased group separation in the Insecta (Misof *et al.* 2014) and Hymenoptera (Johnson *et al.* 2013). Certainly, this new resource for information, together with computational advances, will make it possible to resolve various current uncertainties in the Formicidae.

One of the obstacles to the knowledge and protection of biodiversity is precisely knowing what we have and where. The enormous diversity of insects, which even in the most conservative estimates approximates 3 million species, shows that there are a large number of species to be described. Added

to this problem is the progressive disappearance of qualified taxonomists who cover known groups. Many groups of insects remain virtually unknown or were reviewed decades ago. Even in economic interest groups (such as parasitoid moths or wasps), there is an urgent need for specialists. This has led some biologists to propose faster ways to describe the world’s entomofauna. Some proposals are controversial, such as describing species without control specimens (*i.e.*, no types; for example, see Marshall and Evenhuis 2015). Others are very recent and will surely generate many controversies. An example of this is the description of species with gene characters that accompany focus stacking photographs, biology, and distribution data [Hoenle *et al.* (2020) for *Odontomachus davidsoni*]. In this sense, the morphological information is reduced to one image or a few images. Can this work for other ants? As in ants, there are many modern studies in DNA sequencing; in principle, it is not a problem to characterize species by specific sequences of chosen genes. However, many ants form or appear to form complexes in which one image is not sufficient to delimit or separate species. The presence of castes, the reproductive male or female may also be required. In other cases, there is morphological variation to such a degree that many images would be required, some in scanning electron microscopy (SEM), which in the long run would go against quick and simple descriptions. On the other hand, in ants like *Solenopsis*, genetic variation is high, so much so that there are conflicts between morphology and genes in defining a supposed species. Integrative taxonomy is a strategy that has the strength to accomplish a revision of a given taxon, although it requires time and resources that justify, in part, the so-called “taxonomic impediment”.

The impediment for specimen exportation

A final problem in ant systematics, as well as that of any other group of organisms, is more one of government regulation than of biology. The majority of countries in the Neotropical region have created and promoted laws and regulations of such complexity that it is practically impossible to collect and exchange specimens between countries and, therefore, between colleagues (Fernández 2011; Acosta and Pérez-González 2019). When cooperation between institutions and countries is the best strategy to face the “taxonomic impediment” and the gradual lack of resources in biodiversity studies, the governments themselves have created a tangle of regulations that discourages even the most persevering. This has forced some biologists to move between relative “illegality” and civil disobedience, and others to carry out local studies where the need to resort to sources of specimens outside the country is minimized.

Challenges and tasks

Although recent years have witnessed great advances in ant phylogeny and systematics, there are still several challenging tasks to achieve a better understanding of the diversity of Neotropical ants.

1) What is the sister group of ants? Recently, it has been proposed that the sphecid wasps (broad sense) and bees (Anthophila), that is the Apoidea superfamily, is the sister clade of Formicidae. Boudinot *et al.* (2020) propose the clade Formicapoidina for this association. This has several consequences in terms of visualizing the ancestry of both groups that probably

lived towards the end of the Jurassic, and exploring the morphological changes has led to apoids and ants since the Cretaceous. It is interesting that *Cariridris bipetiolata* has been considered an Ampulicidae and not as an ant as proposed since this hymenopteran is from the Lower Cretaceous of Brazil and this family is at the base of Apoidea, not far from the supposed ancestor of Apoidea + Formicidae. On the other hand, if Apoidea is a superfamily, ants must be upgraded to a superfamily, like Formicoidea, with only one family, depending on the status of the fossil taxon Armaniidae * / Armaniinae *. Boudinot *et al.* (2020) propose new names, clades, and diagnosis within the Formicoidea.

2) Where is the root of the ant tree? The latest proposals have varied between Leptanillinae and Martialinae. The most recent phylogeny by Borowiec *et al.* (2019) propose both subfamilies as the sister group of the rest of ants (Poneroid plus Formicoid clades). It is important to highlight the observation of Borowiec *et al.* (2019) in the sense that it is not about increasing samples and genes, but about performing analyses by changing external groups and analytical methods. For example, several of the outer groups and Leptanillinae are richer in DNA nucleotide adenine (A) and thymine T pairs, while Martialinae and the rest of ants are richer in DNA nucleotide guanine (G) and cytosine (C) pairs.

3) Although the Poneroid clade appears to be monophyletic, some of its internal relationships are not entirely clear, especially the position of Proceratiinae. Within the Proceratiinae the genus *Proceratium* appears to be paraphyletic, and within the Ponerinae the genera *Mesoponera*, *Euponera*, and *Cryptopone* are not monophyletic (Borowiec *et al.* 2019). Likewise, the status of *Mayaponera* and *Rasopone* merits further study (Longino, pers. comm.).

4) In Dorylinae, the phylogenetic relationships and delimitation in the genera around the old concept of "Cerapachyinae" still needs to be explored in greater depth (Borowiec 2016, 2019). For the Neotropical region, the genus *Eciton*, one of the most studied in natural history needs a thorough revision to resolve the status of several of its subspecies.

5) The subfamily Myrmicinae is the most diverse and therefore it is where most problems have arisen as barriers to a clear picture of its phylogeny and diversity. The phylogeny of Ward *et al.* (2015) is a huge step forward, but some groups inevitably require further study. One of them is the internal phylogeny of the Solenopsidini tribe, especially the genera that have been described around *Monomorium*. The resurrections of *Epelysidris* and *Trichomyrmex* in Ward *et al.* (2015) and *Chelaner* (Sparks *et al.* 2019) clearly show the artificiality of *Monomorium* and the status of *Nothidris*, *Antichthonidris* (both from southern South America); and *Monomorium delabieei* Fernández, 2007 (Northeast of Brazil) should be studied, which surely corresponds to more than one genus.

6) The hyperdiverse genera: Under this category are genera such as *Pheidole* (Wilson 2003) and *Camponotus* in the Neotropical region. *Pheidole* can have between 600 - 700 species or more in the Neotropical region, taking into account the large number of new species awaiting study and description. In addition to being diverse, both genera are very common at the regional level (*Camponotus* even approaches the level of the páramo in the Andes cordillera); therefore, they represent a great challenge in species identification. Taxonomic keys in Wilson (2003) are based on major workers, but since many field samples and museum collections contain mainly specimens with only minor workers, we need other tools to

identify such material. John Longino's *Pheidole* interactive taxonomic keys (Longino 2009b) are a great help, although they require a lot of detailed measurements and time, which a non-specialist user might consider a handicap. A similar situation occurs in *Camponotus*, where the published reviews and/or keys (e.g., Mackay and Mackay 2019) are mainly based on major workers. For *Pheidole* and *Camponotus*, new and varied identification tools are needed, without forgetting the elaboration of traditional and illustrated keys based on minor workers, at least on the local scale of the country (as *Pheidole* Project in Colombia).

7) *Atta* genus group. This is a group of great importance at the level of ecology and agricultural economics, but especially for its long and complex evolutionary history with mutualistic fungi, parasites, and bacteria. These ants comprised the *Atta* genus group in Attini tribe (*sensu* Ward *et al.* 2015). Many studies have been conducted on coevolution among the group, in its associated fungi, and between both. Also, and as a consequence of these studies, some genera have been redefined and described, especially *Mycetosoritis*, *Cyphomyrmex* and *Trachymyrmex*. A consequent challenge with these changes is to offer keys or tools that allow the general user to correctly identify the genera of these ants, especially the leaf-cutter ants (*Acromyrmex*, *Amoimyrmex* and *Atta*).

8) Genera in need of revision. In the systematics section, genera were mentioned in need of revision but with ongoing projects in phylogeny and systematics. Many others are equally important but we do not know of global or local revision plans. Among these, we highlight *Discothyrea*, *Odontomachus*, *Platythyrea*, *Neivamyrmex*, *Rogeria*, *Nesomyrmex* and *Xenomyrmex*.

9) Critical areas. Ants in the Neotropical region has been unevenly studied, with well-inventoried countries like Costa Rica in contrast to others like Bolivia with modern studies practically non-existent. In addition to the importance of knowing the biodiversity of the entire Neotropics for biogeographic and conservation studies, the gradual destruction of forests forces an acceleration of research to increase our knowledge of critical and threatened areas, such as the Central American and inter-Andean valleys, the Chocó-Darién moist forests, the foothills of the Andean mountain range, large areas of Bolivia, the Yungas rainforest, northwestern Brazil, portions of the Cerrado savanna, and the Chaco region.

Acknowledgements

We thank our colleague Thibaut Delsinne with whom we initially worked the chapters on which we base part of this contribution. To Consejo Nacional de Investigaciones Científicas y Técnicas de Argentina (Conicet) for the financial support of AFSR. RJJ receives financial support from the Fondo para la Ciencia de la Universidad del Magdalena (Fonciencias) through the project "Taxonomy and distribution of *Pheidole* ants (Formicidae: Myrmicinae) in Colombia". Thanks to the editor of the RCdE, James Montoya-Lerma for the invitation and to editorial team for their assistanship. The English language review was performed by Dr. Thomas Defler. Thanks also to the two anonymous reviewers for their comments. Credits of photographs from www.antweb.org in figure 2 corresponds to: April Nobile (Aneuretinae: *Aneuretus simoni*, CASENT0007014; Myrmecinae: *Myrmecia* sp., CASENT0010660; Pseudomyrmecinae: *Pseudomyrmex gracilis*, CASENT0173760;

Ectatomminae: *Ectatomma edentatum*, CASENT0173376; Formicinae: *Nylanderia fulva*, CASENT0173491; Agroecomyrmecinae: *Tatuidris* sp., CASENT0001968; Paraponerinae: *Paraponera clavata*, CASENT0006789; Ponerinae: *Dinoponera* sp., CASENT0004661; Amblyoponinae: *Fulakora orizabana*, CASENT0172807; Apomyrminae: *Apomyrma stygia*, CASENT0000077; Proceratiinae: *Discothyreareticulata*, CASENT0178697; Leptanillinae: *Protanilla* sp., CASENT0178845), Will Ericson (Dolichoderinae: *Azteca chartifex*, CASENT0249593; Dorylinae: *Eciton burchellii*, CASENT0905935), Zach Lieberman (Myrmicinae: *Acromyrmex ambiguus*, CASENT0922019), and Michael Branstetter (Martinaliae: *Martialis heureka*, CASENT0106181).

Literature cited

- ACOSTA, L. E.; PÉREZ-GONZÁLEZ, A. 2019. ¿Quién va a describir nuestra biodiversidad?: el impedimento taxonómico frente al Protocolo de Nagoya y las normativas vigentes. Revista del Museo Argentino de Ciencias Naturales N. S. 21 (1): 17-27. <http://dx.doi.org/10.22179/REVMACN.21.629>
- AGOSTI, D.; GRIMALDI, D.; CARPENTER, J. M. 1998. Oldest known ant fossils discovered. Nature 391: 447. <https://doi.org/10.1038/35051>
- ALBUQUERQUE, N. L.; BRANDÃO, C. R. F. 2004. A revision of the Neotropical Solenopsidini ant genus *Oxyepoecus* Santschi, 1926 (Hymenoptera: Formicidae: Myrmicinae). 1. The Vezenyii species-group. Papéis Avulsos de Zoologia 44 (4): 55-80. <https://doi.org/10.1590/S0031-10492004000400001>
- ALBUQUERQUE, N. L.; BRANDÃO, C. R. F. 2009. A revision of the Neotropical Solenopsidini ant genus *Oxyepoecus* Santschi, 1926 (Hymenoptera: Formicidae: Myrmicinae). 2. Final. Key for species and revision of the Rastratus species-group. Papéis Avulsos de Zoologia 49 (23): 289-309. <https://doi.org/10.1590/S0031-10492009002300001>
- ANDERSEN, A. N.; HOFFMANN, B. D.; OBERPRIELER, S. 2016. Diversity and biogeography of a species-rich ant fauna of the Australian seasonal tropics. Insect Science 25 (3): 519-526. <https://doi.org/10.1111/1744-7917.12402>
- ANTWEB. 2021. California Academy of Science. <https://www.antweb.org/> [Review date: 21 February 2021].
- ARIAS-PENNA, T. M. 2008a. Subfamilia Amblyoponinae. pp. 41-51. In: Jiménez, E.; Fernández, F.; Arias, T. M.; Lozano-Zambrano, F. H. (Eds.). Sistemática, biogeografía y conservación de las hormigas cazadoras de Colombia. Instituto de Investigación de Recursos Biológicos Alexander von Humboldt. Bogotá, D. C., Colombia. 609 p.
- ARIAS-PENNA, T. M. 2008b. Subfamilia Ectatomminae. pp. 53-107. In: Jiménez, E.; Fernández, F.; Arias, T. M.; Lozano-Zambrano, F. H. (Eds.). Sistemática, biogeografía y conservación de las hormigas cazadoras de Colombia. Instituto de Investigación de Recursos Biológicos Alexander von Humboldt. Bogotá, D. C., Colombia. 609 p.
- ARIAS-PENNA, T. M.; FERNÁNDEZ, F. 2008. Subfamilia Heteroponerinae. pp. 109-117. In: Jiménez, E.; Fernández, F.; Arias, T. M.; Lozano-Zambrano, F. H. (Eds.). Sistemática, biogeografía y conservación de las hormigas cazadoras de Colombia. Instituto de Investigación de Recursos Biológicos Alexander von Humboldt. Bogotá, D. C., Colombia. 609 p.
- BARDEN, P. 2017. Fossil ants (Hymenoptera: Formicidae): Ancient diversity and the rise of modern lineages. Myrmecological News 24: 1-30. https://doi.org/10.25849/myrmecol.news_024:001
- BARDEN, P.; GRIMALDI, D. 2013. A new genus of highly specialized ants in Cretaceous Burmese amber (Hymenoptera: Formicidae). Zootaxa 3681 (4): 405-412. <https://doi.org/10.11646/zootaxa.3681.4.5>
- BARDEN, P.; GRIMALDI, D. 2014. A diverse ant fauna from the mid-Cretaceous of Myanmar (Hymenoptera: Formicidae). PLoS ONE 9 (4): e93627. <https://doi.org/10.1371/journal.pone.0093627>
- BARDEN, P.; HERHOLD, H. W.; GRIMALDI, D. A. 2017. A new genus of hell ants from the Cretaceous (Hymenoptera: Formicidae: Haidomyrmecini) with a novel head structure. Systematic Entomology 42 (4): 837-846. <https://doi.org/10.1111/syen.12253>
- BARONI URBANI, C.; DE ANDRADE, M. L. 2003. The ant genus *Proceratium* in the extant and fossil record (Hymenoptera: Formicidae). Monografie 36. Museo Regionale di Scienze Naturali. Torino, Italy. 492 p.
- BLAIMER, B. B.; BRADY, S. G.; SCHULTZ, T. R.; LLOYD, M. W.; FISHER, B. L.; WARD, P. S. 2015. Phylogenomic methods outperform traditional multi-locus approaches in resolving deep evolutionary history: a case study of formicine ants. BMC Evolutionary Biology 15: 271. <https://doi.org/10.1186/s12862-015-0552-5>
- BOLTON, B. 2000. The ant tribe Dacetini. Memoirs of the American Entomological Institute 65 (1): 1-1028.
- BOLTON, B. 2003. Synopsis and classification of Formicidae. Memoirs of the American Entomological Institute 71: 1-370.
- BOLTON, B. 2007. Taxonomy of the dolichoderine ant genus *Technomyrmex* Mayr (Hymenoptera: Formicidae) based on the worker caste. Contributions of the American Entomological Institute 35 (1): 1-150.
- BOLTON, B. 2021. An online catalog of the ants of the world. Available in: <https://antcat.org>. [Review date: 21 February 2021].
- BOROWIEC, M. L. 2016. Generic revision of the ant subfamily Dorylinae (Hymenoptera, Formicidae). ZooKeys 608: 1-280. <https://doi.org/10.3897/zookeys.608.9427>
- BOROWIEC, M. L. 2019. Convergent evolution of the army ant syndrome and congruence in big-data phylogenetics. Systematic Biology 68 (4): 642-656. <https://doi.org/10.1093/sysbio/syy088>
- BOROWIEC, M. L.; LONGINO, J. T. 2011. Three new species and reassessment of the rare Neotropical ant genus *Leptanilloides* (Hymenoptera, Formicidae, Leptanilloidinae). ZooKeys 133: 19-48. <https://doi.org/10.3897/zookeys.133.1479>
- BOROWIEC, M. L.; RABELING, C.; BRADY, S. G.; FISHER, B. L.; SCHULZ, T. R.; WARD, P. S. 2019. Compositional heterogeneity and outgroup choice influence the internal phylogeny of ants. Molecular Phylogenetics and Evolution 134: 111-121. <https://doi.org/10.1016/j.ympev.2019.01.024>
- BORYSENKO, L. H. 2017. Description of a new genus of primitive ants from Canadian amber, with the study of relationships between stem -and crown- group ants (Hymenoptera: Formicidae). Insecta Mundi 0570: 1-57. <https://journals.flvc.org/mundi/article/view/0570/100602>
- BOUDINOT, B. E. 2015. Contributions to the knowledge of Formicidae (Hymenoptera, Aculeata): a new diagnosis of the family, the first global male-based key to subfamilies, and a treatment of early branching lineages. European Journal of Taxonomy 120: 1-62. <https://doi.org/10.5852/ejt.2015.120>
- BOUDINOT, B. E. 2019. Clave para las subfamilias y géneros basada en machos. pp. 487-499. In: Fernández, F.; Guerrero, R. J.; Delsinne, T. (Eds.). Hormigas de Colombia. Universidad Nacional de Colombia, Bogotá, D. C., Colombia. 1198 p.
- BOUDINOT, B. E.; SUMNICHTY, T. P.; ADAMS, R. M. M. 2013. Central American ants of the genus *Megalomyrmex* Forel (Hymenoptera: Formicidae): six new species and keys to workers and males. Zootaxa 3732 (1): 1-82. <https://doi.org/10.11646/zootaxa.3732.1.1>
- BOUDINOT, B. E.; PROBST, R. S.; BRANDÃO, C. R. F.; FEITOSA, R. M.; WARD, P. S. 2016. Out of the Neotropics: newly discovered relictual species sheds light on the biogeographical history of spider ants (*Leptomyrmex*, Dolichoderinae, Formicidae). Systematic Entomology 41 (3): 658-671. <https://doi.org/10.1111/syen.12181>

- BOUDINOT, B. E.; PERRICHOT, V.; CHAUL, J. C. M. 2020. †*Camelosphecia* gen. nov., lost ant-wasp intermediates from the mid-Cretaceous (Hymenoptera, Formicoidea). *ZooKeys* 1005: 21-55. <https://doi.org/10.3897/zookeys.1005.57629>
- BRADY, S. G.; SCHULTZ, T. R.; FISHER, B. L.; WARD, P. S. 2006. Evaluating alternative hypothesis for the early evolution and diversification of ants. *Proceedings of the National Academy of Sciences* 103 (48): 18172-18177. <https://doi.org/10.1073/pnas.0605858103>
- BRADY, S. G.; FISHER, B. L.; SCHULTZ, T. R.; WARD, P. S. 2014. The rise of army ants and their relatives: diversification of specialized predatory doryline ants. *BMC Evolutionary Biology* 14: 93. <https://doi.org/10.1186/1471-2148-14-93>
- BRANDÃO, C. R. F. 1990. Systematic revision of the Neotropical ant genus *Megalomyrmex* Forel (Hymenoptera: Formicidae: Myrmicinae), with the description of thirteen new species. *Arquivos de Zoologia* 31 (5): 411-481. <https://doi.org/10.11606/issn.2176-7793.v31i5p1-91>
- BRANDÃO, C. R. F. 2003. Further revisionary studies on the ant genus *Megalomyrmex* Forel (Hymenoptera: Formicidae: Myrmicinae: Solenopsidini). *Papéis Avulsos de Zoologia* 43 (8): 145-159. <https://doi.org/10.1590/S0031-10492003000800001>
- BRANDÃO, C. R. F.; DINIZ, J. L. M.; AGOSTI, D.; DELABIE, J. H. C. 1999. Revision of the Neotropical ant subfamily Leptanilloidinae. *Systematic Entomology* 24 (1): 17-36. <https://doi.org/10.1046/j.1365-3113.1999.00064.x>
- BRANDÃO, C. R. F.; FEITOSA, R. M.; DINIZ, J. L. M. 2015. Taxonomic revision of the Neotropical Myrmicinae ant genus *Blepharidatta* Wheeler. *Zootaxa* 4012 (1): 33-56. <https://doi.org/10.11646/zootaxa.4012.1.2>
- BRANSTETTER, M. G. 2013. Revision of the Middle American clade of the ant genus *Stenamma* Westwood (Hymenoptera, Formicidae, Myrmicinae). *ZooKeys* 295: 1-277. <https://doi.org/10.3897/zookeys.295.4905>
- BRANSTETTER, M. G.; DANFORTH, B. N.; PITTS, J. P.; FAIRCLOTH, B. C.; WARD, P. S.; BUFFINGTON, M. L.; GATES, M. W.; KULA, R. R.; BRADY, S. G. 2017. Phylogenomic insights into the evolution of stinging wasps and the origins of ants and bees. *Current Biology* 27 (7): 1019-1025. <https://doi.org/10.1016/j.cub.2017.03.027>
- BROTHERS, D. J. 1999. Phylogeny and evolution of wasps, ants and bees (Hymenoptera, Chrysoidea, Vespoidea and Apoidea). *Zoologica Scripta* 28 (1-2): 233-250. <https://doi.org/10.1046/j.1463-6409.1999.00003.x>
- BROWN, W. L. Jr. 1954. Remarks on the internal phylogeny and subfamily classification of the family Formicidae. *Insectes Sociaux* 1: 21-31. <https://doi.org/10.1007/BF02223148>
- BROWN, W. L. Jr. 1975. Contributions toward a reclassification of the Formicidae. V. Ponerinae, tribes Platythyreini, Cerapachyini, Cylindromyrmecini, Acanthostichini, and Aenictogitini. *Search Agriculture* 5 (1): 1-115. <https://doi.org/10.15468/ga63dr>
- CAMACHO, G. P.; PIE, M. R.; FEITOSA, R. M.; BARBEITOS, M. S. 2018. Exploring gene tree incongruence at the origin of ants and bees (Hymenoptera). *Zoologica Scripta* 48 (2): 215-225. <https://doi.org/10.1111/zsc.12332>
- CANTONE, S.; VON ZUBEN, C. J. 2019. The hind wings of ants: A phylogenetics analysis. *Psyche: A Journal of Entomology* 2019: 1-11. <https://doi.org/10.1155/2019/7929717>
- CARSTENS, B. C.; PELLETIER, T. A.; REID, N. M.; SATLER, J. D. 2013. How to fail at species delimitation. *Molecular Ecology* 22 (17): 4369-4383. <https://doi.org/10.1111/mec.12413>
- CHAUL, J. C. M. 2020. A new species of *Discothyrea* Roger (Hymenoptera: Formicidae) from the Brazilian Atlantic Rainforest. *Boletim do Museu Paraense Emílio Goeldi. Ciências Naturais* 15 (1): 199-217. <https://doi.org/10.46357/bcnaturais.v15i1.266>
- CRISTIANO, M. P.; CARDOSO, D. C.; SANDOVAL-GÓMEZ, V. E.; SIMÓES-GOMES, F. C. 2020. *Amoimyrmex* Cristiano, Cardoso, Sandoval, gen. nov. (Hymenoptera: Formicidae): a new genus of leaf-cutting ants revealed by multilocus molecular phylogenetic and morphological analyses. *Austral Entomology* 59 (4): 643-676. <https://doi.org/10.1111/aen.12493>
- CUEZZO, F. 2000. Revisión del género *Forelius* (Hymenoptera: Formicidae: Dolichoderinae). *Sociobiology* 35 (2A): 197-275. https://www.antwiki.org/wiki/images/9/92/Cuezzo,_F._2000.pdf
- CUEZZO, F.; CLAVER, S. 2009. Two new species of the ant genus *Pogonomyrmex* (Hymenoptera: Formicidae) from Argentina. *Revista de la Sociedad Entomológica Argentina* 68 (1-2): 97-106. https://ri.conicet.gov.ar/bitstream/handle/11336/79386/CONICET_Digital_Nro.72428f58-a54d-4fa2-9288-d42c01f92d67_A.pdf?sequence=2&isAllowed=y
- CUEZZO, F.; GUERRERO, R. J. 2012. The ant genus *Dorymyrmex* Mayr (Hymenoptera: Formicidae: Dolichoderinae) in Colombia. *Psyche: A Journal of Entomology* 2012: 1-24. <https://doi.org/10.1155/2012/516058>
- CUEZZO, F. C.; CALCATERRA, L.; CHIFFLET, L.; FOLLET, P. 2015. *Wasmannia* Forel (Hymenoptera: Formicidae: Myrmicinae) in Argentina. *Systematics and Distribution. Sociobiology* 62 (2): 246-265. <http://dx.doi.org/10.13102/sociobiology.v62i2.246-265>
- DASH, S. T. 2011. A Taxonomic revision of the New World *Hypoponera* Santschi, 1938 (Hymenoptera: Formicidae). Open Access Theses & Dissertations. University of Texas at El Paso. Texas, USA. 295 p.
- DE ANDRADE, M. L. 1998. Fossil and extant species of *Cylindromyrmex* (Hymenoptera: Formicidae). *Revue Suisse de Zoologie* 105: 581-664. <https://doi.org/10.5962/bhl.part.80052>
- DE ANDRADE, M. L. 2004. A new species of *Platythyrea* from Dominican amber and description of a new extant species from Honduras (Hymenoptera: Formicidae). *Revue Suisse de Zoologie* 111: 643-655. <https://doi.org/10.5962/bhl.part.80258>
- DE ANDRADE, M. L.; BARONI URBANI, C. 1999. Diversity and adaptation in the ant genus *Cephalotes*, past and present (Hymenoptera, Formicidae). *Stuttgarter Beiträge zur Naturkunde. Serie B (Geologie und Paläontologie)* 271: 1-889. <https://ia600206.us.archive.org/23/items/biostor-102758/biostor-102758.pdf>
- DEJACO, T.; GASSNERI, M.; ARTHOFERI, W.; SCHLICK-STEINERI, B. C.; STEINER, F. M. 2016. Taxonomist's nightmare... evolutionist's delight: an integrative approach resolves species limits in jumping bristletails despite widespread hybridization and parthenogenesis. *Systematic Biology* 65 (6): 947-974. <https://doi.org/10.1093/sysbio/syw003>
- DELSINNE, T.; FERNÁNDEZ, F. 2012. First record of *Lenomyrmex inusitatus* (Formicidae: Myrmicinae) in Ecuador and description of the queen. *Psyche: A Journal of Entomology* 2012: 1-5. <https://doi.org/10.1155/2012/145743>
- DELSINNE, T.; MACKAY, W. P.; WILD, A.; LE蓬CE, M. 2012. Distribution and diversity of the cryptic ant genus *Oxyepoecus* (Hymenoptera: Formicidae: Myrmicinae) in Paraguay with descriptions of two new species. *Psyche: A Journal of Entomology* 2012: 1-8. <https://doi.org/10.1155/2012/594302>
- DONOSO, D. A.; VIEIRA, J. M.; WILD, A. L. 2006. Three new species of *Leptanilloides* Mann from Andean Ecuador (Formicidae: Leptanilloidinae). *Zootaxa* 1201 (1): 47-62. <https://doi.org/10.11646/zootaxa.1201.1.2>
- ESCÁRRAGA, M. E.; GUERRERO, R. J. 2016. The ant genus *Linepithema* (Formicidae: Dolichoderinae) in Colombia. *Zootaxa* 4208 (5): 446-458. <https://doi.org/10.11646/zootaxa.4208.5.3>
- ESCÁRRAGA, M. E.; LATTKE, J. E.; PIE, M.; GUERRERO, R. J. 2021. Morphological and genetic evidence supports the separation of two *Tapinoma* ants (Formicidae: Dolichoderinae) from the Atlantic Forest biome. *Zookeys* 1026. in press.
- FEITOSA, R. M.; BRANDÃO, C. R. F. 2008. A taxonomic revision of the Neotropical myrmicine ant genus *Lachnomyrmex* Wheeler (Hymenoptera: Formicidae). *Zootaxa* 1890 (1): 1-49. <https://doi.org/10.11646/zootaxa.1890.1.1>

- FEITOSA, R. M.; PRADA-ACHIARDI, F. C. 2019. Subfamilia Ectatomminae. pp. 659-679. In: Fernández, F.; Guerrero, R. J.; Delsinne, T. (Eds.). Hormigas de Colombia. Universidad Nacional de Colombia, Bogotá, D. C., Colombia. 1198 p.
- FEITOSA, R. M.; BRANDÃO, C. R. F.; DINIZ, J. L. M. 2008. Revisionary studies on the enigmatic Neotropical ant genus *Stegomyrmex* Emery, 1912 (Hymenoptera: Formicidae: Myrmicinae), with the description of two new species. *Journal of Hymenoptera Research* 17 (1): 64-82.
- FEITOSA, R. M.; BRANDÃO, C. R. F.; FERNÁNDEZ, F.; DELABIE, J. C. H. 2012. The ant genus *Sphinctomyrmex* Mayr (Hymenoptera, Formicidae, Cerapachyinae) in the Neotropical region, with the description of two new species. *Psyche: A Journal of Entomology* 2012: 1-9. <https://doi.org/10.1155/2012/342623>
- FERNANDES, I. O.; SOUZA, J. L. P.; FERNÁNDEZ, F.; DELABIE, J. H. C.; SCHULTZ, T. R. 2015. A new species of *Simopelta* (Hymenoptera: Formicidae: Ponerinae) from Brazil and Costa Rica. *Zootaxa* 3956 (2): 295-300. <https://doi.org/10.11646/zootaxa.3956.2.10>
- FERNANDES, I. O.; DELABIE, J. H. C.; FERNÁNDEZ, F. 2019. Contribution to the knowledge of the Genus *Proceratium* Roger (Hymenoptera: Formicidae: Proceratiinae) in the New World. *Sociobiology* 66 (4): 551-559. <https://doi.org/10.13102/sociobiology.v66i4.4484>
- FERNÁNDEZ, F. 1991. Las hormigas cazadoras del género *Ectatoma* (Formicidae: Ponerinae) en Colombia. *Caldasia* 16 (79): 551-564. <https://revistas.unal.edu.co/index.php/cal/article/view/35681/36258>
- FERNÁNDEZ, F. 2001. Hormigas de Colombia. IX: Nueva especie de *Lenomyrmex* (Formicidae: Myrmicinae). *Revista Colombiana de Entomología* 27 (3-4): 201-204.
- FERNÁNDEZ, F. (Ed.). 2003a. Introducción a las hormigas de la región Neotropical. Instituto de Investigación de Recursos Biológicos Alexander von Humboldt. Bogotá, D. C., Colombia. 424 p. <http://repository.humboldt.org.co/bitstream/handle/20.500.11761/32961/978-958-8151-23-6.pdf?>
- FERNÁNDEZ, F. 2003b. Myrmicine ants of the genera *Ochetomyrmex* and *Tranopelta* (Hymenoptera: Formicidae). *Sociobiology* 41 (3): 633-661.
- FERNÁNDEZ, F. 2003c. Revision of the myrmicine ants of the *Adelomyrmex* genus-group (Hymenoptera: Formicidae). *Zootaxa* 361 (1): 1-52. <https://doi.org/10.11646/zootaxa.361.1.1>
- FERNÁNDEZ, F. 2004a. The American species of the myrmicine ant genus *Carebara* Westwood. *Caldasia* 26 (1): 191-238. <https://revistas.unal.edu.co/index.php/cal/article/view/39362>
- FERNÁNDEZ, F. 2004b. Adelomyrmecini new tribe and *Cryptomyrmex* new genus of myrmicine ants (Hymenoptera, Formicidae). *Sociobiology* 44 (2): 325-335. <http://antbase.org/ants/publications/publications/21105/21105.pdf>
- FERNÁNDEZ, F. 2007a. The myrmicine ant genus *Allomerus* Mayr (Hymenoptera: Formicidae). *Caldasia* 29 (1): 159-175. <https://revistas.unal.edu.co/index.php/cal/article/view/39221/62889>
- FERNÁNDEZ, F. 2007b. Two new South American species of *Monomorium* Mayr with taxonomic notes on the genus. *Memoirs of the American Entomological Institute* 80: 128-145. In: Snelling, R. R.; Fisher, B. L.; Ward, P. S. (Eds.). Advances in ant systematics (Hymenoptera: Formicidae): homage to E. O. Wilson – 50 years of contributions. *Memoirs of the American Entomological Institute*. Gainesville, USA. 690 p. <http://antbase.org/ants/publications/21280/21280.pdf>
- FERNANDEZ, F. 2008. Subfamilia Ponerinae s.str. pp. 123-218. In: Jiménez, E.; Fernández, F.; Arias, T. M.; Lozano-Zambrano, F. H. (Eds.). Sistemática, biogeografía y conservación de las hormigas cazadoras de Colombia. Instituto de Investigación de Recursos Biológicos Alexander von Humboldt. Bogotá, D. C., Colombia. 609 p. https://www.antwiki.org/wiki/images/f/f6/Jimenez_et_al_2008.pdf
- FERNÁNDEZ, F. 2011. The greatest impediment to the study of biodiversity in Colombia. *Caldasia* 33: iii-v. http://www.gfnun.unal.edu.co/uncienas/data-file/user_16/file/caldasia/3302/cld3302_ii.pdf
- FERNÁNDEZ, F.; GUERRERO, R. J. 2008. *Technomyrmex* (Formicidae: Dolichoderinae) in the New World: synopsis and description of a new species. *Revista Colombiana de Entomología* 34 (1): 110-115. <http://www.scielo.org.co/pdf/rcen/v34n1/v34n1a16.pdf>
- FERNÁNDEZ, F.; MACKAY, W. P. 2003. The myrmicine ants of the *Adelomyrmex laevigatus* species complex (Hymenoptera: Formicidae). *Sociobiology* 41 (3): 593-604.
- FERNÁNDEZ, F.; PALACIO G., E. E. 1997. Clave para las *Pogonomyrmex* (Hymenoptera: Formicidae) del Norte de Suramérica, con la descripción de una nueva especie. *Revista de Biología Tropical* 45 (4): 1649-1661.
- FERNÁNDEZ C., F.; PALACIO G., E. E. 1999. *Lenomyrmex*, an enigmatic new ant genus from the Neotropical region (Hymenoptera: Formicidae: Myrmicinae). *Systematic Entomology* 24 (1): 7-16. <https://doi.org/10.1046/j.1365-3113.1999.00063.x>
- FERNÁNDEZ, F.; BAENA, M. L.; PALACIO, E. E. 1996. Hormigas de Colombia V: el género *Stenamma* Westwood (Hymenoptera: Formicidae: Myrmicinae). *Tacaya* 5: 9-10.
- FERNÁNDEZ, F.; CASTRO-HUERTAS, V.; SERNA, F. 2015. Hormigas cortadoras de hojas de Colombia: *Acromyrmex* y *Atta* (Hymenoptera: Formicidae). *Fauna de Colombia, Monografía No.5. Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Bogotá, D. C., Colombia.* 350 p.
- FISHER, B. L. 2010. Biogeography. pp. 18-37. In: Lach, L.; Parr, C.; Abbot, K. (Eds.). *Ant ecology*. Oxford University. Oxford University Press. Oxford, United Kingdom. 402 p.
- FISCHER, G.; AZORSA, F.; FISHER, B. L. 2014. The ant genus *Carebara* Westwood (Hymenoptera, Formicidae): synonymisation of *Pheidologeton* Mayr under *Carebara*, establishment and revision of the *C. polita* species group. *ZooKeys* 438: 57-112. <https://doi.org/10.3897/zookeys.438.7922>
- FISCHER, G.; SARNAT, E. M.; ECONOMO, E. P. 2016. Revision and Microtomography of the *Pheidole knowlesi* Group, an endemic ant radiation in Fiji (Hymenoptera, Formicidae, Myrmicinae). *PLoS ONE* 11 (7): e0158544. <https://doi.org/10.1371/journal.pone.0158544>
- GALVIS, J. P.; FERNÁNDEZ, F. 2009. Ants of Colombia X. *Acanthognathus* with the description of a new species (Hymenoptera: Formicidae). *Revista Colombiana de Entomología* 35 (2): 245-249. <https://doi.org/10.25100/socolen.v35i2.9226>
- GARCIA-AVENDAÑO, E. I.; GUERRERO, R. J. 2018. Taxonomía y distribución de las hormigas del género *Tapinoma* (Formicidae: Dolichoderinae) en Colombia. *Revista Colombiana de Entomología* 44 (2): 223-237. <https://doi.org/10.25100/socolen.v44i2.7324>
- GONZÁLEZ-CAMPERO, M. C. L.; ELIZALDE, L. 2008. A new species of *Anochetus* (Hymenoptera: Formicidae: Ponerini) from Argentina and Paraguay, associated with leaf cutter ants. *Entomotropica* 23 (1): 97-102.
- GUERRERO, R. J. 2021. Transfer of two South American ant species from *Tapinoma* Foerster, 1850 to *Forelius* Emery, 1888 (Hymenoptera: Formicidae: Dolichoderinae). *Zootaxa* 4920 (3): 428-438. <https://doi.org/10.11646/zootaxa.4920.3.8>
- GUERRERO, R. J.; FERNÁNDEZ, F. 2008. A new species of the ant genus *Forelius* (Formicidae: Dolichoderinae) from the dry forest of Colombia. *Zootaxa* 1958 (1): 51-60. <https://doi.org/10.11646/zootaxa.1958.1.5>
- HITA GARCIA, F.; FISCHER, G.; LIU, C.; AUDISIO, T. L.; ALPERT, G. D.; FISHER, B. L.; ECONOMO, E. P. 2017. X-Ray microtomography for ant taxonomy: An exploration and case study with two new *Terataner* (Hymenoptera, Formicidae, Myrmicinae) species from Madagascar. *PLoS ONE* 12(3): e0172641. <https://doi.org/10.1371/journal.pone.0172641>

- HOENLE, P.; LATTKE, J. E.; DONOSO, D. A.; VON BEEREN, C.; HEETHOFF, M.; SCHMELZLE, S.; ARGOTI, A.; CAMACHO, L.; STRÖBEL, B.; BLÜTHGEN, N. 2020. *Odontomachus davidsoni* sp. nov. (Hymenoptera, Formicidae), a new conspicuous trap-jaw ant from Ecuador. ZooKeys 948: 75-105. <https://doi.org/10.3897/zookeys.948.48701>
- JEŠOVNIK, A.; SCHULTZ, T. R. 2017. Revision of the fungus-farming ant genus *Sericomyrmex* Mayr (Hymenoptera, Formicidae, Myrmicinae). ZooKeys 670: 1-119. <https://doi.org/10.3897/zookeys.670.11839>
- JEŠOVNIK, A.; SOSA-CALVO, J.; LLOYD, M. W.; BRANSTETTER, M. G.; FERNANDEZ, F.; SCHULTZ, T. R. 2017. Phylogenomic species delimitation and host-symbiont coevolution in the fungus-farming ant genus *Sericomyrmex* Mayr (Hymenoptera: Formicidae): Ultraconserved Elements (UCEs) resolve a recent radiation. Systematic Entomology 42 (3): 523-542. <https://doi.org/10.1111/syen.12228>
- JOHNSON, R. A.; MOREAU, C. S. 2016. A new ant genus from southern Argentina and southern Chile, *Patagonomyrmex* (Hymenoptera: Formicidae). Zootaxa 4139 (1): 1-31. <https://doi.org/10.11646/zootaxa.4139.1.1>
- JOHNSON, B. R.; BOROWIEC, M. L.; CHIU, J. C.; LEE, E. K.; ATALLAH, J.; WARD, P. S. 2013. Phylogenomics resolves evolutionary relationships among ants, bees, and wasps. Current Biology 23 (20): 2058-2062. <https://doi.org/10.1016/j.cub.2013.08.050>
- KELLER, R. A. 2011. A phylogenetic analysis of ant morphology (Hymenoptera: Formicidae) with special reference to the poneromorph subfamilies. Bulletin of the American Museum of Natural History 2011 (355): 1-90. <https://doi.org/10.1206/355.1>
- KINZNER, M. C.; WAGNER, H. C.; PESKOLLER, A.; MODER, K.; DOWELL, F. E.; ARTHOFER, W.; SCHLICK-STEINER, B. C.; STEINER, F. M. 2015. A near-infrared spectroscopy routine for unambiguous identification of cryptic ant species. PeerJ 3: e991. <https://doi.org/10.7717/peerj.991>
- KLINGENBERG, C.; BRANDÃO, C. R. F. 2009. Revision of the fungus-growing ant genera *Mycetophylax* Emery and *Paramycetophylax* Kusnezov rev. stat., and description of *Kalathomyrmex* n. gen. (Formicidae: Myrmicinae: Attini). Zootaxa 2052 (1): 1-31. <https://doi.org/10.11646/zootaxa.2052.1.1>
- KUGLER, C. 1994. A revision of the ant genus *Rogeria* (Hymenoptera: Formicidae) with description of the sting apparatus. Journal of Hymenoptera Research 3: 17-89.
- LACAU, S.; DELABIE, J. H. C. 2002. Description de trois nouvelles espèces d'*Amblyopone* avec quelques notes biogéographiques sur le genre au Brésil (Formicidae, Ponerinae). Bulletin de la Société Entomologique de France 107 (1): 33-41.
- LADINO, N.; FEITOSA, R. M. 2020. Taxonomic revision of the genus *Prionopelta* Mayr, 1866 (Formicidae: Amblyoponinae) for the Neotropical region. Zootaxa 4821 (2): 201-249. <https://doi.org/10.11646/zootaxa.4821.2.1>
- LAPOLLA, J. S. 2004. *Acropyga* (Hymenoptera: Formicidae) of the world. Contributions of the American Entomological Institute 33 (3): 1-130. <http://antcat.org/documents/3924/21151.pdf>
- LAPOLLA, J. S.; SOSA-CALVO, J. 2006. Review of the ant genus *Rogeria* (Hymenoptera: Formicidae) in Guyana. Zootaxa 1330 (1): 59-68. <https://doi.org/10.11646/zootaxa.1330.1.5>
- LAPOLLA, J. S.; KALLAL, R. J.; BRADY, S. G. 2012. A new ant genus from the Greater Antilles and Central America, *Zatania* (Hymenoptera: Formicidae), exemplifies the utility of male and molecular character systems. Systematic Entomology 37 (1): 200-214. <https://doi.org/10.1111/j.1365-3113.2011.00605.x>
- LAPOLLA, J. S.; DLUSSKY, G. M.; PERRICHOT, V. 2013. Ants and the fossil record. The Annual Review of Entomology 58: 609-630. <https://doi.org/10.1146/annurev-ento-120710-100600>
- LATTKE, J. E. 1991. Studies of Neotropical *Amblyopone* Erichson (Hymenoptera: Formicidae). Contributions in Science (Los Angeles) 428: 1-7.
- LATTKE, J. E. 1997. Revisión del género *Apterostigma* Mayr (Hymenoptera: Formicidae). Arquivos de Zoología 34 (5): 121-221. <https://doi.org/10.11606/issn.2176-7793.v34i5p121-221>
- LATTKE, J. E. 2006. A new species of *Pogonomyrmex* (Hymenoptera: Formicidae) from gallery forests of the Orinoco Watershed, Venezuela. Myrmecologische Nachrichten 8: 53-57.
- LATTKE, J. E. 2011. Revision of the New World species of the genus *Leptogenys* Roger (Insecta: Hymenoptera: Formicidae: Ponerinae). Arthropod Systematics and Phylogeny 69 (3): 127-264.
- LATTKE, J. E.; MELO, G. A. R. 2020. New haidomyrmecine ants (Hymenoptera: Formicidae) from mid-Cretaceous amber of northern Myanmar. Cretaceous Research 114: 104502. <https://doi.org/10.1016/j.cretres.2020.104502>
- LATTKE, J. E.; FERNÁNDEZ, F.; PALACIO, E. E. 2007. Identification of the species of *Gnamptogenys* Roger in the Americas. pp. 254-270. In: Snelling, R. R.; Fisher, B. L.; Ward, P. S. (Eds.). Advances in ant systematics (Hymenoptera: Formicidae): homage to E. O. Wilson – 50 years of contributions. Memoirs of the American Entomological Institute 80. Gainesville, USA. 690 p. <http://antbase.org/ants/publications/21283/21283.pdf>
- LATTKE, J. E.; FERNÁNDEZ, F.; ARIAS-PENNA, T. M.; PALACIO, E. E.; MACKAY, W. P.; MACKAY, E. 2008. Género *Gnamptogenys* Roger. pp. 66-107. In: Jiménez, E.; Fernández, F.; Arias, T. M.; Lozano-Zambrano, F. H. (Eds.). Sistemática, biogeografía y conservación de las hormigas cazadoras de Colombia. Instituto de Investigación de Recursos Biológicos Alexander von Humboldt. Bogotá, D. C., Colombia. 609 p.
- LATTKE, J. E.; DELSINNE, T.; ALPERT, G. D.; GUERRERO, R. J. 2018. Ants of the genus *Protalaridris* (Hymenoptera: Formicidae), more than just deadly mandibles. European Journal of Entomology 115: 268-295. <https://doi.org/10.14411/eje.2018.027>
- LEAVITT, S. D.; MOREAU, C. S.; LUMBSCH, H. T. 2015. The dynamic discipline of species delimitation: progress toward effectively recognizing species boundaries in natural populations. pp. 11-44. In: Upreti, D. K.; Divakar, P. K.; Shukla, V.; Bajpai, R. (Eds.). Recent advances in lichenology: modern methods and approaches in lichen systematics and culture techniques, Volume 2. Springer, New Delhi, India. 232 p. https://doi.org/10.1007/978-81-322-2235-4_2
- LENHART, P. A.; DASH, S. T.; MACKAY, W. P. 2013. A revision of the giant Amazonian ants of the genus *Dinoponera* (Hymenoptera, Formicidae). Journal of Hymenoptera Research 31: 119-164. <https://doi.org/10.3897/jhr.31.4335>
- LONGINO, J. T. 2003. The *Crematogaster* (Hymenoptera, Formicidae, Myrmicinae) of Costa Rica. Zootaxa 151 (1): 1-150. <https://doi.org/10.11646/zootaxa.151.1>
- LONGINO, J. T. 2006a. A taxonomic review of the genus *Myrmelachista* (Hymenoptera: Formicidae) in Costa Rica. Zootaxa 1141 (1): 1-54. <https://www.mapress.com/zt/article/view/zootaxa.1141.1.1>
- LONGINO, J. T. 2006b. New species and nomenclatural changes for the Costa Rican ant fauna (Hymenoptera: Formicidae). Myrmecologische Nachrichten 8: 131-143. <http://antbase.org/ants/publications/21107/21107.pdf>
- LONGINO, J. T. 2009a. Additions to the taxonomy of New World *Pheidole* (Hymenoptera: Formicidae). Zootaxa 2181 (1): 1-90. <https://doi.org/10.11646/zootaxa.2181.1.1>
- LONGINO, J. T. 2009b. *Pheidole* working group. Available in: <https://ants.biology.utah.edu/pheidoleworkinggroup/index.htm> [Review date: 21 February 2021].
- LONGINO, J. T. 2010. A taxonomic review of the ant genus *Megalomyrmex* Forel (Hymenoptera: Formicidae) in Central America. Zootaxa 2720 (1): 35-58. <https://doi.org/10.11646/zootaxa.2720.1>
- LONGINO, J. T. 2012. A review of the ant genus *Adelomyrmex* Emery 1897 (Hymenoptera, Formicidae) in Central America. Zootaxa 3456 (1): 1-35. <https://doi.org/10.11646/zootaxa.3456.1.1>

- LONGINO, J. T. 2013. A revision of the ant genus *Octostruma* Forel, 1912 (Hymenoptera, Formicidae). *Zootaxa* 3699 (1): 1-61. <https://doi.org/10.11646/zootaxa.3699.1.1>
- LONGINO, J. T.; BOUDINOT, B. E. 2013. New species of Central American *Rhopalothrix* Mayr, 1870 (Hymenoptera, Formicidae). *Zootaxa* 3616 (4): 301-324. <https://doi.org/10.11646/zootaxa.3616.4.1>
- LONGINO, J. T.; BRANSTETTER, M. G. 2020. Phylogenomic species delimitation, taxonomy, and ‘Bird Guide’ identification for the Neotropical ant genus *Rasopone* (Hymenoptera: Formicidae). *Insect Systematics and Diversity* 4 (2): 1-33. <https://doi.org/10.1093/isd/ixaa004>
- LONGINO, J. T.; BRANSTETTER, M. G. 2021. Integrating UCE phylogenomics with traditional taxonomy reveals a trove of New World *Syscia* species (Formicidae: Dorylinae). *Insect Systematics and Diversity* 5 (2): 1-51. <https://doi.org/10.1093/isd/ixab001>
- LONGINO, J. T.; COVER, S. 2004. A revision of the *Aphaenogaster phalangium* complex (Hymenoptera: Formicidae: Myrmicinae). *Zootaxa* 655 (1): 1-12. <https://doi.org/10.11646/zootaxa.655.1.1>
- LONGINO, J. T.; FERNÁNDEZ, F. 2007. Taxonomic review of the genus *Wasemannia*. pp. 271-289. In: Snelling, R. R.; Fisher, B. L.; Ward, P. S. (Eds.). Advances in ant systematics (Hymenoptera: Formicidae): homage to E. O. Wilson – 50 years of contributions. Memoirs of the American Entomological Institute 80. Gainesville, USA. 690 p. <http://antbase.org/ants/publications/21284/21284.pdf>
- LONGINO, J. T.; SNELLING, R. R. 2002. A taxonomic revision of the *Procryptocerus* (Hymenoptera: Formicidae) of Central America. *Contributions in Science (Los Angeles)* 495: 1-30. <https://antcat.org/references/131733>
- LUCKY, A.; TRAUTWEIN, M. D.; GUÉNARD, B. S.; WEISER, M. D.; DUNN, R. R. 2013. Tracing the rise of ants-out of the ground. *PLoS ONE* 8 (12): e84012. <https://doi.org/10.1371/journal.pone.0084012>
- MACKAY, W. P. 1995. New distributional records for the ant genus *Cardiocondyla* in the New World (Hymenoptera: Formicidae). *Pan-Pacific Entomologist* 71 (3): 169-172. <http://antbase.org/ants/publications/14430/14430.pdf>
- MACKAY, W. P. 1996. A revision of the ant genus *Acanthostichus* (Hymenoptera: Formicidae). *Sociobiology* 27 (2): 129-179. <http://antbase.org/ants/publications/14660/14660.pdf>
- MACKAY, W. P. 2019a. New World carpenter ants of the hyper-diverse genus *Camponotus* (Hymenoptera: Formicidae), Volume 1: introduction, subgenera, species complexes, and subgenus *Camponotus*. Lambert Academic Publishing, 420 p.
- MACKAY, W. P. 2019b. Género *Camponotus*. pp. 743-789. In: Fernández, F.; Guerrero, R. J.; Delsinne, T. (Eds.). Hormigas de Colombia. Universidad Nacional de Colombia. Bogotá D. C., Colombia. 1198 p.
- MACKAY, W. P.; MACKAY, E. 2002. The ants of New Mexico (Hymenoptera: Formicidae). Edwin Mellen Press. Lewiston, New York, USA. 408 p.
- MACKAY, W. P.; MACKAY, E. 2008. Revision of the ants of the genus *Simopelta* Mann. pp. 285-328. In: Jiménez, E.; Fernández, F.; Arias, T. M.; Lozano-Zambrano, F. H. (Eds.). Sistemática, biogeografía y conservación de las hormigas cazadoras de Colombia. Instituto de Investigación de Recursos Biológicos Alexander von Humboldt. Bogotá, D. C., Colombia. 609 p.
- MACKAY, W. P.; MACKAY, E. 2010. The systematics and biology of the New World ants of the genus *Pachycondyla* (Hymenoptera: Formicidae). Edwin Mellen Press. Lewiston, New York, USA. 642 p.
- MACKAY, W. P.; MACKAY, E. 2019. Género *Camponotus*. pp. 743-790. In: Fernández, F.; Guerrero, R. J.; Delsinne, T. (Eds.). Hormigas de Colombia. Universidad Nacional de Colombia, Bogotá, D. C., Colombia. 1198 p.
- MACKAY, W. P.; MAES, J.-M.; FERNÁNDEZ, P. R.; LUNA, G. 2004. The ants of North and Central America: the genus *Mycoceropurus* (Hymenoptera: Formicidae). *Journal of Insect Science* 4 (27): 1-7. <https://doi.org/10.1673/031.004.2701>
- MARSHALL, S. A.; EVENHUIS, N. L. 2015. New species without dead bodies: A case for photobased descriptions, illustrated by a striking new species of *Marleyomyia* Hesse (Diptera, Bombyliidae) from South Africa. *ZooKeys* 2015 (525): 117-127. <https://doi.org/10.3897/zookeys.525.6143>
- MAYHÉ-NUNES, A. J.; BRANDÃO, C. R. F. 2002. Revisionary studies on the attine ant genus *Trachymyrnex* Forel. Part 1: definition of the genus and the *Opulentus* group (Hymenoptera: Formicidae). *Sociobiology* 40: 667-698.
- MAYHÉ-NUNES, A. J.; BRANDÃO, C. R. F. 2005. Revisionary studies on the attine ant genus *Trachymyrnex* Forel. Part 2: the *Iheringi* group (Hymenoptera: Formicidae). *Sociobiology* 45: 271-305.
- MAYHÉ-NUNES, A. J.; BRANDÃO, C. R. F. 2007. Revisionary studies on the attine ant genus *Trachymyrnex* Forel. Part 3: The *Jamaicensis* group (Hymenoptera: Formicidae). *Zootaxa* 1444 (1): 1-21. <https://doi.org/10.11646/zootaxa.1444.1.1>
- MISOF, B.; LIU, S.; MEUSEMANN, K.; PETERS, R. S.; DONATH, A.; MAYER, C.; FRANDSEN, P. B.; WARE, J.; FLOURI, T.; BEUTEL, R. G.; NIEHUIS, O.; PETERSEN, M.; IZQUIERDO-CARRASCO, F.; WAPPLER, T.; RUST, J.; ABERER, A. J.; ASPÖCKI, U.; ASPÖCK, H.; BARTEL, D.; BLANKE, A.; BERGER, S.; BÖHM, A.; BUCKLEY, T. R.; CALCOTT, B.; CHEN, J.; FRIEDRICH, F.; FUKUI, M.; FUJITA, M.; GREVE, C.; GROBE, P.; GU, S.; HUANG, Y.; JERMIIN, L. S.; KAWAHARA, A. Y.; KROGMANN, L.; KUBIAK, M.; LANFEAR, R.; LETSCH, H.; LI, Y.; LI, Z.; LI, J.; LU, J.; MACHIDA, R.; MASHIMO, Y.; KAPLI, P.; MCKENNA, D. D.; MENG, G.; NAKAGAKI, Y.; NAVARRETE-HEREDIA, J. L.; OTT, M.; OU, Y.; PASS, G.; PODSIADLOWSKI, L.; POHL, H.; VON REUMONT, B. M.; SCHÜTTE, K.; SEKIYA, K.; SHIMIZU, S.; SLIPINSKI, A.; STAMATAKIS, A.; SONG, W.; SU, X.; SZUCSICH, N. U.; TAN, M.; TAN, X.; TANG, M.; TANG, J.; TIMELTHALER, G.; TOMIZUKA, S.; TRAUTWEIN, M.; TONG, X.; UCHIFUNE, T.; WALZL, M. G.; WIEGMANN, B. M.; WILBRANDT, J.; WIPFLER, B.; WONG, T. K. F.; WU, Q.; WU, G.; XIE, Y.; YANG, S.; YANG, Q.; YEATES, D. K.; YOSHIZAWA, K.; ZHANG, Q.; ZHANG, R.; ZHANG, W.; ZHANG, Y.; ZHAO, J.; ZHOU, C.; ZHOU, L.; ZIESMANN, T.; ZOU, S.; LI, Y.; XU, X.; ZHANG, Y.; YANG, H.; WANG, J.; WANG, J.; KJER, K. M.; ZHOU, X. 2014. Phylogenomics resolves the timing and pattern of insect evolution. *Science* 346 (6210): 763-767. <https://doi.org/10.1126/science.1257570>
- MOREAU, C. S.; BELL, C. D. 2013. Testing the museum versus cradle tropical biological diversity hypothesis: phylogeny, diversification, and ancestral biogeographic range evolution of the ants. *Evolution* 67 (8): 2240-2257. <https://doi.org/10.1111/evo.12105>
- MOREAU, C. S.; BELL, C. D.; VILA, R.; ARCHIBALD, S. B.; PIERCE, N. E. 2006. Phylogeny of the ants: diversification in the age of angiosperms. *Science* 312 (5770): 101-104. <https://doi.org/10.1126/science.1124891>
- OLIVEIRA, A. M.; FEITOSA, R. M. 2019. Taxonomic revision of the genus *Probolomyrmex* Mayr, 1901 (Hymenoptera: Formicidae: Proceratiinae) for the Neotropical Region. *Zootaxa* 4614 (1): 061-094. <https://doi.org/10.11646/zootaxa.4614.1.3>
- ORTIZ, C. M.; FERNÁNDEZ, F. 2011. Hormigas del género *Dolichoderus* Lund (Formicidae: Dolichoderinae) en Colombia. (Monografías de Fauna de Colombia 3). Universidad Nacional de Colombia. Bogotá, D. C., Colombia. 116 p. https://www.phd-eep.com/wp-content/uploads/2020/11/8_HORMIGAS_DOLICHODERUS-COLOMBIA-1.pdf
- ORTIZ, C. M.; FERNÁNDEZ, F. 2014. *Brachymyrmx* species with tumuliform metathoracic spiracles: description of three new species and discussion of dimorphism in the genus

- (Hymenoptera, Formicidae). ZooKeys 371: 13-33. <https://doi.org/10.3897/zookeys.371.6568>
- ORTIZ-SEPULVEDA, C. M.; VAN BOCXLAER, B.; MENESES, A. D.; FERNÁNDEZ, F. 2019. Molecular and morphological recognition of species boundaries in the neglected ant genus *Brachymyrmex* (Hymenoptera: Formicidae): toward a taxonomic revision. Organism Diversity and Evolution 19: 447-542. <https://doi.org/10.1007/s13127-019-00406-2>
- PACHECO, J. A.; MACKAY, W. P. 2013. The systematics and biology of the New World thief ants of the genus *Solenopsis* (Hymenoptera: Formicidae). Edwin Mellen Press. Lewiston, New York, USA. 501 p.
- PALACIO, E. E. 1997. Hormigas de Colombia VI. Dos nuevas especies de *Ocstrostruma* (Hymenoptera: Formicidae: Basicerotini). Caldasia 19 (3): 409-418. <https://revistas.unal.edu.co/index.php/cal/article/view/17442/18317>
- PALACIO, E. E. 1999. Hormigas legionarias (Hymenoptera: Formicidae: Ecitoninae) de Colombia. pp. 117-189. In: Amat, G.; Andrade, G.; Fernández, F. (Eds.). Insectos de Colombia Vol. II. Academia Colombiana de Ciencias Exactas, Físicas y Naturales y Facultad de Ciencias Universidad Nacional de Colombia, Bogotá, D. C., Colombia. 138 p.
- PALACIO, E. E. 2019. Subfamilia Dorylinae. pp. 571-630. In: Fernández, F.; Guerrero, R. J.; Delsinne, T. (Eds.). Hormigas de Colombia. Universidad Nacional de Colombia. Bogotá, D. C., Colombia. 1198 p.
- PETERS, R. S.; KROGMANN, L.; MAYER, C.; DONATH, A.; GUNKEL, S.; MEUSEMANN, K.; KOZLOV, A.; PODSIADLOWSKI, L.; PETERSEN, M.; LANFEAR, R.; DIEZ, P. A.; HERATY, J.; KJER, K. M.; KLOPFSTEIN, S.; MEIER, R.; POLIDORI, C.; SCHMITT, T.; LIU, S.; ZHOU, X.; WAPPLER, T.; RUST, J.; MISOF, B.; NIEHUIS, O. 2017. Evolutionary history of the Hymenoptera. Current Biology 27 (7): 1013-1018. <http://dx.doi.org/10.1016/j.cub.2017.01.027>
- PILGRIM, E. M.; VON DOHLEN, C. D.; PITTS, J. P. 2008. Molecular phylogenetics of Vespoidea indicate paraphyly of the superfamily and novel relationships of its component families and subfamilies. Zoologica Scripta 37 (5): 539-560. <https://doi.org/10.1111/j.1463-6409.2008.00340.x>
- PITTS, J. P.; CAMACHO, G. P.; GOTZEK, D.; MCHUGH, J. V.; ROSS, K. G. 2018. Revision of the fire ants of the *Solenopsis saevissima* species-group (Hymenoptera: Formicidae). Proceedings of the Entomological Society of Washington 120 (2): 308-411. <https://doi.org/10.4289/0013-8797.120.2.308>
- PREBUS, M. M. 2020. Phylogenomic species delimitation in the ants of the *Temnothorax salvinii* group (Hymenoptera: Formicidae): an integrative approach. Systematic Entomology 46 (2): 307-326. <https://doi.org/10.1111/syen.12463>
- PROBST, R. S.; WRAY, B. D.; MOREAU, C. S.; BRANDÃO, C. R. F. 2019. A Phylogenetic analysis of the dirt ants, *Basiceros* (Formicidae: Myrmicinae): Inferring life histories through morphological convergence. Insect Systematics and Diversity 3 (4): 1-12. <https://doi.org/10.1093/isd/ixz013>
- QUEIROZ, K. 2007. Species concepts and species delimitation. Systematic Biology 56 (6): 879-886. <https://doi.org/10.1080/10635150701701083>
- RABELING, C.; BROWN, J. M.; VERHAAGH, M. 2008. Newly discovered sister lineage sheds light on early ant evolution. Proceedings of the National Academy of Science of the United State of North America 105 (39): 14913-14917. <https://doi.org/10.1073/pnas.0806187105>
- RODRÍGUEZ, J. 2008. Género *Odontomachus* Latreille. pp. 148-218. In: Jiménez, E.; Fernández, F.; Arias, T. M.; Lozano-Zambrano, F. H. (Eds.). Sistemática, biogeografía y conservación de las hormigas cazadoras de Colombia. Instituto de Investigación de Recursos Biológicos Alexander von Humboldt. Bogotá, D. C., Colombia. 609 p.
- SANDOVAL-GÓMEZ, V. E.; SÁNCHEZ-RESTREPO, A. F. 2019. Género *Cephalotes*. pp. 899-915. In: Fernández, F.; Guerrero, R. J.; Delsinne, T. (Eds.). Hormigas de Colombia. Universidad Nacional de Colombia. Facultad de Ciencias. Instituto de Ciencias Naturales. Bogotá, D. C., Colombia. 1198 p.
- SARNAT, E. M.; FISCHER, G.; ECONOMO, E. P. 2016. Inordinate spinescence: taxonomic revision and Microtomography of the *Pheidole cervicornis* species group (Hymenoptera, Formicidae). PLoS ONE 11 (7): e0156709. <https://doi.org/10.1371/journal.pone.0156709>
- SCHLICK-STEINER, B. C.; STEINER, F. M.; SEIFERT, B.; STAUFFER, C.; CHRISTIAN, E.; CROZIER, R. H. 2010. Integrative taxonomy: A multisource approach to exploring biodiversity. Annual Review of Entomology 55 (1): 421-438. <https://doi.org/10.1146/annurev-ento-112408-085432>
- SCHMIDT, C. 2013. Molecular phylogenetics of ponerine ants (Hymenoptera: Formicidae: Ponerinae). Zootaxa 3647 (2): 201-250. <https://doi.org/10.11646/zootaxa.3647.2.1>
- SNELLING, R. R.; LONGINO, J. T. 1992. Revisionary notes on the fungus-growing ants of the genus *Cyphomyrmex, rimosus* group (Hymenoptera: Formicidae: Attini). pp. 479-494. In: Quintero, D.; Aiello, A. (Eds.). Insects of Panama and Mesoamerica: selected studies. Oxford University. Oxford University Press, Oxford, United Kingdom. 692 p.
- SOSA-CALVO, J.; LONGINO, J. T. 2008. Subfamilia Proceratiinae. pp. 219-237. In: Jiménez, E.; Fernández, F.; Arias, T. M.; Lozano-Zambrano, F. H. (Eds.). Sistemática, biogeografía y conservación de las hormigas cazadoras de Colombia. Instituto de Investigación de Recursos Biológicos Alexander von Humboldt. Bogotá, D. C., Colombia. 609 p. https://www.antwiki.org/wiki/images/a/aa/Longino_J_Sosa-Calvo_J_2008.pdf
- SOSA-CALVO, J.; SCHULTZ, T. R. 2010. Three remarkable new fungus-growing ant species of the genus *Myrmicocrypta* (Hymenoptera: Formicidae), with a reassessment of the characters that define the genus and its position within the Attini. Annals of the Entomological Society of America 103 (2): 181-195. <https://doi.org/10.1603/AN09108>
- SOSA-CALVO, J.; JEŠOVNIK, A.; VASCONCELOS, H. L.; BACCI, M.; SCHULTZ, T. R. 2017. Rediscovery of the enigmatic fungus-farming ant “*Mycetosoritis asper*” Mayr (Hymenoptera: Formicidae): Implications for taxonomy, phylogeny, and the evolution of agriculture in ants. PLoS ONE 12 (7): e0176498. <https://doi.org/10.1371/journal.pone.0176498>
- SPARKS, K. S.; ANDERSEN, A. N.; AUSTIN, A. D. 2019. A multi-gene phylogeny of Australian *Monomorium* Mayr (Hymenoptera: Formicidae) results in reinterpretation of the genus and resurrection of *Chelaner* Emery. Invertebrate Systematics 33 (1): 225-236. <https://doi.org/10.1071/IS16080>
- TAYLOR, R. W. 2007. Bloody funny wasps! Speculations on the evolution of eusociality in ants. pp. 580-609. In Snelling, R. R.; Fisher, B. L.; Ward, P. S. (Eds.). Advances in ant systematics (Hymenoptera: Formicidae): homage to E. O. Wilson – 50 years of contributions. Memoirs of the American Entomological Institute. Gainesville, USA. 690 p. <http://antbase.org/ants/publications/21292/21292.pdf>
- TRAGER, J. C. 1991. A revision of the fire ants, *Solenopsis geminata* group (Hymenoptera: Formicidae: Myrmicinae). Journal of the New York Entomological Society 99 (2): 141-198. <https://www.jstor.org/stable/25009890>
- WARD, P. S. 1989. Systematic studies on pseudomyrmecine ants: revision of the *Pseudomyrmex oculatus* and *P. subtilissimus* species groups, with taxonomic comments on other species. Quaestiones Entomologicae 25 (4): 393-468. <https://zenodo.org/record/24960#.YF4RtGQzYdU>
- WARD, P. S. 1999. Systematics, biogeography and host plant associations of the *Pseudomyrmex viduus* group (Hymenoptera: Formicidae), *Triplaris-* and *Tachigali-*inhabiting ants. Zoological Journal of the Linnean Society 126 (4): 451-540. <https://doi.org/10.1111/j.1096-3642.1999.tb00157.x>
- WARD, P. S. 2010. Taxonomy, phylogenetics and evolution. pp. 3-17. In: Lach, L.; Parr, C.; Abbott, K. (Eds.). Ant

- ecology. Oxford University. Oxford University Press, Oxford, United Kingdom. 402 p. <https://doi.org/10.1093/acprof:oso/9780199544639.003.0001>
- WARD, P. S. 2014. The phylogeny and evolution of ants. *Annual Review of Ecology, Evolution, and Systematics* 45: 23-43. <https://doi.org/10.1146/annurev-ecolsys-120213-091824>
- WARD, P. S. 2017. A review of the *Pseudomyrmex ferrugineus* and *Pseudomyrmex goeldii* species groups: acacia-ants and relatives (Hymenoptera: Formicidae). *Zootaxa* 4227 (4): 524-542. <https://doi.org/10.11646/zootaxa.4227.4.3>
- WARD, P. S.; DOWNIE, D. A. 2005. The ant subfamily *Pseudomyrmecinae* (Hymenoptera: Formicidae): phylogeny and evolution of big-eyed arboreal ants. *Systematic Entomology* 30 (2): 310-335. <https://doi.org/10.1111/j.1365-3113.2004.00281.x>
- WARD, P. S.; FISHER, B. L. 2016. Tales of Dracula ants: the evolutionary history of the ant subfamily *Amblyoponinae* (Hymenoptera: Formicidae). *Systematic Entomology* 41 (3): 683-693. <https://doi.org/10.1111/syen.12186>
- WARD, P. S.; BRADY, S. G.; FISHER, B. L.; SCHULTZ, T. R. 2010. Phylogeny and biogeography of dolichoderine ants: effects of data partitioning and relict taxa on historical inference. *Systematic Biology* 59 (3): 342-362. <https://doi.org/10.1093/sysbio/syq012>
- WARD, P. S.; BRADY, S. G.; FISHER, B. L.; SCHULTZ, T. R. 2015. The evolution of myrmicine ants: phylogeny and biogeography of a hyperdiverse ant clade (Hymenoptera: Formicidae). *Systematic Entomology* 40 (1): 61-81. <https://doi.org/10.1111/syen.12090>
- WHITFIELD, J. B.; KJER, K. M. 2008. Ancient rapid radiations of insects: challenges for phylogenetic analysis. *Annual Review of Entomology* 53: 449-472. <https://doi.org/10.1146/annurev.ento.53.103106.093304>
- WILD, A. L. 2002. The genus *Pachycondyla* (Hymenoptera: Formicidae) in Paraguay. *Boletín del Museo Nacional de Historia Natural del Paraguay* 14 (1-2): 1-18.
- WILD, A. L. 2004. Taxonomy and distribution of the Argentine ant, *Linepithema humile* (Hymenoptera: Formicidae). *Annals of the Entomological Society of America* 97 (6): 1204-1215. [https://doi.org/10.1603/0013-8746\(2004\)097\[1204:TADOTA\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2004)097[1204:TADOTA]2.0.CO;2)
- WILD, A. L. 2005. Taxonomic revision of the *Pachycondyla apicalis* species complex (Hymenoptera: Formicidae). *Zootaxa* 834 (1): 1-25. <https://doi.org/10.11646/zootaxa.834.1.1>
- WILD, A. L. 2007. Taxonomic revision of the ant genus *Linepithema* (Hymenoptera: Formicidae). *University of California Publications in Entomology*. Volumen 126. University of California Press, Ltd. London, England. 151 p.
- WILSON, E. O. 2003. *Pheidole* in the New World. A dominant, hyperdiverse ant genus. Harvard University Press, Cambridge, Massachusetts, USA. 794 p.
- WILSON, E. O.; CARPENTER, F. M.; BROWN, W. L. Jr. 1967. The first Mesozoic ants, with the description of a new subfamily. *Psyche: A Journal of Entomology* 74: 1-19. <https://doi.org/10.1155/1967/89604>
- ZABALA, G. 2008. Género *Anochetus* Mayr. pp. 124-142. In: Jiménez, E.; Fernández, F.; Arias, T. M.; Lozano-Zambrano, F. H. (Eds.). *Sistemática, biogeografía y conservación de las hormigas cazadoras de Colombia*. Instituto de Investigación de Recursos Biológicos Alexander von Humboldt. Bogotá, D. C., Colombia. 609 p.

Origin and funding

The origin of this review dates back several years, when the authors started gathering information on the diversity of ants at both the national and Neotropical levels, and considered that a synthesis of valid taxa and updated literature could be relevant for a group of insects as important as the ants. In this sense, there are several sources of funding for institutions such as the Universidad Nacional de Colombia, Pontificia Universidad Javeriana and Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Universidad del Magdalena, Colciencias (today Ministerio de Ciencias of Colombia), Smithsonian Institution and many others.

Author contribution

The manuscript is based on texts written by Fernando Fernández, Roberto J. Guerrero and Thibaut Delsinne, published in Spanish in the book Hormigas de Colombia in 2019. From parts of this source, Fernando Fernández and Roberto J. Guerrero made a first update and addition of figures, and then Andrés F. Sánchez-Restrepo added more text, references and a redo of the figures. Fernando Fernández wrote a first version in English. All authors reviewed the final version, add changes and revise the references equally.