

Crabronidae (Hymenoptera: Apoidea) in Mexico: Occurrence and Potential Uses in Biological Control

Crabronidae (Hymenoptera: Apoidea) en México: Ocurrencia y usos potenciales en el control biológico

 MAXIMILIANO VANOE-ELIGIO^{1,*},  JORGE VICTOR HORTA-VEGA²,
 VENANCIO VANOE-ELIGIO³,  LUDWIN JAIME ESTRADA-RAMÍREZ¹,
 BLANCA DEL ROSARIO MARTÍN-CANCHÉ¹

¹ Tecnológico Nacional de México/Instituto Tecnológico Superior de Escárcega, Campeche, México. C.P. maxvanoye@itsescarcega.edu.mx, estrada_ramirez@itsescarcega.edu.mx, blanca_martin@itsescarcega.edu.mx

² Tecnológico Nacional de México/Instituto Tecnológico de Ciudad Victoria, Tamaulipas, México. jorge.hv@cdvictoria.tecnm.mx

³ Universidad Autónoma de Tamaulipas, Tamaulipas, México. vvanoye@docentes.uat.edu.mx

Abstract: The family Crabronidae (order Hymenoptera) includes excavating and hunting wasps of more than 9,009 species within 249 genera that are distributed globally, especially in the tropics. The objective of this research was to improve the knowledge of the occurrence of the family Crabronidae in Mexico from a review of the scientific literature and to provide data on species of importance in biological control. The results revealed that 769 species representing 99 genera have been reported in Mexico. With the current review, 8.5% of the species known worldwide were found to be present in the country and the list of species was increased by 32%. The genera *Cerceris* and *Trypoxyylon* had the highest number of species, 130 species and 58 species, respectively. The Mexican states of Baja California Norte, Sonora and Tamaulipas have the highest number of Crabronidae species. A total of 211 species were added to the list of Crabronidae in Mexico. Twelve genera and eight species of Crabronidae used as biological control agents of various pests were recognized. Further research on the diversity of this group of insects is suggested, given the importance that crabonids have as predators and natural enemies of insect pests that affect economically important crops.

* Corresponding Author

Tecnológico Nacional de México/Instituto Tecnológico Superior de Escárcega. Escárcega. Calle 85 s/n entre 10B, Colonia Unidad, Esfuerzo y Trabajo No. 1. Campeche, México. C.P. maxvanoye@itsescarcega.edu.mx

Suggested Citation

VANOYE-ELIGIO, M.; HORTA-VEGA, J. V.; VANOYE-ELIGIO, V.; ESTRADA-RAMÍREZ, L. J.; MARTÍN-CANCHÉ, B. R. 2022. Crabronidae (Hymenoptera: Apoidea) in Mexico: Occurrence and Potential Uses in Biological Control. Revista Colombiana de Entomología 48 (1): e11654. <https://doi.org/10.25100/socolen.v48i1.11654>

Received: Oct-15-2021

Accepted: Mar-22-2022

Published: jun-11-2022

Revista Colombiana de Entomología

ISSN (Print): 0120-0488

ISSN (On Line): 2665-4385

<https://revistacolombianaintomologia.univalle.edu.co>

Open access

 BY-NC-SA 4.0 creativecommons.org/licenses/by-nc-sa/4.0/

Publishers: Sociedad Colombiana de Entomología SOCOLEN (Bogotá, D. C., Colombia)

<https://www.socolen.org.co>

Universidad del Valle (Cali, Colombia)

<https://www.univalle.edu.co>

© 2021 Sociedad Colombiana de Entomología - SOCOLEN y Universidad del Valle - Univalle

Keywords: *Cerceris*, distribution, predators, wasps.

Resumen: La familia Crabronidae (orden Hymenoptera) incluye avispas excavadoras y caza-doras de más de 9.009 especies dentro de 249 géneros que se distribuyen globalmente, especialmente en los trópicos. El objetivo de esta investigación fue mejorar el conocimiento de la ocurrencia de la familia Crabronidae en México a partir de una revisión de la literatura científica y aporta datos sobre especies de importancia en el control biológico. Los resultados revelaron que en México se han reportado 769 especies que representan 99 géneros. Con la revisión actual, se identificó que el 8.5% de las especies conocidas a nivel mundial están presentes en el país y la lista de especies se incrementó en un 32%. Los géneros *Cerceris* y *Trypoxyylon* tienen el mayor número de especies, 130 y 58 especies, respectivamente. Los estados mexicanos de Baja California Norte, Sonora y Tamaulipas reportaron el mayor número de especies de Crabronidae. Se agregaron un total de 211 especies a la lista de Crabronidae en México. Se reconocieron doce géneros y ocho especies de Crabronidae utilizados como agentes de control biológico en diversas plagas. Se sugiere realizar más investigaciones sobre la diversidad de este grupo de insectos, dada la importancia que los crabónidos tienen como depredadores y enemigos naturales de insectos plaga que afectan cultivos de importancia económica.

Palabras clave: Avispas, *Cerceris*, depredadores, distribución.

Introduction

Mexico is a shelter of 10 to 12% of the total species reported in the world and is one of the main mega-diversity countries, along with Brazil, Peru, Colombia, Indonesia, and China (Toledo 1988; Mittermeier and Goetsch 1992; González 2006; SEMARNAT 2008; Sarukhán *et al.* 2009; Martínez-Meyer *et al.* 2014). Such diversity includes an estimated 125,585 species of invertebrates, most of which are insects (SEMARNAT 2008). According to Morón and Valenzuela-González (1993) and Luis *et al.* (2000), 110,000 species of insects have been reported in Mexico, which involves 6,000 species of Hemiptera, 20,000 species of Diptera, 35,000 species of Coleoptera,

and between 20,000 and 30,000 species of Hymenoptera (Michener 2000; Griswold *et al.* 2006).

The order Hymenoptera includes many species of ecological and economic importance (Fernández 2000; Della 2003; Guzmán 2014; Myartseva *et al.* 2014, 2015). Predatory wasps included in the superfamily Apoidea are distinguished because of their diversity of habits and behavioral complexity, ranging from solitary to eusocial species. Apoidea are advanced aculeates considered the most economically important insect group in the biological control (Brothers and Finnimore, 1993) and pollinators. The taxonomy of Apoidea is complicated due to the size of the insects, the relative scarcity of literature in some regions of the world, and the morphological similarity among closely related species (Ruiz *et al.* 2010).

The taxonomical criteria to separate groups vary according to different authors. For example, Bohart and Menke (1976) mentioned 11 subfamilies within the family Sphecidae, but further studies have elevated these subfamilies to the family level within the superfamily Apoidea (Gauld and Bolton, 1988). Finnimore and Michener (1993) classified these wasps into nine families grouped in the series Spheciformes, whereas Melo (1999) proposes four families, plus the family Apidae within the superfamily Apoidea. These four families are Ampulicidae, Crabronidae, Heterogynaidae, and Sphecidae, within the sphecid wasps (Spheciformes) (Melo 1999), now apoid wasps. Moreover, Pulawski (2021) considered only the families Ampulicidae, Apidae, Crabronidae, Heterogynaidae and Sphecidae.

About 8,000 species are known in Ampulicidae, Crabronidae, Heterogynaidae, and Sphecidae (Bohart and Menke 1976; Finnimore and Michener 1993). In the family Crabronidae, Pulawski (2022) cited about 9,106 species representing 249 genera and eight subfamilies, mainly from the tropics. The family Crabronidae represents 89% of the species present in the Americas from the apoid wasps, followed by Sphecidae (8%) and Ampulicidae with 2% (Table 1). The family Heterogynaidae have only been recorded in the Old World (Amarante, 2006; Pulawski, 2022).

Previous studies of Crabronidae in Mexico recorded 558 species and 80 genera of crabronids (Horta *et al.* 2013), whereas Ruiz *et al.* (2002) listed 463 species, and Amarante (2002) reported 511 species. Such studies indicate that the

families Crabronidae, Sphecidae, and Ampulicidae account for 8% of the species distributed in the Americas, highlighting Crabronidae as the most diverse family. This study aimed to update the number of species of the family Crabronidae in Mexico and provide data on species with potential use as biological control agents.

Materials and methods

A review of the available literature on the occurrence of the family Crabronidae in Mexico was conducted. This review included revising books, documents, and publications of scientific bases in the Journal Citation Reports (JCR), Latindex, and Scielo. Also, it involved other sources, such as reports of local studies in Mexico and databases available worldwide (Scullen 1972; Bohart and Menke 1976; Bohart 1990, 1993, 1997, 2000; Antropov 1999; Ohl 1999; Leclercq 2000, 2002a, 2002b, 2007; Fernandez 2001; Amarante 2002, 2005; Finnimore 2002; Ruiz and Coronado 2002; Ruiz *et al.* 2002; Genaro 2006; Holliday and Coelho 2006; Nemkov 2006; Horta *et al.* 2007, 2013; Holliday *et al.* 2009; González *et al.* 2010; Looney *et al.* 2014; Vanoye-Eligio *et al.* 2014; Cancino *et al.* 2015; Rosa 2015; Ruiz-Cancino 2015; Vanoye-Eligio *et al.* 2015; Mokrousov and Popov 2016; Discovery Life 2018; GBIF 2018; Pulawski 2022; Rosa and Melo 2020). The main criterion for the selection of species was that the record for Mexico is determined at the species level and preferably the place of collection in the different investigations. Any omission of relevant literature available was not intentional. In this study, we followed the classification of Melo (1999).

Results

According to the reviewed literature, a total of 769 species representing 99 genera of the family Crabronidae are reported in Mexico (Table 2, Supplementary File). The subfamilies Crabroninae and Philanthinae exhibited the highest numbers of species, with 338 and 187 species, respectively. The genera *Cerceris* and *Trypoxylon* were the most diverse with 130 and 58 species, respectively (Table 3). Baja California Norte, Sonora, and Tamaulipas represented the most diverse Mexican states with 144, 125, and 106 species, respectively (Fig. 1).

Table 1. Number of species of the superfamily Apoidea including the family Crabronidae reported in Mexico, the Nearctic and Neotropical regions, and worldwide (Bohart and Menke 1976; Finnimore and Michener 1993; LaSalle and Gauld 1993; Wasbauer 1995; Fernández 2000, 2002; Ruiz *et al.* 2002; González 2006; Pitts *et al.* 2006; Horta *et al.* 2007, 2009, 2013; Morrone and Márquez 2008; Pulawski 2022; Orr *et al.* 2021).

Superfamily	Family	World	Nearctic	Neotropical	Mexico
Apoidea	Sphecidae	787	102	159	87
	Crabronidae	9,106	781	1,516	558
	Ampulicidae	202	4	20	3
	Apidae	20505	6273	5929	1878

Table 2. Number of genera and species of the family Crabronidae reported from Mexico.

Subfamily	Genera	Species
Astatinae	4	26
Bembicinae	39	170
Crabroninae	40	338
Pemphredoninae	11	48
Philanthinae	5	187
Total	99	769

Table 3. Number of species of the family Crabronidae reported from Mexico and organized by subfamily and genus.

Subfamily	Genera	Species
Astatinae	<i>Astata</i> Latreille, 1796	15
	<i>Dryudella</i> Spinola, 1843	6
	<i>Diploplectron</i> W. Fox, 1893	4
	<i>Uniplectron</i> F. Parker, 1966	1
Bembicinae	<i>Bembix</i> Fabricius, 1775	19
	<i>Glenostictia</i> Gillaspay, 1962	16
	<i>Hoplisoides</i> Gribodo, 1884	16
	<i>Pseudoplisus</i> Ashmead, 1899	13
	<i>Steniola</i> Say, 1837	11
	<i>Bicyrtes</i> Lepeletier, 1845	9
	<i>Stictiella</i> J. Parker, 1917	9
	<i>Clitemnestra</i> Spinola, 1851	6
	<i>Mellinus</i> Fabricius, 1790	6
	<i>Microbembex</i> Patton, 1879	6
	<i>Stictia</i> Illinger, 1807	6
	<i>Microstictia</i> Gillaspay, 1963	5
	<i>Epinysson</i> Pate, 1935	4
	<i>Psammaletes</i> Pate, 1936	4
	<i>Bembecinus</i> A. Costa, 1859	3
	<i>Didineis</i> Wesmael, 1852	3
	<i>Sphecius</i> Dahlbom, 1843	3
	<i>Stizoides</i> Guérin-Méneville, 1844	3
	<i>Zanyssson</i> Rohwer, 1921	3
Crabroninae	<i>Alysson</i> Panzer, 1806	2
	<i>Argogorytes</i> Ashmead, 1899	2
	<i>Harpactus</i> Shuckard, 1837	2
	<i>Sagenista</i> R. Bohart, 1967	2
	<i>Xerostictia</i> Gillaspay, 1963	2
	<i>Arigorytes</i> Rohwer, 1912	1
	<i>Dienoplus</i> W. Fox 14	1
	<i>Foxia</i> Ashmead, 1898	1
	<i>Hapalomellinus</i> Ashmead, 1899	1
	<i>Leiogorytes</i> R. Bohart, 2000	1
	<i>Megistommum</i> Schulz, 1906	1
	<i>Metanyssson</i> Ashmead, 1899	1
	<i>Neoplisus</i> R. Bohart, 1967	1
	<i>Nyssson</i> Latreille, 1802	1
	<i>Oryttus</i> Spinola, 1836	1
	<i>Rubrica</i> J. Parker, 1929	1
	<i>Saygorytes</i> Nemkov 2007	1
	<i>Stenogorytes</i> Schrottky, 1911	1
	<i>Stizus</i> Latreille, 1802	1
	<i>Tanyoprymnus</i> Cameron, 1905	1
Philanthinae	<i>Trypoxyylon</i> Latreille, 1796	58
	<i>Tachysphex</i> Kohl, 1883	43
	<i>Oxybelus</i> Latreille, 1796	31
	<i>Liris</i> Fabricius, 1804	27
Pemphredoninae	<i>Crossocerus</i> Lepeletier & Brullé, 1835	26

Subfamily	Genera	Species
...	<i>Tachytes</i> Panzer, 1806	23
	<i>Rhopalum</i> Stephens, 1829	18
	<i>Ectemnius</i> Dahlbom, 1845	16
	<i>Solierella</i> Spinola, 1851	16
	<i>Crabro</i> Fabricius, 1775	8
	<i>Entomognathus</i> Dahlbom, 1844	8
	<i>Larropsis</i> Patton, 1892	7
	<i>Pison</i> Jurine In Spinola, 1808	7
	<i>Enoplolindeni</i> Rohwer, 1911	6
	<i>Belomicrus</i> A. Costa, 1871	5
	<i>Pleneculus</i> W. Fox, 1893	5
	<i>Anacrabro</i> Packard, 1866	4
	<i>Lestica</i> Billberg, 1820	4
	<i>Foxita</i> Pate, 1942	2
	<i>Huavea</i> Pate, 1948	2
	<i>Larra</i> Fabricius, 1793	2
	<i>Moniaecea</i> Ashmead, 1899	2
	<i>Aulacophilus</i> F. Smith, 1869	1
	<i>Aphilanthops</i> Patton, 1880	1
	<i>Aulacophilus</i> F. Smith, 1869	1
	<i>Bothynostethus</i> Kohl, 1884	1
	<i>Chimila</i> Pate, 1944	1
	<i>Holcorrhopalum</i> Cameron, 1904	1
	<i>Leerenierus</i> Leclercq, 1977	1
	<i>Lindenius</i> Lepeletier & Brullé, 1835	1
	<i>Lyroda</i> Say, 1837	1
	<i>Miscophus</i> Jurine, 1807	1
	<i>Nitela</i> Latreille, 1809	1
	<i>Pae</i> Pate, 1944	1
	<i>Parataruma</i> Kimsey, 1982	1
	<i>Pisonopsis</i> W. Fox, 1893	1
	<i>Podagritus</i> Spinola, 1851	1
	<i>Prosopigastra</i> Costa, 1867	1
	<i>Taruma</i> Pate, 1944	1
	<i>Tracheliodes</i> A. Morawitz, 1866	1
	<i>Pluto</i> Pate, 1937	17
Pemphredoninae	<i>Pseneo</i> Malloch, 1933	6
	<i>Stigmus</i> Panzer, 1804	6
	<i>Psen</i> Latreille, 1796	4
	<i>Mimesa</i> Shuckard, 1937	3
	<i>Psenulus</i> Kohl, 1897	3
	<i>Spilomena</i> Shuckard, 1838	3
	<i>Ammoplanops</i> Gussakovskij, 1931	2
	<i>Incastigmus</i> Finnimore, 1995	2
	<i>Polemistus</i> Saussure, 1892	1
	<i>Pulverro</i> Pate, 1937	1
Philanthinae	<i>Cerceris</i> Latreille, 1802	130
	<i>Eucerckeris</i> Cresson, 1865	36
	<i>Philanthus</i> Fabricius, 1790	15
	<i>Clypeadon</i> Patton, 1897	3
	<i>Trachypus</i> Klug, 1810	3




Figure 1. Numbers of Crabronidae species by state in Mexico.

Regarding the agroecological importance, the genera *Astata*, *Bicyrtes*, *Cerceris*, *Larrá*, *Liris*, *Oxybelus*, *Rubrica*, *Solierella*, *Tachysphex*, and *Tachytes*, which include species that are used as biological control agents, are reported in Mexico. Also, at species level, seven species (*Astata unicolor* Say, 1824, *Cerceris californica* Cresson, 1865, *C. dilatata* Spinola, 1841, *C. fumipennis* Say, 1837, *C. grandis* Banks, 1913, *Solierella blaisdelli* (Bridwell, 1920) and *S. peckhami* (Ashmead, 1897)) have been recorded in the country.

Discussion

The 769 species of Crabronidae reported from Mexico account for 8.5% of the total species reported worldwide (Pulawski 2022). Our report of 769 species also represents a 32% increase in the number of species previously provided by Horta *et al.* (2013), who listed 558 species and 80 genera. Other previous reports were 511 species by Amarante (2002) and 463 species by Ruiz *et al.* (2002). Also, the study provides evidence of 11 genera and seven species present in Mexico that are recognized as biological control agents.

At the subfamily level, the Crabroninae had the most significant number of species in Mexico (338 species) (Tables 2, 3, Supplementary File), which coincides with other studies worldwide (Ruiz *et al.* 2002; Horta *et al.* 2013; Pulawski 2022). According to Pulawski (2022), the diversity ranking of subfamilies of Crabronidae places Crabroninae as the most

diverse subfamily, followed by Bembicinae and Philanthinae. However, in our review, Philanthinae and Bembicinae were the second and third most diverse subfamilies, respectively. Five genera (*Cerceris* Latreille, *Eucerceris* Cresson, *Philanthus* Fabricius, *Clypeadon* Patton, and *Trachypus* Klug) and 187 species comprised the subfamily Philanthinae. *Cerceris* was the most diverse genus with 70% of the species in that subfamily and represented 24% of the total species of Crabronidae. Such results are consistent with Horta *et al.* (2013), who reported *Cerceris* and *Trypoxylon* as the most representative genera of the family Crabronidae. Moreover, Vanoye *et al.* (2015) stated that *Liris* and *Tachytes* genera are also representatives of the family in Mexico. In this study, however, 36 of 99 genera are associated with only one species. In terms of species richness, the rarity of species within genera may suggest many unknown species inhabiting different regions of Mexico.

The states of Baja California Norte, Sonora, and Tamaulipas in the north of Mexico had the highest number of species with 144, 125, and 106 species, respectively. One explanation of this might be that the number of studies on Crabronidae is higher in this region than in other states of Mexico. In addition, another possible determining factor in the number of species of this family is the vegetation present in these states of the country, which have large plains and plains with hills forming part of the Sonoran and Chihuahuan desert (Gutiérrez-Ruacho *et al.* 2012). In these states the vegetation is

varied, being the transition between the mountain vegetation and the vegetation of arid zones (Hernández *et al.* 2005). Toledo (1995) mentioned that these transition areas between various types of vegetation are rich in biodiversity. In addition to the above, the dominant vegetation in these areas is a desert scrub, which has open spaces that favor the abundance and richness of the family Crabronidae (Rzedowski and Huerta 1994; Comas *et al.* 2009; SEMARNAT 2011).

This vegetation is characterized by thorny vegetation such as cactus and bromeliads and shrubs that ranged from 3 to 5 m high and in semi-desert pasture. Such environmental conditions provide open spaces preferred by wasps, as observed by Comas *et al.* (2009). Indeed, the species diversity of Crabronidae declines in closed or canopied habitats. Hymenoptera associated with open environments exploits the herbaceous plants and shrubs to provide food and nesting resources. Candia and MÁyida (2005) found that structural differences between the arboreal strata and shrub vegetation influence the abundance and species richness of Crabronidae. In this regard, Genaro (2009) reported that nests of species of *Cerceris* are constructed in clay and compact soils without vegetation or sometimes with isolated low vegetation, as noted with *C. victori* (Genaro, 2009).

Of the 769 species reported from Mexico, records of 487 species (63%) provide information on state locations where they were collected, while reports on the remaining 282 species (27%) did not indicate specific locality data (Supplementary File). Overall, this review addressed the current knowledge of the diversity of species and its distribution of Crabronidae species in Mexico.

Regarding the use of insects in the natural control of pest insects, in agriculture, one of the objectives is to maintain biodiversity and promote the regulation of the abundance of harmful organisms, through predation, parasitoidism, parasitism and competition between species to reduce damage to crops (Brito and Camacho 2007). Agricultural and forest crops and other plant species of interest to humans offer excellent opportunities to develop integrated pest management programs, since there is a diversity of harmful and beneficial insect species in them that are in ecological balance and with healthy populations (León 2005). Pest insects limit the production of different crops. Ayala *et al.* (2013) mentioned that the damage they cause in crops is due to different environmental factors, the phenology of the crop and the habits of the pest insect, including the biological characteristics of the pest (Reséndiz *et al.* 2016). In biological control, parasitoids are the most widely used natural enemies of insect pests, as they play a fundamental role as natural regulators. Parasitoid wasps (Hymenoptera) constitute one of the most diverse orders of the class Insecta, which are used as natural control against insect pests (Llave 2015). Regarding predatory wasps, these have acquired great importance as predators of harmful organisms that affect agricultural crops, due to the pedotrophic behavior of these wasps, which includes several species of insects, caterpillars and spiders, and their nesting habits are variable, constructing them on different surfaces; in addition, there are wasps with solitary and social habits, which are considered to have important ecological characteristics which can help design conservation strategies and use in biological control (Brito and Camacho 2007; Meléndez-Ramírez *et al.* 2021). Hence, the use of Crabronidae species in the biological control of pests in forestry and agriculture has been important, since at least 22 cases have been documented for this

purpose worldwide. In the subfamily Philanthinae, 10 species have been recorded (*Cerceris arenaria* (Linnaeus, 1758); *C. californica*; *C. cerverae* Giner-Mari, 1941; *C. completa* Banks, 1919; *C. dilatata*; *C. fumipennis*; *C. grandis*; *C. jeiti* Genaro, 2009; and *C. victori*) (Careless 2009; Genaro 2009; Careless *et al.* 2014; Looney *et al.* 2014; Polidori 2014; Klingeman 2015) that help in the control of pest species of Brupestidae (Coleoptera), which mainly affect wood. Five species and four genera of Crabroninae have been reported (*Larra bicolor* Fabricius, 1804; *Oxybelus fritzi* Bohart, 1993; *O. platensis* Brèthes, 1901; *Solierella blaisdelli*; *S. peckhami*; *Holotachysphex*; *Liris*; *Tachysphex* and *Tachytes*) which have been observed to include species of Hemiptera and Orthoptera (*Nysius raphanus* Howard; *N. tenellus* Barber; *Gryllotalpa* Latreille; *Scapteriscus* Scudder) in their pedotrophic diet, which affect different agricultural crops (Frank *et al.* 1995; Capinera and Leppa 2001; Seyoum and Pulawski 2001; Arévalo and Frank 2005; Ferreira *et al.* 2014; Polidori 2014; Gallardo-Granados *et al.* 2016; Carabali *et al.* 2017; Acosta 2018; De Carvalho *et al.* 2018; Gómez-Agudelo and Romero-Zúñiga 2018; Rial 2019; Sáenz-Romo *et al.* 2019). For Bembicinae two species have been recorded (*Bicyrtes quadrifasciata* (Say) and *Rubrica surinamensis* De Geer, 1778) preying on pest species that attack fruit and vegetable crops (Quispe 2015; Lara *et al.* 2016; Biddinger and Joshi 2017; Lowenstein *et al.* 2018). The subfamilies Astatininae and Pemphredoninae are each recorded with a species (*Astata unicolor* Say, 1824, and *Pemphredon* sp., respectively) considered as natural predators of pest insects in agricultural crops.

In Mexico, of the crabronid species for the purposes of induced or natural biological control that have been documented, seven species belonging to three genera have been recorded (*Astata* Latreille, 1796; *Solierella* Spinola, 1851; *Cerceris* Latreille, 1802), being the species *Cerceris californica californica* Cresson, 1865, with the greatest distribution in the country (11 states), mainly in the northeastern and northwestern states, and the same as the rest of the species. It is important to mention that the states of these regions are part of northern Mexico, where agriculture have managed to expand during the 20th century in millions of hectares with sophisticated irrigation systems, being a multiregional space that occupies about 60% of the Mexican territory and whose main characteristic is the aridity of the soil (Cerutti 2015), which is one of the determining factors in the richness and abundance of the family Crabronidae (Candia and MÁyida 2005).

Conclusions

The results suggest that crabronids are a group of wasps with high species richness in Mexico, which is promoted by the combination of Nearctic and Neotropical elements that characterize this part of the Americas. This work added a total of 211 species to the Crabronidae species list and recorded seven species that have been used in the natural biological control of pests that affect the forestry and agricultural sector; however, given the ecological and economic importance of this group of insects, more research is needed on the diversity and biology of these wasps. Finally, it is important to highlight that despite the efforts that the scientific community worldwide makes by proposing species of insects as natural pest control agents, currently the main method of insect pest control is based on the use of chemical insecticides, which are of immediate action, because they have been effectively adapted.

However, the use of broad-spectrum chemical pesticides negatively affects both beneficial and harmful insects, including those with great potential in natural pest control.

Literature cited

- ACOSTA, M. A. 2018. Fauna benéfica asociada al cultivo orgánico de tomate (*Solanum lycopersicum*) en el fundo de la Universidad Nacional Agraria La Molina. Tesis de Maestría. Universidad Nacional Agraria la Molina. Lima, Perú. 109 p. <http://repositorioslatinoamericanos.uchile.cl/handle/2250/1370566>
- AMARANTE, S. T. P. 2002. A synonymic catalog of the Neotropical Crabronidae and Sphecidae (Hymenoptera: Apoidea). *Arquivos de Zoología* 37 (1): 1-139. <https://doi.org/10.11606/issn.2176-7793.v37i1p1-139>.
- AMARANTE, S. T. P. 2006. Familia Crabronidae. Introducción a los Hymenoptera de la Región Neotropical. Bogotá DC, Sociedad Colombiana de Entomología y Universidad Nacional de Colombia 457-470.
- AMARANTE, S. T. P. 2005. Addendum and corrections to a synonymic catalog of Neotropical Crabronidae and Sphecidae. *Papéis Avulsos de Zoología* 45 (1): 1-18. <https://doi.org/10.1590/S0031-10492005000100001>.
- ANTROPOV, A. V. 1999. Digger wasps of the genus *Aulacophilus* (Hymenoptera, Sphecidae, Trypoxylini). *Entomological Review* 79 (3): 332-343.
- AREVALO, H. A.; FRANK, J. H. 2005. Nectar sources for *Larra bicolor* (Hymenoptera: Sphecidae), a parasitoid of *Scapteriscus mole* crickets (Orthoptera: Gryllotalpidae), in northern Florida. *Florida Entomologist* 88 (2): 146-151. [https://doi.org/10.1653/0015-4040\(2005\)088\[0146:NSFLBH\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2005)088[0146:NSFLBH]2.0.CO;2)
- AYALA, R. O.; NAVARRO, F.; VIRLA, E. G. 2013. Evaluation of the attack rates and level of damages by the fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae), affecting corn-crops in the northeast of Argentina. *Revista de la Facultad de Ciencias Agrícolas* 45 (2): 1-12. <http://hdl.handle.net/11336/971>.
- BIDDINGER, D. J.; JOSHI, N. K. 2017. First report of native *Astata unicolor* (Hymenoptera: Crabronidae) predation on the nymphs and adults of the invasive brown marmorated stink bug (Hemiptera: Pentatomidae). *Florida Entomologist* 100 (4): 809-812. <https://doi.org/10.1653/024.100.0408>.
- BOHART, R. M.; MENKE, A. S. 1976. Sphecid wasp of the world. A generic revision. University of California Press. 695 pp.
- BOHART, R.M. 1990. New species and a key to North American *Soliarella* in the inermis group (Hymenoptera: Sphecidae: Larinae: Mischophini). *Psyche* 97(3-4): 229-240.
- BOHART, R. M. 1993. A synopsis of Central American and Caribbean *Oxybelus* (Hymenoptera, Sphecidae). *Insecta Mundi* 7 (3): 159-168.
- BOHART, R. M. 1997. A review of the genus *Hoplisoides* Gribodo (Hymenoptera: Sphecidae: Gorytini) in North America. *Proceedings of the Entomological Society of Washington* 99 (4): 645-660.
- BOHART, R. M. 2000. A review of Gorytini in the Neotropical Region (Hymenoptera: Sphecidae: Bembicinae). Revisión de Gorytini (Hymenoptera: Sphecidae: Bembicinae) en la región neotropical. *Contributions on Entomology International* 4 (2): 113-259.
- BRITO, Y. M.; CAMACHO, E. R. . 2007. Las avispas. Su papel en el control biológico de organismos nocivos. Instituto de Investigaciones de Sanidad Vegetal. Agricultura Orgánica 1-2.
- BROTHERS, D. J.; FINNAMORE, A. T. 1993. Superfamily Vespoidea. pp. 161-278. In: GOULET, H.; HUBER, J. T. (eds.), *Hymenoptera of the World: An identification guide to families*. Research Branch Agric. Canada Publ., 1894/E. 680 p. https://www.researchgate.net/profile/Denis-Brothers/publication/259840899_Superfamily_Vespoidea/links/56f4f6ce08ae7c1fda2d80d3/Superfamily-Vespoidea.pdf.
- CANCINO, E. R.; KASPARYAN, D. R.; BLANCO, J. M. C.; MYARTSEVA, S. N.; TRJAPITZIN, V. A.; AGUILAR, S. G. H.; JIMÉNEZ, J. G. 2015. Himenópteros de la Reserva "El Cielo", Tamaulipas, México. *Dugesiana* 17 (1): 53-71.
- CANDIA, R.; MÁYIDA, S. 2005. Efecto de borde de dos tipos de matriz sobre la estructura comunitaria de himenópteros aculeados en fragmentos de Bosque Seco Tropical en época de sequía. *Entomotropica* 20 (2): 127-204.
- CAPINERA, J. L.; LEPPA, N. C. 2001. Shortwinged mole cricket, *Scapteriscus abbreviatus* Scudder; Southern mole cricket, *Scapteriscus borellii* Giglio-Tos; and tawny mole cricket, *Scapteriscus vicinus* Scudder (Insecta: Orthoptera: Gryllotalpidae). IFAS Extension, University of Florida EENY-235. <https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.549.2504&rep=rep1&type=pdf>.
- CARABALI-MUÑOZ, A.; PINCHAO-TENGANAN, S. Y. ; LAMPREA-RODRÍGUEZ, I. M.; PEÑA-MOJICA, J. F. ; CARABALI-BANGUERO, D. J. 2016. Insectos polinizadores del aguacate (*Persea americana* Mill.) cv. Hass en Colombia. Colección Nuevo conocimiento agropecuario. Corporación Colombiana de Investigación Agropecuaria [CORPOICA], Mosquera, Colombia. 98. <http://hdl.handle.net/20.500.12324/13146>
- CARELESS, P. D. S. 2009. Biosurveillance: utilizing a beetle hunting wasp (*Cerceris fumipennis*, Hymenoptera: Crabronidae) to detect infestations of emerald ash borers (*Agrilus planipennis*, Coleoptera: Buprestidae) (Doctoral dissertation). <https://hdl.handle.net/10214/20216>.
- CARELESS, P. D. S.; MARSHALL, S. A.; GILL, B. D. 2014. The use of *Cerceris fumipennis* (Hymenoptera: Crabronidae) for surveying and monitoring emerald ash borer (Coleoptera: Buprestidae) infestations in eastern North America. *The Canadian Entomologist* 146 (1): 90-105. <https://doi.org/10.4039/tce.2013.53>.
- CERUTTI, M. 2015. La agriculturización del desierto: Estado, riego y agricultura en el norte de México (1925-1970). *Apuntes* 42 (77): 91-127. <https://doi.org/10.21678/apuntes.77.740>.
- COMAS, L.; ARNAN, X.; GRACIA, M.; RETANA, J.; BOSCH, J. 2009. Importancia del grado de madurez del bosque sobre las comunidades de himenópteros voladores y micromamíferos en el Parque Nacional de Aiguëstortes i Estany de Sant Maurici. 5º Congreso Forestal Español, Montes y Sociedad: Saber que hacer 1-10. http://secforestales.org/publicaciones/index.php/congresos_forestales/article/view/16659/16502.
- DE CARVALHO, A. S.; MOLINA-RUGAMA, A. J.; SILVA, F. E. L.; COSTA, E. M.; ARAUJO, E. L. 2018. Predator insect families associated with melon crop in a semiarid region in Brazil. *Journal of Agricultural Science (Toronto)* 10 (10): 267-273. <https://doi.org/10.5539/jas.v10n10p267>.
- DELLA, L. T. M. C. 2003. Hormigas de importancia económica en la región Neotropical. pp. 337-349. In: FERNÁNDEZ, F. (eds.), *Introducción a las Hormigas de la Región Neotropical*. Instituto de Investigación de Recursos Biológicos Alexander von Humboldt. Bogotá, Colombia, 398 pp. <http://repository.humboldt.org.co/bitstream/handle/20.500.11761/32961/978-958-8151-23-6.pdf?sequence=1>.
- DISCOVERYLIFE. (2018). Super Family Sphecoidea. <http://www.discoverylife.org/proceedings/0000/6/html/Pemphredonidae>. Last accessed 02 July 2020.B
- FERNÁNDEZ, C. 2001. Checklist of genera and subgenera of aculeate Hymenoptera of the Neotropical Region (Hymenoptera: Vespolidae). *Biota Colombiana* 2 (2): 87-130.
- FERNÁNDEZ, C. 2002. Sinopsis de las Familias y Subfamilias de Hymenoptera en La región Neotropical. Red Iberoamericana de Biogeografía y Entomología Sistemática PrIBES 2: 239-243. https://www.researchgate.net/profile/Fernando-Fernandez-4/publication/270283661_Sinopsis_de_las_familias_y_subfamilias_de_Hymenoptera_en_la_Región_Neotropical/links/54a6c4060cf256bf8bb6985d/Sinopsis-de-las-familias-y-subfamilias-de-Hymenoptera-en-la-Region-Neotropical.pdf.

- FERNÁNDEZ, F. 2000. Sistemática y filogenia de los himenópteros de la región Neotropical: Estado del conocimiento y perspectivas. PrIBES: Trabajos del Primer Taller Iberoamericano de Entomología Sistemática. Monografías Tercer Milenio 1: 211-231. http://sea-entomologia.org/PDF/M3M_PRIBES_2002/101_138_Aculeata.pdf
- FERREIRA, M. L. G.; JAHNKE, S. M.; MORAIS, R. M. D.; SILVA, G. D. S. D. 2014. Diversidad de insectos depredadores en área orizícola orgánica y de conservación, en Viamão, RS, Brasil. Revista Colombiana de Entomología 40 (1): 120-128. <https://www.proquest.com/openview/5668ddae20994dc714b933b2fbaab43c/1?pq-origsite=gscholar&cbl=60421>.
- FINNAMORE, A. T.; MICHENER, C. 1993. Serie Spheciformes. p. 668. In: Goulet H, Huber JT (eds.) Hymenoptera of the World: An Identification Guide to Families. Research Branch Agric. Canada Publ., 1894/E. Ottawa. <https://d1ied5g1xfgpx8.cloudfront.net/pdfs/35617.pdf>
- FINNAMORE, A. T. 2002. Revision of the world genera of tribe Stigmini (Hymenoptera: Apoidea: Crabronidae: Pemphredoninae), Part 2. Species of *Incastigmus* Finnimore. Journal of Hymenoptera Research 11 (1): 12-71. <https://ia600208.us.archive.org/29/items/biostor-95401/biostor-95401.pdf>.
- FRANK, J. H.; PARKMAN, J. P.; BENNETT, F.D. 1995. *Larra bicolor* (Hymenoptera: Sphecidae), a biological control agent of *Scapteriscus mole* crickets (Orthoptera: Gryllotalpidae), established in northern Florida. The Florida Entomologist 78 (4): 619-623. <https://doi.org/10.2307/3496051>.
- GALLARDO-GRANADOS, S.; SALAZAR-SOLÍS, E.; SALAS-ARAIZA, M. D.; MARTÍNEZ-JAIME, O. A. 2016. Incidencia de especies de hemípteros en fresa bajo dos sistemas de cultivo en Irapuato, Guanajuato, México. Southwestern Entomologist 41 (2): 547-560. <https://doi.org/10.3958/059.041.0223>.
- GAULD, I. D.; BOLTON, B. 1988. The Hymenoptera. Oxford University Press. British Museum, 332 pp. <https://doi.org/10.1163/187631289x00410>.
- GBIF (Global Biodiversity Information Facility). 2018. Free and open access to biodiversity data. <https://www.gbif.org/>. Last accessed 28 July 2020.
- GENARO, J. A. 2006. Checklist and distribution patterns of apoid wasps (Hymenoptera: Apoidea: Sphecidae and Crabronidae) of Cuba. Zootaxa 1171: 47-68. <https://doi.org/10.11646/zootaxa.1171.1.5>.
- GENARO, J. A. 2009. Dos especies nuevas de *Cerceris* de las Antillas Mayores (Hymenoptera: Apoidea, Crabronidae), con notas sobre su historia natural. Solenodon 8: 99-117. [http://scholar.googleusercontent.com/scholar?q=cache:7HCWOM_bGwJ:scholar.google.com/+GENARO,+J.+A.+2009.+Dos+especies+nuevas+de+Cerceris+de+las+Antillas+Mayores+\(Hymenoptera:+Apoidea,+Crabronidae\),+con+notas+sobre+su+historia+natural.+Solenodon+8:+99-117&hl=es&s_dtt=0,5](http://scholar.googleusercontent.com/scholar?q=cache:7HCWOM_bGwJ:scholar.google.com/+GENARO,+J.+A.+2009.+Dos+especies+nuevas+de+Cerceris+de+las+Antillas+Mayores+(Hymenoptera:+Apoidea,+Crabronidae),+con+notas+sobre+su+historia+natural.+Solenodon+8:+99-117&hl=es&s_dtt=0,5).
- GÓMEZ-AGUDELO, S. A.; ROMERO-ZÚÑIGA, R. I. 2018. Hymenoptera (Apoidea-Chalcidoidea-Chrysidoidea-Ichneumonoidea) asociados a siete cultivos promisorios del departamento del Quindío (Colombia). Revista de Investigaciones Carmenta 1: 23-31. <https://hdl.handle.net/11404/6760>.
- NZALEZ, H. A. 2006. Generación del catálogo de autoridad taxonómica para Hymenoptera: Aculeata no apiformes de la República Mexicana. Universidad Autónoma de Nuevo León, Facultad de Ciencias Biológicas. Informe final SNIB-CONABIO proyecto No. DE019. México, D. F., 17 pp.
- GONZÁLEZ, J. M.; ACOSTA, V. E.; SORMANI, C.; VINSON, S. B. 2010. *Melittobia digitata* Dahms (Hymenoptera: Eulophidae) and *Monodontomerus mexicanus* Gahan (Hym.: Pteromalidae) on a nest of *Trypoxylon (Trypargilum) mexicanum* (Saussure) (Hym.: Crabronidae) collected near Xalapa, Veracruz, Mexico. Acta Zoológica Mexicana 26 (1): 215-218. <https://doi.org/10.21829/azm.2010.261688>.
- GRISWOLD, T.; HANSON, P. E.; ALVES-DOS-SANTOS, I. 2006. Las abejas. Memoirs of the American Entomological Institute 77: 734-785.
- GUTIÉRREZ-RUACHO, O. G.; BRITO-CASTILLO, L.; SAHAGÚN, L. V.; TROYO-DIÉGUEZ, E. 2012. Distribución espacial de la temperatura y precipitación y su relación con la vegetación del Estado de Sonora. Revista Chapingo Serie Zonas Áridas 11 (2): 106-111. <https://www.redalyc.org/comocitar.oa?id=455545059007>.
- GUZMÁN, D. L. 2014. Diversidad de formícidos (Hymenoptera: Formicidae) en el bosque mesófilo de montaña de la reserva de la biosfera El Cielo, Tamaulipas, México. Tesis de Maestría, Colegio de Postgraduados, Montecillo, Texcoco, Edo. De México, 85 pp. <https://www.repositorionacionalcti.mx/recurso/oai:colposdigital.colpos.mx:10521/2445>.
- HERNÁNDEZ, L.; TREVIÑO-CARREÓN, J.; MORA, A.; MARTÍNEZ, M. 2005. Diversidad florística y endemismos. En: G. Sánchez; R. Dirzo y P. Reyes (Eds.) Historia Natural de la Reserva de la Biosfera el Cielo, Tamaulipas, México. México. Universidad Autónoma de Tamaulipas. Impreso en Hong Kong. Pp. 244-259.
- HOLLIDAY, C. W.; COELHO, J. R. 2006. Improved key to new world species of *Sphecius* (Hymenoptera: Crabronidae). Annals of the Entomological Society of America 99 (5): 793-798. [https://doi.org/10.1603/0013-8746\(2006\)99\[793:IKTNWS\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2006)99[793:IKTNWS]2.0.CO;2).
- HOLLIDAY, C. W.; HASTINGS, J. M.; COELHO, J. R. 2009. Cicada prey of New World cicada killers, *Sphecius* spp. (Dahlbom, 1843) (Hymenoptera: Crabronidae). Entomological News 120 (1): 1-17. <https://doi.org/10.3157/021.120.0101>.
- HORTA, V. J. V.; VANOEY, E. M.; GARCÍA, G. M. E.; CORONADO, B. J. M.; BARRIENTOS, L. L. 2013. Crabronidae (Hymenoptera) de la localidad Cañón del Novillo, Victoria, Tamaulipas, México. Acta Zoológica Mexicana 29 (2): 376-387. <http://www.scielo.org.mx/pdf/azm/v29n2/v29n2a9.pdf>.
- HORTA, V. J. V.; GARCÍA, G. M. E.; BENAVIDES, M. M. I.; CORREA, S. A. 2009. Pompilidae (Hymenoptera) de algunos Municipios del Centro y Sur de Tamaulipas, México. Acta Zoológica Mexicana 25 (1): 71-82. <https://doi.org/10.21829/azm.2009.251592>.
- HORTA, V. J. V.; DOMÍNGUEZ, P. N. O.; BARRIENTOS, L. L.; CORREA, S. A. 2007. Sphecidae y Crabronidae (Hymenoptera) de algunos municipios del centro y sur de Tamaulipas, México. Acta Zoológica Mexicana 23 (3): 35-48. <https://doi.org/10.21829/azm.2007.233584>.
- KLINGEMAN, W. E.; HANSEN, J. A.; BASHAM, J. P.; OLIVER, J. B.; YOUSSEF, N. N.; SWINK, W.; MOULTON, J. K. 2015. Seasonal flight activity and distribution of metallic woodboring beetles (Coleoptera: Buprestidae) collected in North Carolina and Tennessee. Florida Entomologist 98 (2): 579-587. <https://doi.org/10.1653/024.098.0230>.
- LARA, J.; PICKETT, C.; INGELS, C.; HAVILAND, D.; GRAFTON-CARDWELL, E.; DOLL, D.; HODDLE, M. 2016. Biological control program is being developed for brown marmorated stink bug. California Agriculture 70 (1): 15-23. <https://doi.org/10.3733/ca.v070n01p15>.
- LASALLE, J.; GAULD, I. D. 1993. Hymenoptera: their biodiversity, and their impact on the diversity of other organisms. Hymenoptera and biodiversity 1-26. <https://www.cabdirect.org/cabdirect/abstract/19931171651>.
- LECLERCQ, J. 2000. Hyménoptères sphécides crabroniens des Amériques du genre *Crossocerus* Lepetier & Brullé, 1835. Notes Fauniques de Gembloux 40: 3-75.
- LECLERCQ, J. 2002a. Les genres *Minicrabro* gen. n. et *Moniae-cera* Ashmead en Amerique (Hymenoptera: Crabronidae, Crabroninae). Notes Fauniques de Gembloux 49: 3-23.
- LECLERCQ, J. 2002b. Hyménoptères Crabronides Crabroniens des Amériques du genre *Rhopalum* Stephens, 1829. Himenópteros Crabroninae del género *Rhopalum* Stephens, 1829 de las Américas. Notes Fauniques de Gembloux 48: 3-115.

- LECLERCQ, J. 2007. Pour Lecrenierus Leclercq, 1977, genre d’Hyménoptères Crabroniens de l’Amérique Latine (Hymenoptera: Crabronidae: Crabroninae). Information on Lecrenierus Leclercq, 1977, the crabronid hymenopteran from Latin America (Hymenoptera: Crabronidae: Crabroninae). Notes Fauniques de Gembloux 60 (2): 93-98.
- LEÓN, M. 2005. La diversidad de insectos en cítricos y su importancia en los programas de manejo integrado de plagas. Manejo Integrado de Plagas y Agroecología 74: 85-93. <https://agriprofiles.agri-d.net/display/AS-pub-0FED84D5B72D12E6800374B5E5E5BC74>.
- LLAVE, A. A. C. 2015. Registro de nuevos taxones de avispas parasitoides (Hymenoptera) como enemigos naturales para el control de plagas en los cultivos de maní y ají. Agro-Ecológica 2 (1): 180-195. <http://revistas.usfx.bo/index.php/rae/article/view/103>.
- LOONEY, C.; HELLMAN, W.; WESTCOTT, R. 2014. Sampling Buprestidae (Coleoptera) in Washington state with *Cerceris californica* Cresson (Hymenoptera, Crabronidae). Journal of Hymenoptera Research 39: 83-97. <https://doi.org/10.3897/jhr.39.8026>.
- LOWENSTEIN, D. M.; ANDREWS, H.; RUDOLPH, E.; SULLIVAN, E.; MARSHALL, C. J.; WIMAN, N. G. 2018. *Astata unicolor* (Hymenoptera: Crabronidae) population in Oregon with observation of predatory behavior on Pentatomidae. Annals of the Entomological Society of America 111 (3): 122-126. <https://doi.org/10.1093/aesa/say010>.
- LUIS, A.; LLORENTE, J.; VARGAS, I.; GUTIÉRREZ, A. L. 2000. Síntesis preliminar del conocimiento de los Papilionoidea (Lepidoptera: Insecta) de México. pp. 275-285. In: MARTÍN, F.; MORRONE, J. J.; MELIC, A. (eds.) Hacia un proyecto para el inventario y estimación de la diversidad biológica en Iberoamérica. PrIBES-2000. SEA, Zaragoza, España. <https://dialnet.unirioja.es/servlet/articulo?codigo=2221337>.
- MARTÍNEZ-MEYER, E.; SOSA-ESCALANTE, J. E.; ÁLVAREZ, F. 2014. El estudio de la biodiversidad en México: ¿una ruta con dirección? Revista Mexicana de Biodiversidad 85: 1-9. <https://doi.org/10.7550/rmb.43248>.
- MELÉNDEZ-RAMÍREZ, V.; DELFÍN-GONZÁLEZ, H.; MARIQUE-SAIDE, P.; CHABLÉ-SANTOS, J.; SÉLEM-SALAS, C. 2021. Diversidad y ecología de abejas, avispas depredadoras y mariposas: aportaciones del CCBA-UADY†. Tropical and Subtropical Agroecosystems 24 (3): 1-2.
- MELO, G. A. R. 1999. Phylogenetic relationships and classification of the major lineages of Apoidea (Hymenoptera), with emphasis on crabronid wasps. Scientific Papers, University of Kansas Natural History Museum 14: 1-55. <https://www.proquest.com/openview/480e39e02003706699dd64a3a36bd979/1?pq-orignal=gscholar&cbl=18750&diss=y>.
- MICHENER, C. D. 2000. The bees of the world. Baltimore & London: J. Hopkins University Press. 913 pp. <https://doi.org/10.1002/mmzn.20020780209>.
- MITTERMEIER, R.; GOETTSCH, C. 1992. La importancia de la diversidad biológica de México. México ante los retos de la biodiversidad 57-62.
- MOKROUSOV, M. V.; POPOV, I. B. 2016. Digger wasps (Hymenoptera, Apoidea: Ampulicidae, Sphecidae, Crabronidae) of the Black Sea coast of Krasnodar Territory, Abkhazia, and adjacent areas. Entomological Review 96 (5): 559-599. <https://doi.org/10.1134/S0013873816050055>.
- MORON, M. A.; VALENZUELA-GONZÁLEZ, J. E. 1993. Estimación de la biodiversidad de insectos en México; análisis de un caso. Revista de la Sociedad Mexicana de Historia Natural 44: 303-312.
- MORRONE, J. J.; MÁRQUEZ, J. 2008. Biodiversity of Mexican terrestrial arthropods (Arachnida and Hexapoda): a biogeographical puzzle. Acta Zoológica Mexicana 24 (1): 15-41. <https://doi.org/10.21829/azm.2008.241613>.
- MYARTSEVA, S. N.; RUIZ-CANCINO, E.; CORONADO-BLANCO, J. M. 2014. Biodiversidad de Aphelinidae (Hymenoptera: Chalcidoidea) en México. Revista Mexicana de Biodiversidad 85: 379-384. <https://doi.org/10.7550/rmb.32255>.
- MYARTSEVA, S. N.; RUIZ-CANCINO, E.; CORONADO-BLANCO, J. M. 2015. Diversidad del Género *Encarsia* Förster (Hymenoptera: Chalcidoidea: Aphelinidae) en Tamaulipas, México. Encuentro Nacional de Investigación Científica y Tecnológica del Golfo de México 18. https://www.researchgate.net/profile/Juana-Coronado-Blanco/publication/271588845_Diversidad_del_genero_Encarsia_Förster_Hymenoptera_Chalcidoidea_Aphelinidae_en_Tamaulipas_Mexico/links/54cd72770cf29ca810f8354f/Diversidad-del-genero-Encarsia-Förster-Hymenoptera-Chalcidoidea-Aphelinidae-en-Tamaulipas-Mexico.pdf#page=44.
- NEMKOV, P. G. 2006. A new genus of digger wasps (Hymenoptera, Crabronidae, Bembicinae) from North and Central America. Far Eastern Entomologist 167: 1-4.
- OHL, M. 1999. A revision of *Stizoides* Guérin-Méneville, 1844: Taxonomy, phylogenetic relationships, biogeography, and evolution (Hymenoptera: Apoidea, “Sphecidae”). Zoosystematics and Evolution 75 (1): 63-169. <https://doi.org/10.1002/mmnz.19990750108>.
- ORR, M. C.; HUGHES, A. C.; CHESTERS, D.; PICKERING, J.; ZHU, C. D.; ASCHER, J. S. 2021. Global patterns and drivers of bee distribution. Current Biology 31 (3): 451-458. <https://doi.org/10.1016/j.cub.2020.10.053>.
- PITTS, J. P.; WASBAUER, M. S.; VON DOHLEN, C. D. 2006. Preliminary morphological analysis of relationships between the spider wasp subfamilies (Hymenoptera: Pompilidae): revisiting an old problem. Zoologica Scripta 35 (1): 63-84. <https://doi.org/10.1111/j.1463-6409.2005.00217.x>.
- POLIDORI, C. 2014. Himenópteros depredadores y control biológico de plagas: ¿un recurso ignorado? Entomología y Control Biológico 1-12. <https://air.unimi.it/handle/2434/801098#YWROp1VBzIU>.
- PULAWSKI, W. J. 2021. Catalog of Sphecidae. <https://www.calacademy.org/scientists/projects/catalog-of-sphecidae>. Last accessed 19 April 2022.
- PULAWSKI, W. J. 2022. Number of Extant Species of Sphecidae sensu lato. https://researcharchive.calacademy.org/research/entomology/entomology_resources/hymenoptera/sphecidae/number_of_species.pdf. Last accessed 19 April 2022.
- QUISPE, T. R. 2015. Refugios vegetales para el fomento de la entomofauna benéfica en el agroecosistema del cultivo de maíz en La Molina. Tesis de Maestría. Universidad Nacional Agraria la Molina, Lima, Perú. 181 pp. <http://repositorio.lamolina.edu.pe/handle/20.500.12996/1163>.
- RESÉNDIZ, R. Z.; LÓPEZ, S. J. A.; OSORIO, H. E.; ESTRADA D. B.; PECINA, M. J. A.; MENDOZA, C. M. C.; REYES, M. C. A. 2016. Importancia de la resistencia del maíz nativo al ataque de larvas de lepidópteros. Temas de Ciencia y Tecnología 20 (59): 3-14.
- RIAL, J. B. 2019. Artropofauna de sistemas hortícolas periurbano y rural en cultivos de tomate (*Lycopersicum esculentum* Mill.) en el Alto Valle de Río Negro y Neuquén. Bachelor’s Thesis, Universidad Nacional del Comahue. Facultad de Ciencias del Ambiente y la Salud. 183 pp.
- ROSA, B. B.; MELO, G. A. 2020. Systematics of the Neotropical species of the crabronid wasp genus *Psenulus* Kohl, 1897 (Hymenoptera: Crabronidae), with a contribution to the worldwide phylogeny of the genus. Austral Entomology 59 (3): 422-454. <https://doi.org/10.1111/aen.12487>.
- ROSA, B. B. D. 2015. Revisão taxonómica e análise filogenética das espécies neotropicais do gênero *Psenulus* Kohl, 1897 (Hymenoptera, Crabronidae). Tesis de Maestría. Universidad Federal do Paraná, 81 pp. <https://acervodigital.ufpr.br/bitstream/handle/1884/38046/R%20-%20D%20-%20BRUNNO%20BUENO%20DA%20ROSA.pdf?sequence=1&isAllowed=y>.
- RUIZ, C. E.; CORONADO, B. M. J.; VARELA, S. E.; HORTA, V. J. V. 2002. Sphecoidea. pp. 657-670. In: LLORENTE, B. J.;

- MORRONE, J. J. (eds.). Biodiversidad, Taxonomía y Biogeografía de Artrópodos de México. Vol. 3. Facultad de Ciencias de la U.N.A.M. México.
- RUÍZ, C. E.; KASPARYAN, D. R.; CORONADO, B. J. M.; MYARTSEVA, S. V.; TRJAPITZIN, V. A.; HERNÁNDEZ, A. S. G.; GARCÍA-JIMÉNEZ, J. 2010. Himenópteros de la Reserva "El Cielo", Tamaulipas, México. *Dugesiana* 17 (1): 53-71. <https://www.zin.ru/Labs/insects/hymenopt/personalia/Myartseva/pdf/2010d.pdf>.
- RUIZ, C. E.; CORONADO, B. J. M. 2002. Artrópodos terrestres de los estados de Tamaulipas y Nuevo León, México. Universidad Autónoma de Tamaulipas, Unidad Académica Multidisciplinaria Agronomía y Ciencias, Centro de Investigación y Desarrollo Agropecuario, Forestal y de la Fauna, 379 pp.
- RUÍZ-CANCINO, E. 2015. La familia Ichneumonidae (Hymenoptera) en México. *Entomología Mexicana* 2: 1-13.
- RZEDOWSKI, J.; HUERTA, L. 1994. Vegetación de México. No. 582.16 R997r Ej. 1 002135, México. Limusa, Noriega Editores, 492 pp. <https://bibliotecasibe.ecosur.mx/sibe/book/000022943>.
- SÁENZ-ROMO, M. G.; VEAS-BERNAL, A.; MARTÍNEZ-GARCÍA, H.; IBÁÑEZ-PASCUAL, S.; MARTÍNEZ-VILLAR, E.; CAMPOS-HERRERA, R.; PÉREZ-MORENO, I. 2019. Effects of ground cover management on insect predators and pests in a mediterranean vineyard. *Insects* 10 (12) 421: 1-16. <http://dx.doi.org/10.3390/insects10120421>.
- SARUKHÁN, J.; KOLEFF, P.; CARABIAS, J.; SOBERÓN, J.; DIRZO, R.; LLORENTE-BOUSQUETS, J.; ANTA, S. 2009. Capital natural de México. Síntesis: conocimiento actual, evaluación y perspectivas de sustentabilidad. México: Comisión Nacional para el Conocimiento y Uso de la Biodiversidad. (No. 333.951672 COM SIN. CIMMYT), 100 pp. <https://doi.org/10.5962/bhl.title.112130>.
- SCULLEN, H. A. 1972. Review of the genus *Cerceris* Latreille in Mexico and Central America (Hymenoptera: Sphecidae). *Smithsonian Contributions to Zoology* 110: 1-121. <https://doi.org/10.5479/si.00810282.110>.
- SEMARNAT (Secretaría del Medio Ambiente y Recursos Naturales). 2011. El Medio Ambiente en México. Ecosistemas Terrestres. http://apps1.semarnat.gob.mx/dgeia/informe_resumen14/02_ecosistemas/2_1.html. Last accessed 15 May 2020.
- SEMARNAT (Secretaría del Medio Ambiente y Recursos Naturales). 2008. Capítulo IV: Biodiversidad en México. [https://apps1.semarnat.gob.mx:8443/dgeia/informe18/tema/cap4.html](http://apps1.semarnat.gob.mx:8443/dgeia/informe18/tema/cap4.html). Last accessed 02 May 2019.
- SEYOUUM, E.; PULAWSKI, W. J. 2001. Sphecid wasps (Hymenoptera: Apoidea: Sphecidae) as potential control agents of acridid pests (Orthoptera: Acrididae) in Ethiopia. *Archives of Phytopathology & Plant Protection* 34 (5): 319-326. <https://doi.org/10.1080/03235400109383394>.
- TOLEDO, V. M. 1988. La diversidad biológica de Latinoamérica: un patrimonio amenazado. *Ambiente y desarrollo* 4 (3): 13-24. https://www.unich.edu.mx/wp-content/uploads/2014/01/BiodiversidadLatinoamerica_Toledo88.pdf.
- TOLEDO, V. M. 1995. La riqueza florística de México: un análisis para conservacionistas. En: S. Guevara, P. Moreno-Casa-Sola y J. Rzedowski (Eds.), Logros y perspectivas del conocimiento de los recursos vegetales de México en vísperas del siglo XXI. Instituto de Ecología, Xalapa 109-124.
- VANOYE-ELIGIO, M.; MELÉNDEZ-RAMÍREZ, V.; AYALA, R.; NAVARRO-ALBERTO, J.; DELFÍN-GONZÁLEZ, H. 2015. Avispas depredadoras de áreas naturales protegidas del estado de Yucatán, México. *Revista Mexicana de Biodiversidad* 86 (4): 989-997. <http://dx.doi.org/10.1016/j.rmb.2015.04.037>.
- VANOYE-ELIGIO, M.; RAMÍREZ, V.M.; AYALA, R.; NAVARRO, J.; DELFIN-GONZÁLEZ, H. 2014. Predatory Wasps (Hymenoptera) of the Yucatan Peninsula. *Southwestern Entomologist* 39 (3): 635-646. <https://doi.org/10.3958/059.039.0322>
- WASBAUER, M. S. 1995. Pompilidae. The Hymenoptera of Costa Rica 14 (3): 5221161-1170.

Origin and Funding

The present research work derives from a literature review of the species registered for Mexico of the Crabronidae family, as well as their potential use in agriculture. The project did not have financial support.

Author Contribution

MVE and VVE conceived/designed the research and wrote the manuscript. MVE, LJER and BRMC performed the literature search and data analysis. MVE, JVHV and VVE reviewed and edited the manuscript. All authors read, corrected, and approved the manuscript.

Conflict of Interest

The authors declare that they have no conflict of interest.

Ethical approval

The study was a literature review published to date, so insects were not captured at any time. Due to the above, an ethical approval was not required.