

Effect of mating on ovary maturation in *Dactylopius coccus* (Hemiptera: Coccoidea: Dactylopiidae)

Efecto de la cópula sobre la maduración del ovario en *Dactylopius coccus*
(Hemiptera: Coccoidea: Dactylopiidae)

ARTURO RAMÍREZ-CRUZ¹; CELINA LLANDERAL-CÁZARES²

¹ Ph. D. CIIDIR, Unidad Michoacán, Instituto Politécnico Nacional, CP 59510, Michoacán, México, aramirez@ipn.mx, <https://orcid.org/0000-0001-5184-6194>. ² Ph. D. Entomología y Acarología, Colegio de Postgraduados, IFIT, Campus Montecillo, Montecillo, Estado de México, México, llcelina@colpos.mx, <https://orcid.org/0000-0002-7049-2931>.

Abstract: The cochineal *Dactylopius coccus* is an industrially-important insect used to extract carminic acid (a red dye). The effect of mating on the maturation of its ovaries was determined. In order to do this, three-month old mated and virgin females were used which were counted from *Opuntia ficus indica* cladode infestation (in Montecillo, Mexico). The morphophysiological characteristics of the degree of maturation of their ovaries were evaluated. Mating was not needed to start the maturation of the ovaries, because in virgin females, a low percentage of oocytes successfully complete vitellogenesis and choriogenesis; however, mating promotes both processes, since the proportion of mature oocytes produced in mated females noticeably increases. Similarly, lack of mating in *D. coccus* was a factor that induced degeneration of the ovaries, and consequently, the resorption of their oocytes.

Keywords: Oocyte resorption, oogenesis, ovarian maturation, ovoviviparity, cochineal insects, Hemiptera, Coccoidea, Dactylopiidae.

Resumen: La cochinilla *Dactylopius coccus* es un insecto de gran importancia industrial para la extracción del colorante rojo ácido carminico. Se determinó el efecto de la cópula sobre la maduración del ovario. Para ello, se emplearon hembras copuladas y hembras vírgenes de tres meses de edad, contados desde la infestación de los cladodios de *Opuntia ficus indica* (en Montecillo, México). Se evaluaron las características morfofisiológicas del grado de madurez de sus ovarios. La cópula no fue necesaria para que diera inicio la maduración de las ovarios, ya que en las hembras vírgenes un bajo porcentaje de sus ovocitos lograron culminar adecuadamente la vitelogénesis y la coriogénesis; sin embargo, la cópula promovió ambos procesos, ya que aumentó notablemente la proporción de los ovocitos maduros que se producirán en las hembras copuladas. Del mismo modo, la falta de cópula en *D. coccus* fue un factor que indujo la degeneración de las ovarios y, en consecuencia, la reabsorción de sus ovocitos.

Palabras clave: Reabsorción del ovocito, ovogénesis, maduración ovárica, ovoviviparidad, cochinillas, Hemiptera, Coccoidea, Dactylopiidae.

Corresponding author: Arturo Ramírez-Cruz, Ph. D. CIIDIR, Unidad Michoacán, Instituto Politécnico Nacional, CP 59510, Michoacán, México, aramirez@ipn.mx, <https://orcid.org/0000-0001-5184-6194>.

Suggested citation:
RAMÍREZ-CRUZ, A.; LLANDERAL-CÁZARES, C. 2019. Effect of mating on ovary maturation in *Dactylopius coccus* (Hemiptera: Coccoidea: Dactylopiidae). Revista Colombiana de Entomología 45 (2): e7954. <https://doi.org/10.25100/socolen.v45i2.7954>

Received: 16-Aug-2017

Accepted: 21-Mar-2019

Published: 31-Dec-2019

Revista Colombiana de Entomología
ISSN (Print): 0120-0488
ISSN (Online): 2665-4385
<http://revistacolombianaentomologia.univalle.edu.co/>

Open access



BY-NC-SA 4.0
(<https://creativecommons.org/licenses/by-nc-sa/4.0/deed.es>)

Publishers:
Sociedad Colombiana de Entomología
SOCOLEN (Bogotá, D. C., Colombia)
<http://www.socolen.org.co>
Universidad del Valle (Cali, Colombia)
<http://www.univalle.edu.co/>

© 2019 Sociedad Colombiana de Entomología
- SOCOLEN y Universidad del Valle - Univalle

Introduction

Among insects, mating has a huge influence on the reproductive physiology of females. Some of the processes that can be affected by mating are, for instance, vitellogenesis, oviposition, fecundity, and fertility; however, the effect on such processes varies depending on the group of insects (Engelmann 1970; Raabe 1986). Females of the insect *Dactylopius coccus* Costa, 1829 are used to extract carminic acid, a natural red dye currently employed in several industries (Méndez-Gallegos *et al.* 2003). Their ovaries are made up of more than 400 ovarioles of the telotrophic type, where the germarium, located apically and where nutritive cells are located, is connected by the trophic cord to a single oocyte contained in the vitellarium (Ramírez-Cruz *et al.* 2008). From the molt to adult, the female of *D. coccus* has ovarioles with oocytes completely immature since it is a sinovigenic species, and from that moment onwards, maturation starts; this maturation is characterized by the progressive increase of size of the germarium and the vitellarium over several days. When the female reaches 13 days of age, its ovarioles show the first mature oocytes where the chorion is formed (choriogenesis); at this age, the germarium has already entered into a degenerative stage and tends to be resorbed. Only one gonadic cycle takes place (Ramírez-Cruz 2014).

It is known that in *D. coccus* mating is necessary for the oviposition to take place (Marín and Cisneros 1977; Pérez-Guerra and Kosztarab 1992), and it also increases the concentration of carminic acid in females (Briseño-Garzón and Llanderal 2008; Lagowska and Golan 2009). However, there is no information about the effect of mating on ovarian maturation in this species. Due to that, in order to contribute to the better understanding of the reproductive physiology of *D. coccus*, the main objective of this work was to determine how mating affects maturation of the ovary, considering the oocytes of each ovariole.

Materials and methods

Insects were reared on cladodes of *Opuntia ficus indica* (L.) Mill (Cactaceae) using the method of inverted hanging cladode proposed by Aldama-Aguilera and Llanderal-Cázares (2003). From April to July 2013, a greenhouse for rearing cochineal was used, with an average temperature of 21 °C and an average relative humidity of 50 %, at the Colegio de Postgraduados in Montecillo, Mexico, located at 19°29'N and 98°54'W, at an altitude of 2,250 mals. Females from this colony were isolated and grouped into two experimental groups: mated females, and virgin females. Although mating was not experimentally verified in the group of mated females, it was assumed that it took place because they were in the presence of males. To obtain virgin females, some of the infested cladodes were kept in isolation inside wooden cages with glass in two of their walls and organza fabric in the other two; to ensure the virginity of females, it was taken into account that the pupal stage of males lasts about 12 days (Llanderal-Cázares and Nieto-Hernández 2001), so cocoons were carefully removed as soon as they were observed, and the cages were inspected for the emergence of adult males.

To evaluate the effect of mating on maturation, from each of the experimental groups mentioned above, 20 mated females that had not yet oviposited and 20 virgin females were randomly selected; all females were three months of age, counted from the time that cladodes were infested with crawlers. To evaluate the degree of maturation of the ovarioles in both groups of females, they were fixed in Carnoy's solution for 24 hours, and they were rinsed with 70 % ethanol (Martínez 2002); to clear ovarioles, whole females were placed in a mixture of pure glycerin - 70 % ethanol (1:2) (for three days at least) until they were dissected. Finally, under the stereomicroscope, both ovaries of each female were extracted, and all their ovarioles were counted and classified according to their degree of maturation, and some morphophysiological characteristics described for ovariole maturation of *D. coccus* (Ramírez-Cruz and Llanderal-Cázares 2013; Ramírez-Cruz 2014).

Results

The ovarioles of both mated and virgin females were morphophysiological characterized and grouped in at least four maturation stages (Fig. 1): Stage I, which included ovarioles with oocytes in previtellogenesis or vitellogenesis. Stage II, which included ovarioles with the germarium in the middle of the process of being resorbed, and thus, it was no longer functional. Stage III, which showed ovarioles with the germarium completely or almost completely resorbed. Stage IV, comprised degenerating ovarioles or ovarioles being




Figure 1. Schematic representation of the ovarioles morphology of the stages of maturation I – IV taken into account, in three-month-old adult females of *Dactylopius coccus*. The diagrams represent the average sizes of the ovarioles from each stage. g, germarium; v, vitellarium. Scale bar: 200 μ m.

resorbed, with a high degree of resorption of the germarium and its oocyte in the vitellarium.

Stage I. In both mated and virgin females, ovarioles in stage I (oocytes in previtellogenesis or vitellogenesis) were found (Figs. 2A, 2B, respectively); however, in mated females, only about 10 % of their ovarioles (52.1 on average) were in stage I, while in virgin females, the percentage of ovarioles in this stage was almost 40 % (167.2 ovarioles on average) (Fig. 3).

Stage II. Both mated and virgin females had fewer ovarioles in stage II (with a degenerating germarium) than in stage I; in mated females, just a little more than 2 % of their ovarioles (12.75 ovarioles on average) were in this stage, while in virgin females this number was closer to 10 % (38.2 ovarioles on average) (Fig. 3). However, the most important difference between these two groups was the maturation stage of the oocytes, since in mated females, ovarioles in stage II showed from oocytes that were almost mature until eggs with embryos in midblastula stage (Fig. 2C), while virgin females had ovarioles in stage II with either oocytes close to maturity or mature oocytes where choriogenesis had already concluded (Fig. 2D).

Stage III. In mated females, almost 87 % of their ovarioles (441.2 ovarioles on average) were in stage III (germarium completely or almost completely resorbed) (Fig. 3), and they had eggs with embryos from the stage of late blastula until pre-nymphs or completely developed nymphs (Fig. 2E). On the other hand, the percentage of ovarioles in this stage in virgin females was only slightly above 10 % (45.15 ovarioles




Figure 2. Ovarioles in different maturation stages, in three-month-old mated and virgin females of *Dactylopius coccus*. **A.** Stage I, mated; in vitellogenesis. **B.** Stage I, virgin; in vitellogenesis. **C.** Stage II, mated; the vitellarium (v) contains a mature oocyte. **D.** Stage II, virgin; the vitellarium (v) contains an oocyte without chorion. **E.** Stage III, mated; two early embryos can be observed (above) and a completely developed nymph (below). **F.** Stage III, virgin; mature oocyte. **G.** Stage IV, virgin; the germarium (g) and traces of yolk (asterisk) can still be observed in the oocyte. **H.** Three ovarioles from a virgin female where an ovariole (n) can be observed next to two ovarioles in different stages of resorption (r); in the latter, traces of yolk can still be seen in the center of both oocytes. g, germarium; v, vitellarium. Scale bar: 100 μ m.

on average) (Fig. 3); however, these ovarioles contained only mature oocytes and they never had embryos (Fig. 2F). Thus, it is important to notice that the ovarioles of virgin females can adequately complete their maturation and develop oocytes with chorion, although in low numbers.

Stage IV. In mated females, the number of ovarioles in stage IV (degenerating or being resorbed) was insignificant, about 0.16 % (two ovarioles on average), whereas in female virgins, 41.68 % of their ovarioles (193.4 ovarioles on average) were in the stage of degeneration (Fig. 3). In both cases, ovariole resorption occurred only in those that contained oocytes at the end of vitellogenesis or close to maturity, but that had not started choriogenesis yet. In most of them it was possible to observe the germarium, although some only presented the vitellarium; both structures showed evident signs of degeneration and resorption, followed by shrinking of the ovariole (Fig. 2G). In the vitellarium, oocytes also showed different degrees of yolk degradation (Fig. 2H).

Discussion

In this study, the fact that virgin females of *D. coccus* presented ovarioles with oocytes in vitellogenesis shows that this process is independent of mating; however, in these females the process seems to be delayed or even inhibited, because at three months after crawler emergence, about 40 % of their

ovarioles were still in stage I of maturation; this contrasts with mated females, where only about 10 % of their ovarioles were in that stage, and about 87 % were in stage III. This indicates that, although mating in *D. coccus* is not necessary for their oocytes to start vitellogenesis, it is necessary to ensure the adequate conclusion of the process.

In the same way, the fact that in virgin females we found ovarioles in stages II and III with oocytes with chorion indicates that these females are able to complete maturation normally, albeit in lower proportions than mated females. This demonstrates that, although mating in *D. coccus* is not necessary for the complete maturation of its oocytes, it is necessary to increase the number of mature oocytes that will be produced.

Similarly to what was found in this study with *D. coccus*, there are other species in Hemiptera where normal maturation of oocytes is independent of mating, since virgin females produce mature oocytes, although in lower numbers than in mated females; some examples are found in the Miridae *Adelphocoris lineolatus* (Goeze, 1778) (Masner 1966) and *Helopeltis antonii* Signoret, 1858 (Siswanto *et al.* 2009), in the Triatominae *Triatoma infestans* (Klug, 1834) (Asin and Crocco de Ayerbe 1992) and *Triatoma protracta* (Uhler, 1894) (Mundall 1978), and in Pentatomidae *Perillus bioculatus* (Fabricius, 1775) (Adams 2000 a, b), *Podisus maculiventris* (Say, 1832) (De Clercq and Degheele 1997), *Podisus nigrispinus* (Dallas, 1851) (Soares *et al.* 2011) and




Figure 3. Average percentage of the number of ovarioles (\pm SEM) from the four maturation stages that were considered, in mated (black bars) and virgin (gray bars) three-month-old *Dactylopius coccus* females.

Nezara viridula (L., 1758) (Fortes *et al.* 2011). However, in contrast with the latter, there are other Hemiptera where females are incapable of developing any mature oocyte unless mating takes place; such is the case of *Dindymus versicolor* (Herrich-Schaeffer, 1853) (Pyrrhocoridae) (Friedel 1974), *Diaphorina citri* Kuwayama, 1908 (Psyllidae) (Dossi and Consoli 2010) and *Anthocoris tomentosus* Péricart, 1971 (Anthocoridae) (Horton *et al.* 2005). About that, it is known that in Hemiptera, mating affects oocyte maturation by activating the *corpora allata* or maintaining their activity for longer, in order to synthesize juvenile hormone (Davey 1997). Although the hormonal mechanism that regulates oocyte maturation in *D. coccus* is unknown, the fact that only a small percentage of the oocytes of virgin females completed vitellogenesis and choriogenesis normally (mature oocytes in stages II and III) in this study, could suggest that the *corpora allata* of such females are active only for a short period of time (compared with mated females) before being deactivated; therefore, the ovarioles whose oocytes had started but not concluded vitellogenesis would start degenerating or would be resorbed (stage of maturation IV). Something similar has been observed in *T. protracta* (Mundall and Engelmann 1977); however, profound studies are required to really know the endocrine mechanism involved in oocyte maturation in *D. coccus*.

On the other hand, oocyte resorption in insects is a phenomenon produced by several factors, both internal and external that occurs mainly under adverse conditions (Bell and Bohm 1975). In this study, the absence of mating in *D. coccus* induced high levels of ovariole and oocyte resorption (stage of maturation IV), and affected mainly vitellogenesis, since three months after emergence, little more than 40 % of the ovarioles of virgin females started to be resorbed before their oocytes completed that phase and began synthesizing the chorion. In contrast, in mated females, only 0.16 % of the ovarioles showed that phenomenon. Ovariole resorption in *D. coccus* was morphologically characterized by the degeneration of the germarium, vitellarium and its respective oocyte, with the consequent degradation of the yolk, and although the mechanism of yolk degradation in *D. coccus* is unknown, it is probably degraded and released into the hemolymph, similarly to what happens in the bug *Plautia crossota stali* Scott, 1874 (Hemiptera: Pentatomidae); during

oocyte resorption in virgin females of this species, vitellins are quickly degraded by proteases and released into the hemolymph as peptides or amino acids (Kotaki 2003).

It is also known that in many species of insects, virgin females do not oviposit mature oocytes, but they are retained in the ovaries or oviducts, where they can be resorbed, and the released material during that resorption can serve as nutritive reserves, used either to mature more oocytes or to increase the life cycle of the females (Engelman 1970). This could suggest that the constituents of the yolk of the oocytes that degenerate during the resorption of the ovarioles of *D. coccus* are employed as nutritive reserves, especially by virgin females, to sustain their metabolic activity or even to increase their longevity, since this could explain why virgin females of *D. coccus* live longer than mated females, because according to Marín and Cisneros (1977) and Pérez-Guerra and Kosztarab (1992), virgin females of *D. coccus* do not oviposit and live longer.

In this study, the fact that no embryos were found in the ovarioles of virgin females of *D. coccus* confirms that mating is necessary for the species to produce fertile eggs, as previously established by Flores-Flores and Tekelenburg (1995). Besides, the finding of completely developed nymphs inside the ovarioles of mated females of *D. coccus* proves that all their embryonic development is intraovarian, and that places it as a clearly ovoviviparous species. However, in *Dactylopius tomentosus* (Lamarch, 1801), egg incubation after oviposition takes from 15 to 20 days (Mathenge *et al.* 2009), which would indicate that, unlike *D. coccus*, *D. tomentosus* is an oviparous species and not ovoviviparous.

Conclusions

Although mating in *D. coccus* is not necessary for its oocytes to initiate and adequately complete the vitellogenesis and choriogenesis, it does promote both processes, as it greatly increases the proportion of mature oocytes to produce in mated females. The lack of mating induced the resorption of the ovarioles and their oocytes, and the vitellogenesis was the affected phase. It is necessary to deep more into the effect of mating in *D. coccus* to elucidate, among other aspects, the hormonal mechanism involved in oocyte maturation, as well as to determine which other factors, besides mating, condition the resorption of ovarioles and oocytes, and the mechanism involved in this process. Besides, it is important to know the final destination of the components of the yolk released by oocytes during their resorption and establish if these components contribute to the longer life of virgin females.

Literature cited

- ADAMS, T. S. 2000a. Effect of diet and mating status on ovarian development in a predaceous stink bug *Perillus bioculatus* (Hemiptera: Pentatomidae). *Annals of the Entomological Society of America* 93: 529-535. [https://doi.org/10.1603/0013-8746\(2000\)093\[0529:EODAMS\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2000)093[0529:EODAMS]2.0.CO;2)
- ADAMS, T. S. 2000b. Effect of diet and mating on oviposition in the twospotted stink bug *Perillus bioculatus* (F.) (Heteroptera: Pentatomidae). *Annals of the Entomological Society of America* 93: 1288-1293. [https://doi.org/10.1603/0013-8746\(2000\)093\[1288:EODAMO\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2000)093[1288:EODAMO]2.0.CO;2)
- ALDAMA-AGUILERA, C.; LLANDERAL-CÁZARES, C. 2003. Grana cochinilla: Comparación de métodos de producción en penca cortada. *Agrociencia* 37: 11-19.
- ASIN, S.; CROCCO DE AYERBE, L. 1992. Influence of mating on ovarian follicle development in *Triatoma infestans* (Klug, 1834).

- Memórias do Instituto Oswaldo Cruz 87 (3): 369-374. <https://doi.org/10.1590/S0074-02761992000300006>
- BELL, W. J.; BOHM, M. K. 1975. Oosorption in insects. *Biological Reviews* 50: 373-396. <https://doi.org/10.1111/j.1469-185X.1975.tb01058.x>
- BRISEÑO-GARZÓN, A.; LLANDERAL, C. 2008. Contenido de ácido carmínico en hembras de grana cochinilla de diferentes edades. pp. 16-20. En: Llanderal, C.; Zetina, D. H.; Viguera, A. L.; Portillo, L. (Eds.). *Grana cochinilla y colorantes naturales*. Colegio de Postgraduados, Texcoco, Edo. de México. México. 124 p.
- DAVEY, K. G. 1997. Hormonal controls on reproduction in female Heteroptera. *Archives of Insect Biochemistry and Physiology* 35: 443-453. [https://doi.org/10.1002/\(SICI\)1520-6327\(1997\)35:4<443::AID-ARCH7>3.0.CO;2-S](https://doi.org/10.1002/(SICI)1520-6327(1997)35:4<443::AID-ARCH7>3.0.CO;2-S)
- DE CLERCQ, P.; DEGHEELE, D. 1997. Effects of mating status on body weight, oviposition, egg load, and predation in the predatory stinkbug *Podisus maculiventris* (Heteroptera: Pentatomidae). *Annals of the Entomological Society of America* 90 (2): 121-127. <https://doi.org/10.1093/aesa/90.2.121>
- DOSSI, F. C. A.; CONSOLI, F. L. 2010. Desenvolvimento ovariano e influencia da cópula na maturação dos ovários de *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae). *Neotropical Entomology* 39 (3): 414-419. <https://doi.org/10.1590/S1519-566X2010000300015>
- ENGELMANN, F. 1970. *The physiology of insect reproduction*. Pergamon Press, Oxford. 307 p.
- FLORES-FLORES, V.; TEKELENBURG, A. 1995. Dacti (*Dactylopius coccus* Costa) dye production. pp. 167-185. In: Barbera, G.; Inglese, P.; Pimienta Barrios, E. (Eds.). *Agroecology, cultivation and uses of cactus pear*. Plant production and protection. Paper 132. FAO, Rome. Italy. 216 p.
- FORTES, P.; SALVADOR, G.; CONSOLI, F. L. 2011. Ovary development and maturation in *Nezara viridula* (L.) (Hemiptera: Pentatomidae). *Neotropical Entomology* 40 (1): 89-96. <https://doi.org/10.1590/S1519-566X2011000100013>
- FRIEDEL, T. 1974. Endocrine control of vitellogenesis in the harlequin bug, *Dindymus versicolor*. *Journal of Insect Physiology* 20: 717-733. [https://doi.org/10.1016/0022-1910\(74\)90192-9](https://doi.org/10.1016/0022-1910(74)90192-9)
- HORTON, D. R.; LEWIS, T. M.; NEVEN, L. G. 2005. Ovarian development and lipid reserves are affected by mating delays in three species of *Anthocoris* (Hemiptera: Anthocoridae). *The Canadian Entomologist* 137: 328-336. <https://doi.org/10.4039/n04-053>
- KOTAKI, T. 2003. Oosorption in the stink bug, *Plautia crossota stali*: induction and vitellogenin dynamics. *Journal of Insect Physiology* 49: 105-113. [https://doi.org/10.1016/S0022-1910\(02\)00254-8](https://doi.org/10.1016/S0022-1910(02)00254-8)
- LAGOWSKA, B.; GOLAN, K. 2009. Scale insects (Hemiptera, Coccoidea) as source of natural dye and other useful substances. *Aphids and other Hemipterous Insects* 15: 151-167.
- LLANDERAL-CÁZARES, C.; NIETO-HERNÁNDEZ, R. 2001. Características biológicas de la grana cochinilla del nopal *Dactylopius coccus* Costa. pp. 23-30. In: Llanderal Cázares C.; Nieto Hernández, R. (Eds.). *Producción de grana cochinilla*. Colegio de Postgraduados. México. 140 p.
- MARÍN, L. R.; CISNEROS, V. F. 1977. Biología y morfología de la cochinilla del carmín, *Dactylopius coccus* Costa (Homopt.: Dactylopiidae). *Revista Peruana de Entomología* 20 (1): 115-120.
- MARTÍNEZ, M. I. 2002. Técnicas básicas de anatomía microscópica y de morfometría para estudiar los insectos. *Boletín de la Sociedad Entomológica Aragonesa* 30: 187-195.
- MASNER, P. 1966. The structure, function and imaginal development of the female inner reproductive organs of *Adelphocoris lineolatus* (Goeze), (Heteroptera, Miridae). *Acta Entomologica Bohemoslovaca* 63: 177-199.
- MATHENGE, C. W.; HOLFORD, P.; HOFFMANN, J. H.; SPOONER-HART, R.; BEATTIE, G. A. C.; ZIMMERMANN, H. G. 2009. The biology of *Dactylopius tomentosus* (Hemiptera: Dactylopiidae). *Bulletin of Entomological Research* 99: 551-559. <https://doi.org/10.1017/S0007485308006597>
- MÉNDEZ-GALLEGOS, S. de J.; PANZAVOLTA, T.; TIBERI, R. 2003. Carmine cochineal *Dactylopius coccus* Costa (Rhynchota: Dactylopiidae): Significance, production and use. *Advances in Horticultural Science* 17 (3): 165-171.
- MUNDALL, E. 1978. Oviposition in *Triatoma protracta*: role of mating and relationship to egg growth. *Journal of Insect Physiology* 24: 321-323. [https://doi.org/10.1016/0022-1910\(78\)90030-6](https://doi.org/10.1016/0022-1910(78)90030-6)
- MUNDALL, E.; ENGELMANN, F. 1977. Endocrine control of vitellogenin synthesis and vitellogenesis in *Triatoma protracta*. *Journal of Insect Physiology* 23: 825-836. [https://doi.org/10.1016/0022-1910\(77\)90007-5](https://doi.org/10.1016/0022-1910(77)90007-5)
- PÉREZ-GUERRA, G.; KOSZTARAB, M. 1992. Biosystematics of the family Dactylopiidae (Homoptera: Coccoidea) with emphasis on the life cycle of *Dactylopius coccus* Costa. Virginia Agricultural Experiment Station. Bulletin No. 92-1. 90 p.
- RAABE, M. 1986. Insect reproduction: regulation of successive steps. *Advances in Insect Physiology* 19: 29-154. [https://doi.org/10.1016/S0065-2806\(08\)60100-9](https://doi.org/10.1016/S0065-2806(08)60100-9)
- RAMÍREZ-CRUZ, A. 2014. Maduración de la ovariola de la grana cochinilla *Dactylopius coccus* (Hemiptera: Coccoidea: Dactylopiidae). *Revista Colombiana de Entomología* 40 (2): 229-233.
- RAMÍREZ-CRUZ, A.; LLANDERAL-CÁZARES, C. 2013. Fecundidad potencial de *Dactylopius coccus* Costa bajo condiciones de invernadero. *Revista Colombiana de Entomología* 39 (1): 170-173.
- RAMÍREZ-CRUZ, A.; LLANDERAL-CÁZARES, C.; RACOTTA, R. 2008. Ovariola structure of cochineal scale insect, *Dactylopius coccus*. *Journal of Insect Science* 8 (20): 1-5. <https://doi.org/10.1673/031.008.2001>
- SISWANTO; MUHAMAD, R.; OMAR, D.; KARMAWATI, E. 2009. The effect of mating on the eggs' fertility and fecundity of *Helopeltis antonii* (Heteroptera: Miridae). *Tropical Life Sciences Research* 20 (1): 89-97.
- SOARES, M. A.; BATISTA, J. D.; ZANUNCIO, J. C.; LINO-NETO, J.; SERRAO, J. E. 2011. Ovary development, egg production and oviposition for mated and virgin females of the predator *Podisus nigrispinus* (Heteroptera: Pentatomidae). *Acta Scientiarum Agronomy* 33 (4): 597-602. <https://doi.org/10.4025/actasciagr.v33i4.6694>

Origin and funding

This project was an independent initiative, without funding support.

Authors contribution

*Arturo Ramírez-Cruz: designed and executed the laboratory work. Both authors analyzed the results and wrote the manuscript. Celina Llanderal-Cázares: carried out *D. coccus* breeding in laboratory.*