

# Arthropod pests and their management, natural enemies and flora visitors associated with castor (*Ricinus communis*), a biofuel plant: a review

Artrópodos plaga y su manejo, enemigos naturales y visitantes florales asociados a la higuerilla (*Ricinus communis*), un cultivo bioenergético: revisión

GUILLERMO LÓPEZ-GUILLÉN<sup>1</sup>; JAIME GÓMEZ-RUIZ<sup>2</sup>; JUAN F. BARRERA<sup>3</sup>

<sup>1</sup> Ds. C. Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias (INIFAP), Campo Experimental Rosario Izapa, C. P. 30780, Tuxtla Chico, Chiapas, Mexico, [lopez.guillermo@inifap.gob.mx](mailto:lopez.guillermo@inifap.gob.mx), <https://orcid.org/0000-0001-7858-9984>. <sup>2</sup> Ds. C. El Colegio de la Frontera Sur, C. P. 30700, Tapachula, Chiapas, Mexico, [jgomez@ecosur.mx](mailto:jgomez@ecosur.mx), <https://orcid.org/0000-0002-9704-9761>. <sup>3</sup> Ph. D. El Colegio de la Frontera Sur, C. P. 30700, Tapachula, Chiapas, Mexico, [jbarrera@ecosur.mx](mailto:jbarrera@ecosur.mx), <https://orcid.org/0000-0002-8488-7782>.

**Abstract:** Interest in bioenergetic crops, such as the castor oil plant *Ricinus communis* (Euphorbiaceae), for production of biodiesel has increased in recent years. In this paper, phytophagous arthropods, their natural enemies and floral visitors associated with this plant in the world are reviewed. Despite its insecticidal properties, arthropods have been reported feeding on *R. communis* plants. The arthropod pests of *R. communis* damage all parts of the plant, including the seeds, where some toxic compounds are even more concentrated. In the scientific databases, we found reports of 193 arthropods associated to *R. communis* in different parts of the world. This information obtained in the scientific databases was concentrated in a database and analyzed according to the coevolutionary hypothesis, which allows us to predict that the greatest wealth and abundance of phytophagous arthropods is found in the center of origin by *R. communis*. According to this review, *Achaea janata*, *Spodoptera litura*, *Edwardsiana flavescens*, *Liriomyza trifolii*, *L. sativae*, *Spilosoma obliqua*, *Cogenethes punctiferalis*, *Oxyrhachis taranda*, and *Helicoverpa armigera* are the most devastating pests in Asia. In Africa, *Agrotis ipsilon*, *S. exigua*, *Nezara viridula*, *Trialeurodes ricini*, and *Tetranychus urticae* were mentioned as the most important. In Central and South-America, *Phyllophaga* sp., *Agriotes* sp., *Erinnys ello*, *N. viridula*, *Corythucha gossypii*, *Falconia antioquiensis*, and *S. marima* are reported as pests of economic importance. The most commonly reported natural enemies of some of these arthropod pests were species of *Bacillus thuringiensis*, *B. cereus*, *B. popilliae*, *Trichogramma achaeae*, *T. chilonis*, *T. minutum*, *T. australicum*, *T. dendrolimi*, *T. pretiosum*, *T. evanescens*, *Microplitis rufiventris*, *M. maculipennis*, *M. ophiussae*, *Telenomus remus*, *T. proditor*, *Stethorus siphonulus* and *S. histrio*. *Apis mellifera* is recorded as the main insect pollinator of *R. communis*. Pest management methods used against the arthropod pests of *R. communis* include biological, ethological, mechanical, cultural, genetic, and chemical control.

**Keywords:** Castor-oil plant, biodiesel, pests, entomophagous organisms, pollinators.

**Resumen:** El interés por los cultivos bioenérgéticos, tales como *Ricinus communis* (Euphorbiaceae) para producir biodiesel ha aumentado en años recientes. En este documento se hace una revisión sobre los artrópodos fitófagos, enemigos naturales y visitantes florales asociados a esta planta en el mundo. A pesar de las propiedades insecticidas de *R. communis*, existen registros sobre artrópodos que se alimentan de ella. Los artrópodos plaga de *R. communis* dañan toda la planta, incluso las semillas, donde se localizan compuestos tóxicos más concentrados. En las bases de datos científicas, se encontró registro de 193 artrópodos asociados a *R. communis* en diferentes partes del mundo. Esta información se concentró en una base de datos y se analizó de acuerdo con la hipótesis coevolutiva, la cual permite predecir que la mayor riqueza y abundancia de artrópodos fitófagos, se encuentra en el centro de origen de *R. communis*. De esta revisión se desprende que entre las plagas más devastadoras en Asia se encuentran *Achaea janata*, *Spodoptera litura*, *Edwardsiana flavescens*, *Liriomyza trifolii*, *L. sativae*, *Spilosoma obliqua*, *Cogenethes punctiferalis*, *Oxyrhachis taranda* y *Helicoverpa armigera*. En África, las plagas más importantes son *Agrotis ipsilon*, *S. exigua*, *Nezara viridula*, *Trialeurodes ricini* y *Tetranychus urticae*. Entre las plagas de importancia económica que se reportan en Centro y Suramérica, están *Phyllophaga* sp., *Agriotes* sp., *Erinnys ello*, *N. viridula*, *Corythucha gossypii*, *Falconia antioquiensis* y *S. marima*. Los enemigos naturales de algunas plagas comúnmente reportados fueron *Bacillus thuringiensis*, *B. cereus*, *B. popilliae*, *Trichogramma achaeae*, *T. chilonis*, *T. minutum*, *T. australicum*, *T. dendrolimi*, *T. pretiosum*, *T. evanescens*, *Microplitis rufiventris*, *M. maculipennis*, *M. ophiussae*, *Telenomus remus*, *T. proditor*, *Stethorus siphonulus* y *S. histrio*. Se registra a *Apis mellifera* como el insecto más polinizador de *R. communis*. Los métodos de manejo de plagas contra artrópodos de *R. communis* incluyen control biológico, etológico, mecánico, cultural, genético y químico.

**Palabras clave:** Higuerilla, biodiesel, plagas, entomófagos, polinizadores.

Corresponding author: Guillermo López-Guillén, Ds. C. Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias (INIFAP), Campo Experimental Rosario Izapa, C. P. 30780, Tuxtla Chico, Chiapas, Mexico, [lopez.guillermo@inifap.gob.mx](mailto:lopez.guillermo@inifap.gob.mx), <https://orcid.org/0000-0001-7858-9984>.

Suggested citation:  
LÓPEZ-GUILLÉN, G.; JAIME GÓMEZ-RUIZ, J.; BARRERA, J. F. 2020. Arthropod pests and their management, natural enemies and floral visitors associated with castor (*Ricinus communis*), a biofuel plant: a review. Revista Colombiana de Entomología 46 (1): e8604. <https://doi.org/10.25100/socolen.v46i1.8604>

Received: 23-Feb-2018

Accepted: 2-Aug-2019

Published: 31-Aug-2020

Revista Colombiana de Entomología

ISSN (Print): 0120-0488

ISSN (On Line): 2665-4385

<https://revistacolombianaentomologia.univalle.edu.co/>

Open access



BY-NC-SA 4.0  
(<https://creativecommons.org/licenses/by-nc-sa/4.0/deed.es>)

Publishers:

Sociedad Colombiana de Entomología

SOCOLEN (Bogotá, D. C., Colombia)

<http://www.socolen.org.co>

Universidad del Valle (Cali, Colombia)

<http://www.univalle.edu.co/>

© 2020 Sociedad Colombiana de Entomología  
- SOCOLEN y Universidad del Valle - Univalle

## Introduction

The castor-oil plant, *Ricinus communis* L., is an oleaginous plant belonging to the Euphorbiaceae family, which comprises 280 genera. This species has been cultivated for more than 6000 years on the Asian continent, and more recently on the African and American continents (Govaerts *et al.* 2000; Salihu *et al.* 2014). *R. communis* is a non-edible plant, mainly used in chemical, pharmaceutical, and automobile industries, where it has numerous applications (Savy 2005; Barnes *et al.* 2009; Severino *et al.* 2010). All parts of this plant contain lectin ricin – one of the most potent lethal natural poisons known – but is particularly concentrated in the seeds and pods (Audi *et al.* 2005).

In recent years, *R. communis* oil has acquired importance as a biofuel, due to the possibility of its use in producing biodiesel (Baldwin and Cossar 2009; César and Batalha 2010). *R. communis* is distributed in tropical and subtropical regions and is also adaptable to temperate zones (Lima *et al.* 2011). The principal producer countries of *R. communis* seeds are India, China, and Mozambique; whereas the countries with the highest consumption of the products of this plant are Holland, Japan, and Italy (Faostat 2015). India, China, and Brazil contribute approximately 95 % of the world production of seeds (Sailaja *et al.* 2008).

*Ricinus communis* seeds are outstanding for their high oil content, between 40 and 60 %, compared with sunflower (*Helianthus annuus* L.) seeds with 38 to 48 %, soybean (*Glycine max* (L.) Merr.) between 18 and 19 %, moringa (*Moringa oleifera* L.) with 14 to 24 %, neem [*Azadirachta indica* (Juss)] between 17 and 39 %, and cotton (*Gossypium hirsutum* L.) with 15 to 19 % (Kittock and Williams 1970; Severino *et al.* 2006; Nass *et al.* 2007; Baldwin and Cossar 2009; Martín *et al.* 2010), a characteristic that makes this plant very attractive as a source of biofuel, particularly biodiesel.

The extensive cultivation of varieties and hybrids of *R. communis* under different management practices has made the plant vulnerable to biotic and abiotic factors. *R. communis* plants may lose leaves, seeds and pods for different reasons: damage by pests, diseases, wind, hail, traffic of machinery, and inappropriate use of herbicides and defoliation (Severino *et al.* 2010). Even though a castor-oil plant can recover from severe defoliation, the damage suffered by the leaves may reduce the production. It is estimated that for 1 m<sup>2</sup> of lost leaf area, seed production diminishes by 37.8 g and oil production by 24.4 g (Lakshmamma *et al.* 2009; Lakshmi *et al.* 2010; Severino *et al.* 2010). Continuous sowing of *R. communis* in the same areas, as well as the lack of intercropping has increased the occurrence of pests and diseases. There are reports that more than 100 species of insects in different parts of the world feed on *R. communis* and can cause serious damage (Barteneva 1986; Kolte 1995). In India, for example, insect pests caused losses in seed production from 35 to 50 % (Kolte 1995). Integrated pest management programs are therefore important to prevent losses that can affect the economy of producer-countries.

The present literature reviewed focuses on the phytophagous arthropods associated with *R. communis* in different parts of the world, as well as, their natural enemies and floral visitors. The information was obtained through extensive search of scientific literature on these subjects published in the Web of Science database, Ebsco database and Google Scholar, using appropriate key words (e.g. ‘insects on *Ricinus communis*’ ‘arthropods on

*Ricinus communis*’, ‘pests of *Ricinus communis* or castor-oil’); the search was conducted until January 2019. Afterwards, the information collected was analyzed from the perspective of the co-evolutionary hypothesis following the approach of literature review analysis of arthropod herbivory on physic nut (*Jatropha curcas* L.) conducted by Lama *et al.* (2015). Specifically, we set out to answer the following questions regarding the arthropods associated with *R. communis*: (1) What is the diversity of arthropod taxa associated with this plant? (2) In what geographic area does the greatest richness of associated arthropod species occur? (3) What are the parts of the plant most preferred by the herbivorous arthropods? and (4) What mouthpart classes of the arthropods associated with *R. communis* can be identified? According to the co-evolutionary hypothesis, it would be expected to find greater richness of native arthropod species in Asia and Africa, the origin area of *R. communis*, in comparison with those areas where this plant has been introduced or cultivated more recently.

## Phytophagous arthropods associated with *R. communis*

*Ricinus communis* has been considered tolerant and/or resistant to pest attack due to the toxic compounds present in different parts of the plant. Some of the most common compounds found in this plant species are ricin, ricinine, N-demethylricinine, flavonoids, gallic acid, gentisic acid, coumaric acid, syringic acid, cinnamic acid, vanillic acid and rutin, and allergen proteins such as Ric c1 and Ric c3 (Usha Rani *et al.* 2006; Gahukar 2010; Vandenborre *et al.* 2011; Usha Rani and Pratyusha 2014). Some of these are toxic compounds that may even have insecticidal or antifeedant properties against insect pests of other crops (Rossi *et al.* 2012; Amoabeng *et al.* 2014; Dinesh *et al.* 2014). Despite the insecticidal properties of *R. communis*, there are reports of arthropods that feed on several parts of this plant. Ricinine, for example, one of its main alkaloids that has shown insecticidal effect on some insect pests of other plants (Bigi *et al.* 2004; Liu and Li 2006; Rossi *et al.* 2012) does not have any detrimental effect on certain specialist phytophagous insects that are common pests of *R. communis*, such as *Achaea janata* (L., 1758) (Lepidoptera: Noctuidae), *Spodoptera litura* (F., 1775) (Lepidoptera: Noctuidae) and others (Prabhakar *et al.* 2003; Usha Rani and Pratyusha 2014). This is due to the presence of enzymes in the midgut of these insects that are able to degrade toxins and thus breakdown the plants' natural defenses (Yasur *et al.* 2009; Usha Rani and Pratyusha 2014).

The arthropod pests of *R. communis* damage all parts of the plant, including the seeds, where some toxic compounds such as lipases, the alkaloid ricinine (including the protein ricin) and glycosides of ricinoleic, isoricinoleic, stearic and dihydroxystearic acids are even more concentrated (Jena and Gupta 2012). The type of pest and damage varies from place to place; some pests of *R. communis* can be present in different regions. Table 1 presents information published in the literature on arthropods that attack *R. communis*.

According to Table 1, 59 % of the arthropod species feed on foliage, 20 % on roots and seedlings, 17 % on flowers, fruits and seeds, and 5 % on stems and branches. The low percentage of arthropods feeding on seeds and roots can be explained in part by the high concentration of ricinine in these parts of the plant (Salihu *et al.* 2014). To feed on seeds and roots, these arthropods have had to develop highly efficient mechanisms of detoxification (Yasur *et al.* 2009).

**Table 1.** Order, family and geographical distribution of the phytophagous arthropod species that attack cultivated *Ricinus communis*.

Order	Family	Species	Geographical distribution	References
<b>Roots and seedlings</b>				
Coleoptera	Curculionidae	<i>Protostropus</i> spp.	Africa	Salihu <i>et al.</i> (2014)
	Elateridae	<i>Agriotes</i> sp.	Costa Rica	Anónimo (1991)
	Scarabeidae	<i>Amphimallon solstitialis</i> (Linnaeus, 1758)	Russia	Arkhangel'Skii and Romanova (1930)
		<i>Holotrichia consanguinea</i> Blanchard, 1850	India	Gahukar (2018)
		<i>Phyllophaga</i> sp.	Colombia and Costa Rica	Anónimo (1991); Londoño-Zuluaga (2008)
		<i>Holochelus aequinoctialis</i> (Herbst, 1790) [= <i>Rhizotrogus aequinoctialis</i> (Herbst, 1790)]	Russia	Arkhangel'Skii and Romanova (1930)
Diptera	Agromyzidae	<i>Liriomyza trifolii</i> (Burgess, 1880)	India	Anjani <i>et al.</i> (2007)
Lepidoptera	Noctuidae	<i>Agrotis ipsilon</i> (Hüfnagel, 1766)	Colombia and Egypt	Mona <i>et al.</i> (2005); Saldarriaga Cardona <i>et al.</i> (2011)
		<i>Helicoverpa zea</i> (Boddie, 1850)	USA	Wene (1933)
		<i>Spodoptera frugiperda</i> (J. E. Smith, 1797)	Colombia	Saldarriaga <i>et al.</i> (2011)
		<i>Spodoptera marima</i> (Schaus, 1904)	Brazil	Ribeiro and Costa (2008)
	Sphingidae	<i>Erinnyis ello</i> (Linnaeus, 1758)	Brazil	Ribeiro and Costa (2008)
Orthoptera	Gryllidae	<i>Brachytrupes</i> spp.	Africa	Salihu <i>et al.</i> (2014)
	Pyrgomorphidae	<i>Chrotononus</i> spp.	Africa	Salihu <i>et al.</i> (2014)
		<i>Zonocerus variegatus</i> (Linnaeus, 1758)	Africa	Salihu <i>et al.</i> (2014)
Isoptera	Termitidae	<i>Odontotermes obesus</i> (Rambur, 1842)	India	Gahukar (2018)
<b>Leaves</b>				
Coleoptera	Curculionidae	<i>Naupactus glaucus</i> Perty, 1832 [= <i>Pantomorus glaucus</i> (Perty, 1830)]	Brazil	Cavalcante <i>et al.</i> (1974)
Diptera	Agromyzidae	<i>Liriomyza sativae</i> Blanchard, 1938	China	Zhang <i>et al.</i> (2006)
		<i>Liriomyza subpussilla</i> Frost, 1943	USA	Wene (1933); Parkman <i>et al.</i> (1989)
		<i>Liriomyza trifolii</i> (Burgess, 1880)	India	Galande <i>et al.</i> (2005)
Hemiptera	Aleyrodidae	<i>Bemisia tabaci</i> (Gennadius, 1889)	Costa Rica and Africa	Anónimo (1991); Salihu <i>et al.</i> (2014)
		<i>Trialeurodes ricini</i> (Misra, 1924) (= <i>Trialeurodes rara</i> Singh, 1931)	India and Egypt	Idriss <i>et al.</i> (1997); Sarma <i>et al.</i> (2005); Abdullah and Martin (2007); Raghavaiah (2011)
	Aphrophoridae	<i>Ptyelus grossus</i> (Fabricius, 1781)	Uganda	Darling (1946)
	Cicadellidae	<i>Amrasca (Amrasca) biguttula</i> (Ishida, 1913) [= <i>Amrasca biguttula biguttula</i> (Ishida, 1912)]	India	Sharma and Singh (2002); Raghavaiah (2011)
		<i>Agallia</i> sp.	Spain	Durán <i>et al.</i> (2010)
		<i>Edwardsiana flavescens</i> (Fabricius, 1794) [= <i>Empoasca flavescens</i> ((Fabricius, 1794))]	India	Jayaraj (1964); Sarma <i>et al.</i> (2005); Lakshmi <i>et al.</i> (2005); Jyothisna <i>et al.</i> (2009)
		<i>Empoasca (Empoasca) solana</i> Delong, 1931 (= <i>Empoasca solana</i> Delong, 1931)	USA	Wene (1933)
		<i>Empoasca</i> sp.	Costa Rica	Anónimo (1991)
		<i>Empoasca</i> sp.	Africa	Salihu <i>et al.</i> (2014)
		<i>Jacobiasca furcostylus</i> (Ramakrishnan y Menon, 1972)	India	Parmar <i>et al.</i> (2006)
	Miridae	<i>Falconia antioquiana</i> Carvalho, 1987	Colombia	Saldarriaga Cardona <i>et al.</i> (2011)
		<i>Polymerus cognatus</i> (Fieber, 1858) (= <i>Poecilocythus cognatus</i> Fieber, 1858)	Russia	Arkhangel'Skii and Romanova (1930)
	Pentatomidae	<i>Acrosternum pallidoconspersum</i> (Stål, 1858)	Egypt	Jannone (1952)
		<i>Nezara viridula</i> (Linnaeus, 1758)	Costa Rica and Egypt	Jannone (1952); Anónimo (1991)
	Pseudococcidae	<i>Paracoccus marginatus</i> Williams and Granara de Willink, 1992	Cuba	Martínez <i>et al.</i> (2005)
	Tingidae	<i>Corythucha gossypii</i> (Fabricius, 1794)	USA, Colombia, Mexico, and Cuba	Miller and Nagamine (2005); Londoño-Zuluaga (2008); Saldarriaga Cardona <i>et al.</i> (2011), López-Guillén <i>et al.</i> (2012)
Lepidoptera	Arctiidae	<i>Amsacta moorei</i> Butler, 1876	India	Sarma <i>et al.</i> (2005)
		<i>Amsacta albistriga</i> Walker, 1864	India	Sarma <i>et al.</i> (2005)
		<i>Pericallia ricini</i> (Fabricius, 1775)	India	Mathur <i>et al.</i> (1994); Neelanarayanan and Indira (2010)
		<i>Spilosoma obliqua</i> Walker, 1855	India and Pakistan	Singh and Grewal (1982); Khattak <i>et al.</i> (1991); Sarma <i>et al.</i> (2005)

continued...

... cont.

	Dalceridae	<i>Anacraga citrinopsis</i> Dyar, 1927	Brazil	Lourenço <i>et al.</i> (1989)
	Limacodidae	<i>Parasa lepida</i> Cramer, 1799	India	Raghavaiah (2011)
	Lymantriidae	<i>Dasychira</i> sp.	Africa	Salihu <i>et al.</i> (2014)
		<i>Euproctis fraterna</i> Moore, 1883	India	Paul <i>et al.</i> (2000); Suganthy (2010)
	Noctuidae	<i>Achaea janata</i> (Linnaeus, 1758)	India, USA, and China	Hua (1984); Delaya <i>et al.</i> (1985); Basappa and Lingappa (2001); Mau and Kessing (2007)
		<i>Helicoverpa armigera</i> (Hübner, 1803-1808)	India and USA	Wene (1933); Ribeiro and Costa (2008)
		<i>Spodoptera cosmioides</i> (Walker, 1858)	Brazil	Bavaresco <i>et al.</i> (2003)
		<i>Spodoptera exigua</i> (Hübner, 1808)	Egypt	Ribeiro and Costa (2008)
		<i>Spodoptera litura</i> (Fabricius, 1775)	India and Pakistan	Lohar <i>et al.</i> (1997); Usha Rani and Rajasekharreddy (2009)
		<i>Spodoptera ornithogalli</i> (Guenée, 1852) [= <i>Spodoptera marima</i> (Schaus, 1904)]	Brazil	Ribeiro and Costa (2008)
		<i>Spodoptera</i> sp.	Costa Rica	Anónimo (1991)
	Nymphalidae	<i>Ariadne merione</i> Cramer, 1779 [= <i>Ergolis merione</i> Cramer, 1779]	India	Ghosh (1914); Sarma <i>et al.</i> (2005)
	Saturniidae	<i>Samia ricini</i> (Drury, 1773)	Egypt, India, and Brazil	El-Shaarawy <i>et al.</i> (1975); Negreiros <i>et al.</i> (1998)
		<i>Rothschildia jacobaeae</i> Walker, 1855	Brazil	Ribeiro and Costa (2008)
Orthoptera	Acrididae	<i>Chrotogonus (Chrotogonus) trachypterus robertsi</i> Kirby & W. F., 1914 (= <i>Chrotogonus robertsi</i> Kirby & W. F., 1914)	India	Sarma <i>et al.</i> (2005)
Thysanoptera	Thripidae	<i>Retithrips syriacus</i> (Mayet, 1890)	India	Sarma <i>et al.</i> (2005)
		<i>Scirtothrips dorsalis</i> Hood, 1919	India	Patel <i>et al.</i> (2009)
		<i>Zaniothrips ricini</i> Bhatti, 1967	India	Daniel <i>et al.</i> (1983)
Acarina	Tetranychidae	<i>Eutetranychus orientalis</i> (Klein, 1936)	India	Ahuja (1994)
		<i>Eutetranychus</i> sp.	India	Raghavaiah (2011)
		<i>Tetranychus piercei</i> McGregor, 1950	China	Lui and Lui (1986)
		<i>Tetranychus urticae</i> Koch, 1836 [= <i>Tetranychus telarius</i> (Linnaeus, 1758)]	Morocco and India	Cangardel (1954); Rajasekhar <i>et al.</i> (1999); Raghavaiah (2011)
	Tarsonemidae	<i>Polyphagotarsonemus latus</i> (Banks, 1904)	Belgium	Heungens and Degheele (1986)
<b>Stems and branches</b>				
Coleoptera	Buprestidae	<i>Sphenoptera</i> sp.	Africa	Salihu <i>et al.</i> (2014)
	Tenebrionidae	<i>Blapstinus</i> sp.	USA	De Ong (1918)
Hemiptera	Membracidae	<i>Oxyrhachis taranda</i> (Fabricius, 1798)	India	Ali <i>et al.</i> (2006)
Lepidoptera	Cossidae	<i>Strigocossus capensis</i> (Walker, 1856) [= <i>Xyleutes capensis</i> (Walker, 1856)]	Africa	Salihu <i>et al.</i> (2014)
<b>Flowers, fruits and seeds</b>				
Coleoptera	Anobiidae	<i>Lasioderma serricorne</i> (Fabricius, 1792)	India and Africa	Hussain and Khan (1966); Salihu <i>et al.</i> (2014)
	Tenebrionidae	<i>Tribolium castaneum</i> (Herbst, 1797)	Africa	Salihu <i>et al.</i> (2014)
Hemiptera	Cicadellidae	<i>Empoasca</i> sp.	Costa Rica	Anónimo (1991)
	Miridae	<i>Eurystylus</i> sp.	Africa	Salihu <i>et al.</i> (2014)
		<i>Helopeltis</i> sp.	Africa	Salihu <i>et al.</i> (2014)
	Pentatomidae	<i>Nezara viridula</i> (Linnaeus, 1758)	Costa Rica, and USA	Anónimo (1991); Golden and Follett (2006)
	Scutelleridae	<i>Calidea</i> sp.	Africa	Salihu <i>et al.</i> (2014)
Lepidoptera	Crambidae	<i>Conogethes punctiferalis</i> (Guenée, 1854) [= <i>Dichocrocis punctiferalis</i> (Guenée, 1854)]	India and Australia	Anonymous (1913); Sharma <i>et al.</i> (1995); Jyothsna <i>et al.</i> (2009); Patel and Patel (2009); Hedge <i>et al.</i> (2009)
	Noctuidae	<i>Achaea janata</i> (Linnaeus, 1758)	India, USA, and China	Hua (1984); Delaya <i>et al.</i> (1985); Basappa and Lingappa (2001); Mau and Kessing (2007)
		<i>Heliothis</i> sp.	Costa Rica	Anónimo (1991)
		<i>Helicoverpa armigera</i> (Hübner, 1803-1808)	India, and USA	Wene (1933); Geetha <i>et al.</i> (2003); Satyanarayana and Sing (2003)
		<i>Spodoptera</i> sp.	Costa Rica	Anónimo (1991)
	Pyralidae	<i>Cadra cautella</i> (Walker, 1863) [= <i>Epehestia cautella</i> (Walker, 1863)]	Africa	Salihu <i>et al.</i> (2014)
	Tortricidae	<i>Thaumatotibia leucotreta</i> (Meyrick, 1913) [= <i>Cryptophlebia leucotreta</i> Meyrick, 1913]	Africa	Salihu <i>et al.</i> (2014)

A total of 76 species of phytophagous arthropods associated to cultivated plants of *R. communis* is found worldwide (Table 1). Before the present literature review, the report was of 60 species (Raouf *et al.* 2003). The arthropods reported in Table 1 belong to eight orders and 38 families; 40 % of these species belong to Lepidoptera, 27 % to Hemiptera, 14 % to Coleoptera and 19 % to other orders. The species that belong to Lepidoptera, Hemiptera and Coleoptera represent 81 % of the total. These phytophagous arthropods are distributed geographically in Asia (39 %), America (34 %), Africa (25 %) and Europe (2 %). As it was supposed, it was not uncommon to find that the greatest richness of arthropods associated to *R. communis* occurred in Asia and Africa, continents considered as the center of origin of this plant (Govaerts *et al.* 2000). 63 % of the species had mandibulate mouthparts (Lepidoptera, Coleoptera, Orthoptera, Isoptera and Diptera) and 37 % were piercing-and-sucking mouthpart classes (Hemiptera, Thysanoptera and Acarina).

Of the pests listed in Table 1, the castor semilooper *A. janata*, the tobacco caterpillar *S. litura*, the green leafhopper *Edwardsiana flavescens* (F., 1794) [= *Empoasca flavescens* (F., 1794)] (Hemiptera: Cicadellidae), the serpentine leafminer *Liriomyza trifolii* Burgess, 1880, the vegetable leafminer *L. sativae* Blanchard, 1938 (Diptera: Agromyzidae), the Bihar hairy caterpillar *Spilosoma obliqua* Walker, 1855 (Lepidoptera: Arctiidae), the shoot and capsule borer *Conogethes punctiferalis* (Guenée, 1854) [= *Dichocrocis punctiferalis* (Guenée, 1854)] (Lepidoptera: Crambidae), the cowbug *Oxyrhachis taranda* (F., 1798) (Hemiptera: Membracidae), and the cotton bullworm *Helicoverpa armigera* (Hübner, 1803-1808) (Lepidoptera: Noctuidae), among others, are the most devastating pests in Asia. In Africa, the black cutworm *Agrotis ipsilon* (Hüfnagel, 1776), the armyworm *S. exigua* (Hübner, 1808) (Lepidoptera: Noctuidae), the stink bug *Nezara viridula* (L., 1758) (Hemiptera: Pentatomidae), the castor bean whitefly *Trialeurodes ricini* (Misra, 1924) (= *Trialeurodes rara* Singh, 1931) (Hemiptera: Aleyrodidae), the red spider mite *Tetranychus urticae* Koch, 1836 [= *Tetranychus telarius* (Linnaeus, 1758)] (Acarina: Tetranychidae), among others, are mentioned as the most important. In Central and South-America, the white grub *Phyllophaga* sp. (Coleoptera: Scarabaeidae), *Agriotes* sp., *Erinnys ello* (F., 1794) (Lepidoptera: Sphingidae), *N. viridula*, the cotton lace bug *Corythucha gossypii* (Fabricius, 1794) (Hemiptera: Tingidae), the sucking bug *Falconia antioquiana* Carvalho, 1987 (Hemiptera: Miridae), *S. marima* (Schaus, 1904) (Lepidoptera: Noctuidae) and others, are reported as pests of economic importance for *R. communis* (Varón *et al.* 2010; Saldarriaga Cardona *et al.* 2011; López-Guillén *et al.* 2012).

The principal pests in Brazil are *N. viridula*, the leafhopper *Empoasca* spp., some defoliator larvae including *S. frugiperda* Smith (Lepidoptera: Noctuidae), *A. janata*, and *A. ipsilon*, and mites such as *T. urticae*, and *T. ludeni* Zacher, 1913 (Acarina: Tetranychidae) (Soares *et al.* 2001; Ribeiro and Costa 2008). In Colombia, *C. gossypii* is mentioned as the pest of greatest economic importance in *R. communis* crops (Varón *et al.* 2010). In Mexico, *C. gossypii*, *N. viridula*, and *Tetranychus* spp. are reported as the main potential pests of *R. communis* (López-Guillén *et al.* 2012) (Table 1).

## Phytophagous arthropods found on noncultivated *R. communis*

There are reports of phytophagous insects mostly found in noncultivated *R. communis* plants, isolated plants, as well as, in more or less clustered plants or plants growing in urban and suburban areas, and in disturbed landscapes of Egypt, India, Spain, Uganda, USA and other countries (Oshaibah *et al.* 1986; Singh *et al.* 1991; Jacob *et al.* 2000; Pons *et al.* 2002; Ylla *et al.* 2008; Boland 2016; Egonyu *et al.* 2017).

Table 2 presents a total of 20 species of phytophagous arthropods associated with non-cultivated plants of *R. communis* in the world. These species belong to five orders and 16 families. 60 % of the arthropod species belong to Lepidoptera (30 %) and Hemiptera (30 %), while 40% belong to Coleoptera (20 %) and other orders (20 %). The species that belong to Lepidoptera, Hemiptera and Coleoptera represent 80 % of the total. 59 % of the arthropod species registered in Table 2 are distributed geographically in Asia (32 %) and Africa (27 %), while 41 % are registered in America (32 %) and Europe (9 %). 60 % of the species of arthropods are mandibulate mouthpart (Lepidoptera, Coleoptera, and Orthoptera) and 40 % are piercing-and-sucking mouthpart (Hemiptera and Acarina).

Such insects were observed feeding on leaves of *R. communis* plants, and even though some species have been reported as pests of *R. communis* in other countries, most of them cause no considerable damage. However, they have the potential of becoming pests of *R. communis* if it is cultivated as a monoculture or, *R. communis* could be a plant host for important pests as the invasive ambrosia beetle *Euwallacea* sp. (Coleoptera: Curculionidae) (Boland 2016; Egonyu *et al.* 2017). Among these potential pests are insect and mite species of various families of Lepidoptera, Hemiptera, Orthoptera, and others (Table 2).

## Pollinator insects and floral visitors in *R. communis*

*Ricinus communis* is a monoecious cross-pollinating plant, cultivated as a hybrid in India, Brazil, China, and other countries because they produce better yields than pure lines or varieties (Moll *et al.* 1962; Birchler *et al.* 2003; Reif *et al.* 2007). Several studies demonstrate that certain species of pollinator insects may improve seed production of *R. communis*. For example, it is mentioned that *Apis mellifera* (L., 1758) (Hymenoptera: Apidae) contributes to increasing *R. communis* crop productivity by incrementing fruit numbers as well as oil content in seeds (Freitas and Cruz 2010).

Among the pollinator insects of *R. communis*, *A. mellifera* is recorded as the main pollinating insect. It is also mentioned that this insect feeds on the nectar produced by the plant's extrafloral nectar glands (Rizzato *et al.* 2012; Waters *et al.* 2014). *A. mellifera* is the principal pollinating insect of *R. communis*, and laboratory work has demonstrated that the pollen of this plant reduces bee survival (Junior *et al.* 2011). According to these studies, expansion of *R. communis* as a crop in the semiarid region of Brazil for biodiesel production represents a risk for the native and domestic bees used for honey production.

As shown in Table 3, a total of 36 species of pollinator insects and floral visitors of non-cultivated plants of *R. communis* is found in the world. These species belong to four orders and 16 families. 25 % of the species belong

**Table 2.** Order, family and geographical distribution of the phytophagous arthropod species found on non-cultivated *Ricinus communis*.

Order	Family	Species	Geographical distribution	References
<b>Leaves</b>				
Coleoptera	Bostrichidae	<i>Prostephanus truncatus</i> (Horn, 1878)	Mexico	Bourne-Murrieta <i>et al.</i> (2014)
	Chrysomelidae	<i>Diabrotica graminea</i> Baly, 1886	Puerto Rico	Wolcott (1917)
	Scarabaeidae	<i>Lepadoretus sinicus</i> Burmeister, 1855 ( <i>= Adoretus sinicus</i> Burmeister, 1855)	USA	McQuate y Jameson (2011)
	Scolytidae	<i>Euwallacea</i> sp.	Uganda and USA	Boland (2016); Egonyu <i>et al.</i> (2017)
Hemiptera	Aleyrodidae	<i>Aleurodicus dispersus</i> Russell, 1965	Cape Verde	Monteiro <i>et al.</i> (2005)
	Cicadellidae	<i>Amrasca (Amrasca) biguttula</i> (Ishida, 1913) [ <i>= Amrasca devastans</i> (Distant, 1918)]	India	Jacob <i>et al.</i> (2000)
		<i>Empoasca (Empoasca) kerri</i> Singh-Pruthi, 1940 ( <i>= Empoasca kerri</i> Pruthi, 1940)	India	Singh <i>et al.</i> (1991); Jacob <i>et al.</i> (2000)
		<i>Empoasca (Empoasca) motti</i> Singh-Pruthi, 1940 ( <i>= Empoasca motti</i> Singh-Pruthi, 1940)	India	Jacob <i>et al.</i> (2000)
	Flatidae	<i>Metcalfa pruinosa</i> (Say, 1830)	Spain	Pons <i>et al.</i> (2002)
	Miridae	<i>Apolygus lucorum</i> (Meyer-Dür, 1843)	China	Lu <i>et al.</i> (2010)
Lepidoptera	Arctiidae	<i>Amsacta moorei</i> Butler, 1876	India	Singh <i>et al.</i> (1989)
	Cosmopterigidae	<i>Pyroderces rileyi</i> (Walsingham, 1882) ( <i>= Sathrobrota rileyi</i> Walsingham, 1882)	Egypt	Oshaibah <i>et al.</i> (1986)
	Lymantriidae	<i>Euproctis lunata</i> Walker, 1855	Bangladesh	Islam <i>et al.</i> (1988)
	Noctuidae	<i>Agrotis ipsilon</i> (Hüfnagel, 1766)	Egypt	Younis (1992)
	Pyralidae	<i>Phycita diaphana</i> (Staudinger, 1870)	Spain	Huertas Dionisio (2002); Ylla <i>et al.</i> (2008)
	Tortricidae	<i>Thaumatotibia leucotreta</i> (Meyrick, 1913) ( <i>= Cryptophlebia leucotreta</i> Meyrick, 1913)	South Africa	Kirkman and Moore (2007)
Orthoptera	Acrididae	<i>Melanoplus differentialis</i> (Thomas, 1865)	USA	Spain (1940)
Acarina	Tetranychidae	<i>Eutetranychus banksi</i> (McGregor, 1914)	USA	McGregor (1914)
		<i>Eutetranychus orientalis</i> (Klein, 1936)	Palestine and Egypt	Klein (1936)
		<i>Tetranychus gloveri</i> Banks, 1900 ( <i>= Tetranychus quinquenychus</i> McGregor, 1914)	USA	McGregor (1914)

to Lepidoptera (19 %) and Hemiptera (6 %), while 75 % belong to Hymenoptera (67 %) and Diptera (8 %). 55 % of the arthropod species registered in Table 3 are distributed geographically in Asia (33 %) and Africa (22 %), while 45 % are registered in America; no records were found for Europe.

In Mexico, Cameroon, USA, India, and Brazil, entomophagous Hymenoptera, as well as several species of Lepidoptera, Diptera, and Hemiptera have been reported to feed on nectaries and flowers of *R. communis*; however, only *A. mellifera* has been reported as a pollinator. Therefore, it is necessary to carry out studies on pollination and floral ecology in order to determine if there are other insect pollinators of *R. communis* that should be protected or may be used to increase crop yield (Table 3).

Some pests can affect pollinators through herbivory. In the case of *R. communis*, Wäckers *et al.* (2001) showed that plants damaged by larvae of *Spodoptera littoralis* (Boisd., 1833) (Lepidoptera: Noctuidae) increased the total amount of nectar produced by extrafloral nectaries compared to undamaged plants. De Sibio and Rossi (2016) found a similar result for the herbivory of *S. frugiperda* on *R. communis*. The secretion of carbohydrates through extrafloral nectaries is considered an indirect strategy of plant defense because it serves to attract parasitoids and predators (Heil 2008). Unlike floral nectaries, extrafloral nectaries do not participate in pollination,

however, in plants pollinated by insects, extrafloral nectaries can negatively affect the effectiveness of pollination by distracting pollinators away from floral nectaries or when the ants that are attracted by the nectar attack the floral visitors (Wäckers *et al.* 2001; Turlings and Wäckers 2004).

#### Natural enemies of the pests of *R. communis*

Among the natural enemies of the key pests of cultivated *R. communis*, there are parasitoids, predators, and entomopathogens such as fungi, bacteria, nematodes, and viruses, which are used as biological control agents or have been found parasitizing, depredating, or naturally infecting some pests of the crop. An extensive list of natural enemies of phytophagous arthropods of *R. communis* grouped by taxa with information of their host or prey and geographical distribution is shown in Table 4; as it can appreciate in this table, the most commonly reported natural enemies in countries like India, Brazil, China, and USA, are *Bacillus* spp., *Trichogramma* spp., *Microplitis* spp., *Telenomus* spp., *Stethorus* spp., and other species attacking pests such as *A. janata*, *S. litura*, *Anacraga citrinopsis* Dyar, 1927, *S. obliqua*, *Phyllophaga* sp., *Eutetranychus banksi* (McGregor, 1914), *Tetranychus piercei* McGregor, 1950, *Zaniothrips ricini* Bhatti, 1967, and other species. Table 4 shows a total of 61

natural enemies of phytophagous insects of *R. communis*. Three species are bacteria belonging to the same genus; four species are nematodes of different genera; two species are fungi of different genera; two reports are viruses; 36 species are parasitoids of eight families of Hymenoptera and one family of Diptera; and 14 species are predators of six different families and order 74 % of the species is distributed geographically in Asia, 24 % in America, 2 % in Africa and 0 % in Europe.

An example of natural enemies of pest of *R. communis* is presented by Basappa (2009). According to this author, parasitoids, insect predators, spiders, insectivorous birds and some microbial organisms are important natural enemies of the pest complex of *R. communis* ecosystem in India. In the case of *A. janata*, *Trichogramma chilonis* Ishii, 1941, *Trichogramma achaeae* Nagaraja and Nagarkatti, 1970, *Telenomus* sp. and *Trissolcus* sp. were recorded from eggs; *Microplitis maculipennis* (Szepligeti, 1900), *Euplectrus*

**Table 3.** Order, family and geographical distribution of the pollinators and floral visitors, reported on *Ricinus communis* plants.

Order	Family	Species	Geographical distribution	References
Diptera	Muscidae	<i>Musca domestica</i> Linnaeus, 1758	Cameroon	Douka and Tchuenguem (2014)
	Richardiidae	<i>Sepsisoma</i> sp.	Brazil	Souza-Silva <i>et al.</i> (2001)
	Syrphidae	<i>Ischiodon scutellaris</i> (Fabricius, 1805)	India	Navatha and Sreedevi (2012)
Hemiptera	Coreidae	<i>Anoplocnemis curvipes</i> (Fabricius, 1871)	Cameroon	Douka and Tchuenguem (2014)
	Pentatomidae	<i>Nezara viridula</i> Linnaeus, 1758	India	Navatha and Sreedevi (2012)
Lepidoptera	Nymphalidae	<i>Acraea acerata</i> (Hewitson, 1874)	Cameroon	Douka and Tchuenguem (2014)
		<i>Acraea terpsicore</i> (Linnaeus, 1758)	India	Navatha and Sreedevi (2012)
		<i>Hypolimnas misippus</i> (Linnaeus, 1764)	India	Navatha and Sreedevi (2012)
	Pieridae	<i>Catopsilia florella</i> (Fabricius, 1775)	Cameroon	Douka and Tchuenguem (2014)
		<i>Eurema blanda</i> (Boisduval, 1836)	India	Navatha and Sreedevi (2012)
		<i>Eurema</i> sp.	Cameroon	Douka and Tchuenguem (2014)
		<i>Pieris brassicae</i> (Linnaeus, 1758)	India	Navatha and Sreedevi (2012)
Hymenoptera	Apidae	<i>Apis mellifera</i> Linnaeus, 1758	Brazil	Freitas <i>et al.</i> (2009); Freitas and Cruz (2010)
		<i>Apis florea</i> Fabricius, 1973	India	Navatha and Sreedevi (2012)
		<i>Ceratina</i> sp.	India	Navatha and Sreedevi (2012)
		<i>Scaptotrigona</i> sp.	Brazil	Freitas <i>et al.</i> (2009)
		<i>Trigona</i> sp.	India	Navatha and Sreedevi (2012)
		<i>Xylocopa fenerstrata</i> (Fabricius, 1798)	India	Navatha and Sreedevi (2012)
	Braconidae	<i>Bracon</i> spp.	Mexico	Álvarez and Reyes (1987)
		<i>Ephialtux</i> sp.	Mexico	Álvarez and Reyes (1987)
	Chalcididae	<i>Conura igneoides</i> Kirby, 1883 [= <i>Spilochalcis igneoides</i> (Kirby, 1883)]	Mexico	Álvarez and Reyes (1987)
		<i>Conura maria</i> Riley, 1870 [= <i>Spilochalcis mariae</i> (Riley, 1872)]	Mexico	Álvarez and Reyes (1987)
Eurytomidae		<i>Neorileya</i> sp.	Mexico	Álvarez and Reyes (1987)
	Formicidae	<i>Camponotus compressus</i> (Fabricius, 1787)	India	Navatha and Sreedevi (2012)
		<i>Linepithema humile</i> (Mayr, 1868)	USA	Line <i>et al.</i> (2013)
		<i>Polyrachis</i> sp.	Cameroon	Douka and Tchuenguem (2014)
Halictidae		<i>Halictus</i> sp.	India	Navatha and Sreedevi (2012)
	Sphecidae	<i>Liris</i> sp.	Mexico	Álvarez and Reyes (1987)
Torymidae		<i>Sceliphron assimile</i> (Dahlbom, 1843)	Mexico	Álvarez and Reyes (1987)
		<i>Tachysphex</i> sp.	Mexico	Álvarez and Reyes (1987)
		<i>Tachytes</i> sp.	Mexico	Álvarez and Reyes (1987)
		<i>Trypoxylon</i> sp.	Mexico	Álvarez and Reyes (1987)
		<i>Torymus capillaceus</i> (Huber, 1927)	Mexico	Álvarez and Reyes (1987)
Vespidae		<i>Synagris cornuta</i> (Linnaeus, 1758)	Cameroon	Douka and Tchuenguem (2014)
		<i>Delta</i> sp.	Cameroon	Douka and Tchuenguem (2014)
		<i>Polistes</i> sp.	Mexico	Álvarez and Reyes (1987)

**Table 4.** Natural enemies of phytophagous arthropods of *Ricinus communis*.

Species	Host and/or prey	Geographical distribution	References
<b>Entomopathogens</b>			
<b>Bacteria</b>			
<i>Bacillus thuringiensis</i> var. <i>kurstaki</i> (Berliner, 1915)	Larvae of <i>Achaea janata</i>	India	Vimala Devi and Sudhakar (2006)
<i>Bacillus cereus</i> (Manson, Pollock & Tridgell, 1954)	Larvae of <i>Achaea janata</i>	India	Kattegoudar et al. (1994)
<i>Bacillus popilliae</i> Dutky, 1940	Larvae of <i>Phyllophaga</i> sp.	Colombia	Saldarriaga Cardona et al. (2011)
<b>Nematodes</b>			
<i>Hexameris dactylocercus</i> Poinar and Linares, 1985	Larvae of <i>Amsacta albistriga</i>	India	Prabhakar et al. (2010)
<i>Steinernema carpocapsae</i> (Weiser, 1955)	Larvae of <i>Spodoptera litura</i>	India	Raveendranath et al. (2008)
<i>Heterorhabditis indica</i> Poinar, Karunaka y David, 1992	Larvae of <i>Spodoptera litura</i>	India	Raveendranath et al. (2008)
<i>Mermis</i> sp.	Larvae of <i>Achaea janata</i>	India	Sujatha et al. (2011)
<b>Fungi</b>			
<i>Metarhizium rileyi</i> (Farl.) Kepler, S.A.Rehner & Humber, 2014 [= <i>Nomuraea rileyi</i> (Farlow) Samson, 1974]	Larvae of <i>Spodoptera litura</i>	India and USA	Mau and Kessing (2007)
<i>Beauveria bassiana</i> (Balsamo) Vuillemin, 1912	Larvae of <i>Achaea janata</i> and <i>Cogenethes punctiferalis</i>	India	Duraimurugan et al. (2015)
<b>Virus</b>			
Nucleopolyhedrovirus	Larvae of <i>Spodoptera litura</i>	India	Basappa (2009)
Granulovirus	Larvae of <i>Achaea janata</i> and <i>Spodoptera litura</i>	India	Naveen Kumar et al. (2013)
<b>Parasitoids</b>			
<b>INSECTA</b>			
<b>Hymenoptera</b>			
<b>Aphelinidae</b>			
<i>Encarsia formosa</i> Gahan, 1924	Nymphs of <i>Trialeurodes ricini</i>	China	Wang et al. (2016)
<b>Braconidae</b>			
<i>Habrobracon hebetor</i> (Say, 1836)	Larvae of <i>Cogenethes punctiferalis</i>	India	Basappa (2003)
<i>Apanteles hyposidrae</i> Wilkinson, 1928	Larvae of <i>Achaea janata</i>	India	Basappa (2009)
<i>Apanteles ricini</i> Bhatnagar, 1948	Larvae of <i>Cogenethes punctiferalis</i>	India	Basappa (2003)
<i>Cotesia flavipes</i> (Cameron, 1891) [ <i>Apanteles flavipes</i> (Cameron, 1891)]	Larvae of <i>Spilosoma obliqua</i> and <i>Spodoptera litura</i>	India	Yadav et al. (2010); Basappa (2009)
<i>Glyptapanteles dalosoma</i> de Santis, 1987	Larvae of <i>Anacraga citrinopsis</i>	Brazil	Lourenço et al. (1989)
<i>Microplitis (= Microgaster) rufiventris</i> Kokujev, 1914	Larvae of <i>Spodoptera littoralis</i>	Egypt	Shalaby et al. (1988)
<i>Microplitis maculipennis</i> (Szepligeti, 1900) (= <i>Microplitis ophiusa</i> Aiyar, 1921)	Larvae of <i>Achaea janata</i>	India	Suganthy (2010); Naik et al. (2010)
<b>Chalcididae</b>			
<i>Brachymeria euploiae</i> (Westwood, 1837)	Pupae of <i>Cogenethes punctiferalis</i>	India	Sujatha et al. (2011)
<b>Eulophidae</b>			
<i>Ceranisus menes</i> (Walker, 1839)	2º instar nymph of <i>Zaniothrips ricini</i>	India	Daniel et al. (1983)
<i>Euplectrus maternus</i> Bhatnagar, 1952	Larvae of <i>Achaea janata</i>	India	Basappa (2009)
<i>Tetrastichus howardi</i> (Olliff, 1893) (= <i>Tetrastichus ayyari</i> Rohwer, 1921)	Pupae of <i>Spodoptera litura</i> and <i>Achaea janata</i>	India	Basappa (2009)
<i>Trichospilus pupivorus</i> Ferrière, 1930	Pupae of <i>Spodoptera litura</i> and <i>Achaea janata</i>	India	Basappa (2009)
<b>Trichogrammatidae</b>			
<i>Trichogramma achaeae</i> Nagaraja and Nagarkatti, 1970	Eggs of <i>Achaea janata</i>	India	Basappa (2009)
<i>Trichogramma chilonis</i> Ishii, 1941	Eggs of <i>Achaea janata</i> and <i>Spodoptera litura</i>	India	Singh et al. (2008); Suganthy (2010); Naik et al. (2010)
<i>Trichogramma minutum</i> Riley, 1879	Eggs of <i>Achaea janata</i>	USA	Mau and Kessing (2007)
<i>Trichogramma australicum</i> Girault, 1912	Eggs of <i>Achaea janata</i>	China	Hua (1984)
<i>Trichogramma dendrolimi</i> Matsumura, 1926	Eggs of <i>Achaea janata</i>	China	Hua (1984)
<i>Trichogramma pretiosum</i> Riley, 1879	Eggs of <i>S. cosmoides</i>	Brazil	Cabezas et al. (2013)
<i>Trichogramma evanescens</i> Westwood, 1833	Eggs of <i>Achaea janata</i>	India	Basappa (2009)

continued...

cont...

**Scelionidae**

<i>Telenomus remus</i> Nixon, 1937	Eggs of <i>Spodoptera litura</i> , <i>Spodoptera cosmooides</i> and <i>Spodoptera frugiperda</i>	India	Satyanarayana <i>et al.</i> (2005); Pomari <i>et al.</i> (2013)
<i>Telenomus proditor</i> Nixon, 1937	Eggs of Lepidoptera	USA	Mau and Kessing (2007)
<i>Telenomus</i> sp.	Eggs of <i>Achaea janata</i>	India	Basappa (2009)
<i>Trissolcus</i> sp.	Eggs of <i>Achaea janata</i>	India	Basappa (2009)

**Vespidae**

<i>Polistes</i> sp.	Larvae of <i>Phyllophaga</i> sp., <i>Agrotis</i> sp. and <i>Spodoptera</i> spp.	Costa Rica	Anónimo (1991)
---------------------	---	------------	----------------

**Ichneumonidae**

<i>Campoletis chlorideae</i> Uchida, 1957	Larvae of <i>Spodoptera litura</i>	India	Satyanarayana <i>et al.</i> (2005)
<i>Charops obtusus</i> Morley, 1913	Larvae of <i>Spilosoma obliqua</i> and <i>Achaea janata</i>	India	Basappa (2009)
<i>Hyposoter exiguae</i> (Viereck, 1912)	Larvae of <i>Achaea janata</i>	USA	Mau and Kessing (2007)
<i>Diadegma ricini</i> Row & Kurian, 1950	Larvae of <i>Cogenethes punctiferalis</i>	India	Basappa (2003)
<i>Theronia</i> sp.	Larvae of <i>Cogenethes punctiferalis</i>	India	Basappa (2003)
<i>Isdromas monterai</i> (Costa Lima, 1948)	Larvae of <i>Anacraga citrinopsis</i>	Brazil	Lourenço <i>et al.</i> (1989)

**Tachinidae**

<i>Palexorista parachrysops</i> Bezzi, 1925	Larvae of <i>Cogenethes punctiferalis</i>	India	Kalra (1984)
<i>Eucelatoria armigera</i> (Coquillett, 1889)	Larvae and pupae of <i>Achaea janata</i>	USA	Mau and Kessing (2007)
<i>Chaetogaedia monticola</i> (Bigot, 1887)	Larvae and pupae of <i>Achaea janata</i>	USA	Mau and Kessing (2007)

**Predators****Coleoptera****Carabidae**

<i>Calosoma</i> sp.	Larvae of <i>Phyllophaga</i> sp., <i>Agrotis</i> sp. and <i>Spodoptera</i> spp.	Costa Rica	Anónimo (1991)
---------------------	---	------------	----------------

**Coccinellidae**

<i>Cheiromenes sexmaculata</i> (Fabricius, 1781)	Eggs larvae of <i>Achaea janata</i> and <i>Spodoptera litura</i>	India	Basappa (2009)
<i>Micraspis cardoni</i> (Weise, 1892)	<i>Zaniothrips ricini</i>	India	Daniel <i>et al.</i> (1983)
<i>Scymnus</i> sp.	<i>Eutetranychus orientalis</i>	Palestine and Egypt	Klein (1936)
<i>Stethorus</i> sp.	<i>Eutetranychus banksi</i>	USA	McGregor (1914)
<i>Stethorus siphonulus</i> Kapur, 1948	<i>Tetranychus piercei</i>	China	Lui and Lui (1986)
<i>Stethorus histrio</i> Chazeau, 1974	<i>Tetranychus urticae</i>	Chile	Aguilera (1987)

**Hemiptera****Pentatomidae**

<i>Eocanthecona furcellata</i> (Wolff, 1811)	<i>Achaea janata</i>	India	Rao (1977); Usha Rani (2009)
--	----------------------	-------	------------------------------

**Reduviidae**

<i>Rhynocoris kumarii</i> Ambrose and Livingstone, 1986	Eggs larvae of <i>Achaea janata</i> and <i>Spodoptera litura</i>	India	Basappa (2009)
---	--	-------	----------------

**Thysanoptera****Aeolothripidae**

<i>Franklinothrips megalops</i> (Trybom, 1912)	<i>Zaniothrips ricini</i>	India	Daniel <i>et al.</i> (1983)
<i>Myrmorthrips garuda</i> Ramakrishna and Margabandhu, 1931	<i>Zaniothrips ricini</i>	India	Daniel <i>et al.</i> (1983)

**Neuroptera****Chrysopidae**

<i>Chrysoperla carnea</i> (Stephens, 1836)	<i>Tetranychus urticae</i>	India	Rajasekhar <i>et al.</i> (1999)
<i>Chrysoperla</i> sp.	Eggs and larvae of <i>Achaea janata</i> and <i>Spodoptera litura</i>	India	Basappa (2009)

**Mantodea****Mantidae**

<i>Haldimania lilliputana</i> Beier, 1930	<i>Zaniothrips ricini</i>	India	Daniel <i>et al.</i> (1983)
---	---------------------------	-------	-----------------------------

**ARACHNIDA / Acari****Phytoseiidae**

<i>Sciulus</i> sp.	<i>Eutetranychus banksi</i>	USA	McGregor (1914)
--------------------	-----------------------------	-----	-----------------

*maternus* Bhatnagar, 1952, *Rhogas* spp. and *Apanteles hyposidrae* Wilkinson, 1928 were found among larval parasitoids; and pupae were found to be parasitised by *Tetrastichus howardi* (Olliff, 1893) (= *Tetrastichus ayyari* Rohwer, 1921) and *Phorocera* sp. Among insect predators of *A. janata*, *Chrysoperla* sp. and *Cheilomenes sexmaculata* (Fabricius, 1781) were found feeding on the eggs and neonate larvae; other general insect predators like mantids, paper wasps, and sphecid digger wasps were also found preying on larvae; spiders like green lynx spiders, jumping spiders and crab spiders were found feeding on early instar larvae. Among the entomopathogens of *A. janata*, *Spodoptera litura* nucleopolyhedrovirus and granulosis virus were isolated from dead larvae and the fungi *Metarhizium rileyi* (Farl.) Kepler, S.A.Rehner & Humber, 2014 [= *Nomuraea rileyi* (Farlow) Samson, 1974] and *Beauveria bassiana* (Balsamo) Vuillemin were found infecting larvae (Basappa 2009). Many other examples of natural enemies of pests of *R. communis* are shown in Table 4.

### Pest management of phytophagous arthropods in *R. communis*

Pest management methods used to control the principal arthropod pests of *R. communis* include cultural, genetic, ethological, biological, and chemical control.

Cultural control is the use of agronomical practices designed to reduce the presence of pests in crops of *R. communis*. Intercropping is a type of cultural control recommended to diminish the damage caused by insect pests in *R. communis*. Srinivasa Rao *et al.* (2012) found that plants such as *Cyamopsis tetragonoloba* (L.) Taub., 1891, *Vigna unguiculata* (L.) Walp., 1845, *Vigna mungo* (L.) Hepper, 1956, and *Arachis hypogaea* L., 1753, intercropped with *R. communis* in a 1:2 proportion, decreased the incidence of insect pests such as *A. janata*, *E. flavesrens*, and *C. punctiferalis*. Moreover, a more considerable presence of natural enemies of these pests was observed in these intercropping systems. Patel and Patel (2009) recommended intercropping *R. communis* with *Vigna radiata* (L.) Wilczek, 1954, *Sesamum indicum* L., 1753, *Vigna aconitifolia* (Jacq.) Marechal, 1969, and *V. unguiculata*, to reduce damage by *C. punctiferalis*. When *R. communis* was monocropped, *C. punctiferalis* caused 53 % damage, but when intercropped with the above mentioned species, the damage was between 35 and 53 %. Sowing date is another cultural method for reducing damage and the presence of pests. Salihu *et al.* (2014) suggest that the correct time for planting *R. communis* crop must be related to the rainy season, which is more important than any other pest control measure in Africa, since the rains decrease the presence of certain pests.

Genetic control includes the use of cultivars resistant to insect pests, however, according to Singh *et al.* (2015), breeding *R. communis* is complicated by limited sources of pest resistance. In India, there are *R. communis* varieties that are tolerant or resistant to attack by pests of greater economic importance, such as *E. flavesrens*, *T. ricini*, *S. litura*, *A. janata*, *C. punctiferalis*, and *L. trifolii* (Anjani *et al.* 2010; Anjani 2012). Resistant or tolerant plants have high oil content (between 40 and 49 %) and yields that oscillate between 540 and 1,580 kg/ha (Lavanya *et al.* 2012). It is mentioned that the cultivars having purple leaves are resistant to the attack of *L. trifolii*, while those with green leaves are

susceptible (Sarma *et al.* 2006; Anjani *et al.* 2007). Sarma *et al.* (2006) mention that purple-leaf varieties have high levels of anthocyanin, which make the plant more tolerant to *L. trifolii* attack, and the epicuticular wax on their leaves reduces infestation and defoliation by *A. janata* and *S. litura*. There are hybrids, such as GCH4, that are resistant to the attack by *E. flavesrens* due to the high wax content on the plant stems and leaves (Lakshmi *et al.* 2005). Five accessions viz., RG-43, RG-631, RG-1621, RG-3037 and RG-3067, among 165 core set accessions representing diversity in the entire collection maintained at ICAR-Indian Institute of Oilseeds Research, Hyderabad, India, exhibited resistance reaction against *E. flavesrens*; oil content of these accessions was 46, 51, 51, 51, 52 %, respectively (Anjani *et al.* 2018). On the other hand, Severino *et al.* (2012) recommended parallel research to determine the increased potential susceptibility to pests in breeding programs to develop low-ricin, low-ricinine, and low-allergen cultivars to reduce hazardous chemical products found in *R. communis*.

Research is being carried out on the use of transgenic plants of *R. communis*. In India, two transgenic varieties of *R. communis*, Jyothi and VP1, developed by genetic engineering induce *A. janata* mortality above 88 % due to the *Bacillus thuringiensis* gene *CryAb* (Malathi *et al.* 2006).

A little explored method for monitoring and massive trapping of *R. communis* pests has been the use of pheromones, kairomonal attractants and light traps. In India, the pheromone compounds of some pests of economic importance have been identified and used for monitoring and massive trapping of *C. punctiferalis*, *S. litura*, *Amsacta albistriga* Walker, 1864 (Lepidoptera: Arctiidae), *A. janata*, and *S. obliqua* (Cork and Hall 1998). In this country, an important prerequisite for successful management of *S. litura*, the most destructive insect pest of *R. communis* damaging the crop from July- October during the south-west monsoon (kharif season), has been the implementation of an intensive monitoring program of *S. litura* population using sex pheromone traps (Satyagopal *et al.* 2014). Setting twelve traps baited with pheromone compounds per hectare for massive trapping of *S. litura* is recommended (Nandagopal and Rathod 2007; Raghavaiah 2011). In Brazil, researchers are now taking the first steps toward identifying the pheromone compounds of *C. gossypii* (Fregadolli *et al.* 2012) with the aim of developing a commercial pheromone. In India, the kairomonal compounds of the most destructive lepidopteran insect pest of *R. communis*, such as *S. litura*, *A. janata*, and *C. punctiferalis* have been identified for trapping. In field experiment, water trap baited with phenyl acetaldehyde + 2-phenyl ethanol recorded significantly higher moth catches of *S. litura* (6.8 moths/trap/wk) and *C. punctiferalis* (5.8 moths/trap/wk) (Duraimurugan *et al.* 2017). Recently, Duraimurugan and Alivelu (2018), determined the relationship of pheromone trap catches corresponding to the economic threshold level of 25 % defoliation of *S. litura* on *R. communis*, which was estimated to be 81.4 moths/trap/week. Light traps using ultraviolet black-blue spectrum have also been suggested to capture *Phyllophaga* sp. adults as a measure of ethological control Saldarriaga Cardona *et al.* (2011).

Biological control (spraying entomopathogenic microorganisms and releasing entomophagous insects) has been implemented in the control of key *R. communis* pests in countries such as India and Colombia. In India, for example, parasitism rates between 10.4 and 28.7 % of *M. maculipennis* and *Cotesia* sp. were recorded on larvae of *A. janata* and *S.*

*litura*, respectively (Suganthy 2007), while 9.5 % parasitism rates of *Cotesia flavipes* (Cameron, 1891) against *S. obliqua* larvae have been observed (Yadav *et al.* 2010). Also, in India, Rajasekhar *et al.* (1999) mention that when *Chrysoperla carnea* (Stephens, 1836) were released, the *T. urticae* (= *T. telarius*) mite populations diminished by 75 %. In the case of *S. litura*, the release of 150 adults of the parasitoid *Telenomus remus* (Nixon, 1937) per egg mass and the release of the larval parasitoid *Campoletis chlorideae* Uchida, 1957 in a parasitoid: host ratio of 1:15 achieved parasitism rates above 96 % (Satyanarayana *et al.* 2005).

Biological control through entomopathogenic nematodes, bacteria, fungi, and virus exposed to the principal pests of *R. communis* has been evaluated in India. For instance, mortality of *S. litura* pupae was evaluated with two nematode species: *Heterorhabditis indica* Poinar, Karunaka and David, 1992, and *Steinernema carpocapsae* (Weiser, 1955) (Raveendranath *et al.* 2008). Other studies evaluated the mortality of *A. janata* and *Samia ricini* (Drury, 1773) larvae exposed to two species of bacteria: *Bacillus thuringiensis* Berliner, 1915 and *Bacillus cereus* Frankland & Frankland 1887 (Manson *et al.* 1954; Kattegoudar *et al.* 1994; Mathur *et al.* 1994; Vimala Devi and Sudhakar 2006). Duraimurugan *et al.* (2015) conducted research to determine the mortality of *A. janata* larvae and *C. punctiferalis* adults with *Beauveria bassiana* (Balsamo) Vuillemin, 1912 fungus. Mortality of *A. janata* larvae exposed to granulovirus was also evaluated (Naveen Kumar *et al.* 2013). In Colombia, Saldarriaga Cardona *et al.* (2011) reported that the control of *Phyllophaga* sp. larvae was achieved by applying the *B. popilliae* Dutky bacterium at a concentration of 24,000.00 billion spores/ha a year, during five consecutive years.

The use of secondary metabolites derived from plants and other organisms, as well as methods of chemical control, have been assessed for controlling some *R. communis* pests. In laboratory studies, it has been found that methanol extracts of *Clathria longitoxa* (Hentschel, 1912) and *Callyspongia diffusa* (Ridley, 1884), two marine sponges, have insecticidal effect on *A. janata* and *P. ricini* larvae (Joseph *et al.* 2010). Furthermore, *Calotropis gigantea* (L.) W. T. Aiton, 1811, leaf extracts demonstrated strong feeding deterrent activity against larvae of *Pericallia ricini* (Fabricius, 1775) at high concentrations (Neelanarayanan and Indira 2010). Likewise, toxic effects and strong antifeedant activity of raw acetonic extracts of *Mormodica charantia* L., 1753, *Tectona grandis* L. f., 1790, and *Madhuca indica* J. F. Gmel., (1791) against *S. litura* and *A. janata* larvae (Devanand and Usha Rani 2008) were found. Neem, *Azadirachta indica* (Juss, 1830), also evaluated for the control of *S. litura*, induces mortality of larvae at high concentrations (Choudhury and Aizur Rahman 2008).

Chemical control through insecticides is one of the most common practices for control of *R. communis* pests (Gahukar 2018). In Colombia, six pesticides for the control of *C. gossypii* were evaluated. The results of insect control efficacy three days after pesticide application were as follows: (from the least to the most effective) thiamethoxam + lambda-cihalotrin (0.00 %), spinetoram (0.00 %), malathion (20.35 %), thiamethoxam (38.62 %), dimethoate (86.94 %), and imidacloprid (87.33 %); whereas after seven days the following results were obtained: thiamethoxam + lambda-cihalotrin (0.00 %), spinetoram (21.46 %), malathion (38.77 %), thiamethoxam (50.84 %), dimethoate (86.14 %), and imidacloprid (90.37 %) (Varón *et al.* 2010). Mead (1989) suggested the use of carbaryl or malathion

for controlling *C. gossypii* in Florida, USA. In Colombia, Saldarriaga Cardona *et al.* (2011) recommended application of baits poisoned with carbaryl at a dose of 2 to 3 g/L for the control of *A. epsilon* and *S. frugiperda*; the same authors recommended application of liquid chlorpyrifos at the base of the plants at a dose of 1.5 - 2.0 cc/L.

The most recommendable strategy of *R. communis* pest control is Integrated Pest Management (IPM). Most of the IPM programs have been directed against key pests of *R. communis*, such as *S. litura*, *C. punctiferalis*, and *A. janata* (Prabhakar *et al.* 2003; Singh *et al.* 2006; Basappa 2009). In India, the growers increased seed production of *R. communis* up to 28 %, by implementing IPM programs with insecticides, crop rotation, insect traps, application of neem extract, and intercropping (Basappa 2007). The results of research in India demonstrate that IPM is an efficient strategy for the control of *A. janata* and *S. litura*, two of the key pests of *R. communis*. It is possible to decrease populations of these pests by using the recommended IPM program, which includes the use of bird perches for predatory birds to rest and to look for preys, foliar applications of 5 % neem seed extracts, biological insecticide consisting of nuclear polyhedrosis virus (*S. litura* NPV 100 LE/ha), monocrotophos at 0.5 %, and manual removal of larvae (Suganthy 2010). The pest control effectiveness of carbaryl 50W 0.2 %, endosulfan 35 EC 0.05 %, triazophos 40 EC 0.05 %, spinosad 45 SC 0.018 %, fipronil SSC 0.01 %, extract of neem seeds 5 % (weight/volume), *B. thuringiensis* 0.1 %, and a control without applying the dose of 500 L/ha, was evaluated under field conditions 30 and 45 days after establishing a plantation of a *R. communis* variety susceptible to leafminer *L. trifolii*. The results showed that the least damage (lowest number of insect mines) was found when spinosad and triazophos were applied and, at the same time, the best yield was obtained with both treatments (883 and 835 kg seed/ha, respectively) (Akashe *et al.* 2009). On the other hand, natural enemy impact has been proven to be greatest at sites adopting biointensive IPM (BIPM); for example, studies conducted by Basappa (2009) shown that BIPM modules were safer to *A. janata* eggs (*T. chilonis*) and larvae (*M. maculipennis*) parasitoids with 16.1 and 66.1 % average field parasitism, compared to chemical pesticide intensive integrated pest management modules with 6.9 and 21.2 % parasitism, respectively.

## Conclusions

There is a wide range of arthropods that damage *R. communis* in different parts of the world where this plant is cultivated; many of these are considered pests of economic importance. Likewise, there are reports of a great variety of natural enemies, which have been used in biological control programs. According to the coevolutionary hypothesis, it was found that the greatest richness and abundance of arthropods associated with *R. communis* is in Asia and Africa, considered as the center of origin of this plant. Most phytophagous arthropods feed on leaves. The natural enemies with more abundance and richness are the parasitoids that mostly attack the larvae of phytophagous arthropods. With respect to pollinators, *A. mellifera* is the principal pollinating insect, however, more research on pollination and floral ecology in *R. communis* is needed, in order to determine what other floral visitors may act as pollinators, and how they can be protected or manipulated to increase crop yield. The pest management

programs of phytophagous arthropods of *R. communis* must be directed toward promoting and preserving natural enemies and pollinating insects by means of environment-friendly pest management techniques, for which use of wide-spectrum insecticides must be avoided.

### Acknowledgements

We are grateful to Fernando E. Vega [Insect Biocontrol Laboratory, US Department of Agriculture, Agricultural Research Service (USDA-ARS), Beltsville, MD] for suggestions leading to improvement the first version of the manuscript.

### Literature cited

- ABDULLAH, N. M. M.; MARTIN, J. 2007. New record for four additional whiteflies species from Yemen. *Arab Journal of Plant Protection* 25: 33-34.
- AGUILERA, P. A. 1987. New localities for *Stethorus histrionis* Chazeau (Coleoptera Coccinellidae) in Chile. *Revista Chilena de Entomología* 15: 33-36.
- AHUJA, D. B. 1994. Seasonal incidence and chemical control of oriental mite, *Eutetranychus orientalis* (Klein) on castor. *Indian Journal of Entomology* 56: 1-5.
- AKASHE, V. B.; GUD, M. A.; SHINDE, S. K.; DESHPANDE, A. N. 2009. Bio-efficacy of botanicals and chemical insecticides for control of leaf miner (*Liriomyza trifolii* Burges) under dry land condition. *An Asian Journal of Soil Science* 4: 315-317.
- ALI, M. S.; KUMAR, K.; SINGH, R. 2006. Host range of *Oxyrhachis tarandus* Fabricius (Homoptera: Membracidae) in woody trees and shrubs of Bihar. *Environment and Ecology (Kalyani)* 24S: 14-16.
- ÁLVAREZ, P. C.; REYES, F. 1987. Himenópteros entomófagos adultos que se alimentan en los nectarios de algunas malezas comunes en Nuevo León, México. *The Southwestern Entomologist* 12: 205-210.
- AMOABENG, B. W.; GURR, G. M.; GITAU, C. W.; STEVENSON, P. C. 2014. Cost: benefit analysis of botanical insecticide use in cabbage: Implications for smallholder farmers in developing countries. *Crop Protection* 57: 71-76. <https://doi.org/10.1016/j.cropro.2013.11.019>
- ANJANI, K. 2012. Castor genetic resources: A primary gene pool for exploitation. *Industrial Crops and Products* 35 (1): 1-14. <https://doi.org/10.1016/j.indcrop.2011.06.011>
- ANJANI, K.; PALLAVI, M.; SUDHAKARA BABU, S. N. 2007. Uniparental inheritance of purple leaf and the associated resistance to leafminer in castor bean. *Plant Breeding* 126 (5): 515-520. <https://doi.org/10.1111/j.1439-0523.2007.01395.x>
- ANJANI, K.; PALLAVI, M.; SUDHAKARA BABU, S. N. 2010. Biochemical basis of resistance to leafminer in castor (*Ricinus communis* L.). *Industrial Crops and Products* 31 (1): 192-196. <https://doi.org/10.1016/j.indcrop.2009.10.005>
- ANJANI, K.; RAOOF, M. A.; LAKSHMI PRASAD, M. S.; DURAIMURUGAN, P.; LUCOSE, C.; YADAV, P.; PRASAD, R. D.; JAWAHAR LAL, J.; SARADA, C. 2018. Trait-specific accessions in global castor (*Ricinus communis* L.) germplasm core set for utilization in castor improvement. *Industrial Crops and Products* 112: 766-774. <https://doi.org/10.1016/j.indcrop.2018.01.002>
- ANÓNIMO. 1991. Dirección General de Investigación y Extensión Agrícola. Ministerio de Agricultura y Ganadería. San José, Costa Rica. 571 p.
- ANONYMOUS. 1913. Insects injurious to papaw apples in Queensland. *Queensland Agricultural Journal* 27: 33-35.
- ARKHANGEL'SKII, N. N.; ROMANOVA, V. P. 1930. Pests of sunflower and castor in the north Caucasian Region. *Bulletin of the North Caucasian Plant Protection Station* vi-vii: 199-216.
- AUDI, J.; BELSON, M.; PATEL, M.; SCHIER, J.; OSTERLOH, J. 2005. Ricin poisoning. A comprehensive review. *Journal of the American Medical Association* 294 (18): 2342-2351. <https://doi.org/10.1001/jama.294.18.2342>
- BALDWIN, B. S.; COSSAR, R. D. 2009. Castor yield in response to planting date at four locations in the south-central United States. *Industrial Crops and Products* 29 (2-3): 316-319. <https://doi.org/10.1016/j.indcrop.2008.06.004>
- BARNES, D. J.; BALDWIN, B. S.; BRAASCH, D. A. 2009. Degradation of ricin in castor seed meal by temperature and chemical treatment. *Industrial Crops and Products* 29 (2-3): 509-515. <https://doi.org/10.1016/j.indcrop.2008.09.006>
- BARTENEVA, R. V. 1986. Diseases and pests of castor and their control. pp. 101-101. In: Moshkin, V. A. (Ed.). *Castor*. Oxonian Press Pvt Ltd. New Delhi. 284 p.
- BASAPPA, H. 2003. Integrated pest management in Castor. Hyderabad: Directorate of Oilseeds Research (ICAR), India, 52 p.
- BASAPPA, H. 2007. Validation of integrated pest management modules for castor (*Ricinus communis*) in Andhra Pradesh. *The Indian Journal of Agricultural Sciences* 77: 357-362.
- BASAPPA, H. 2009. Impact of integrated pest management modules on the activity of natural enemies in castor ecosystem. *Journal of Biological Control* 23: 221-228.
- BASAPPA, H.; LINGAPPA, S. 2001. Damage potential of *Achaea janata* Linn at different phenological stages of castor. *Indian Journal of Plant Protection* 29: 17-24.
- BAVARESCO, A.; SILVEIRA GARCIA, M.; DIONE GRÜTZMACHER, A.; FORESTI, J.; RINGENBERG, R. 2003. Compared biology of *Spodoptera cosmioides* (Walk.) (Lepidoptera: Noctuidae) in onion, castor oil plant, soybean and bean. *Ciência Rural, Santa Maria* 33 (6): 993-998. <https://doi.org/10.1590/S0103-84782003000600001>
- BIGI, M. F.; TORKOMIAN, V. L. V.; DE GROOTE, S. T. C. S.; HEBLING, M. J. A.; BUENO, O. C.; PAGNOCCA, F. C.; FERNANDES, J. B.; VIEIRA, P. C.; SILVA, M. F. 2004. Activity of *Ricinus communis* (Euphorbiaceae) and ricinine against the leaf-cutting ant *Atta sexdens rubropilosa* (Hymenoptera: Formicidae) and the symbiotic fungus *Leucoagaricus gongylophorus*. *Pest Management Science* 60 (9): 933-938. <https://doi.org/10.1002/ps.892>
- BIRCHLER, J. A.; AUGER, D. L.; RIDDLE, N. C. 2003. In search of the molecular basis of heterosis. *The Plant Cell* 15: 2236-2239. <https://doi.org/10.1105/tpc.151030>
- BOLAND, J. M. 2016. The impact of an invasive ambrosia beetle on the riparian habitats of the Tijuana River Valley, California. *PeerJ* 4: e2141. <https://doi.org/10.7717/peerj.2141>
- BOURNE-MURRIETA, L. R.; WONG-CORRAL, F. J.; BORBOA-FLORES, J.; CINCO-MOROYOQUI, F. J. 2014. Daños causados por el barrenador mayor de los granos *Prostephanus truncatus* (Horn) (Coleoptera: Bostrichidae) en maíz y ramas de plantas silvestres. *Revista Chapingo. Serie ciencias forestales y del ambiente* 20 (1): 63-75. [http://www.scielo.org.mx/scielo.php?script=sci\\_arttext&pid=S2007-40182014000100007](http://www.scielo.org.mx/scielo.php?script=sci_arttext&pid=S2007-40182014000100007)
- CABEZAS, G. F.; MELO, M.; GARCÍA, M. S.; DIEZ-RODRÍGUEZ, G. I.; NAVA, D. E. 2013. Parasitismo de *Trichogramma pretiosum* (Hymenoptera: Trichogrammatidae) sobre *Spodoptera cosmioides* (Lepidoptera: Noctuidae) a diferentes temperaturas. *Revista Colombiana de Entomología* 39: 216-220.
- CANGARDEL, H. 1954. Experiments on the control of the castor bean red spider. *Essais de lutte contre l'araignée rouge du ricin. Terre Mame* 28: 53-55.
- CAVALCANTE, R. D.; PEDROSA, F. N. T.; DE ARAUJO, F. E. 1974. *Pantomorus glaucus* (Perty, 1830), a pest of various crops in the State of Ceará. *Fitossanidade* 1: 22.
- CÉSAR, A. da S.; BATALHA, M. O. 2010. Biodiesel production from castor oil in Brazil: A difficult reality. *Energy Policy* 38 (8): 4031-4039. <https://doi.org/10.1016/j.enpol.2010.03.027>

- CHOUDHURY, P.; AIZUR RAHMAN, S. M. 2008. Neem effects of growth and development of *Spodoptera litura* Fab. *Progressive Research* 3: 2011-2012.
- CORK, A.; HALL, D. R. 1998. Application of pheromones for crop pest management in the Indian Sub-continent. *Journal of Asia-Pacific Entomology* 1 (1): 35-49. [https://doi.org/10.1016/S1226-8615\(08\)60005-9](https://doi.org/10.1016/S1226-8615(08)60005-9)
- DANIEL, A. M.; KUMAR, N. S.; BAKTHAVATSALAM, N. 1983. Bioecology of *Zaniothrips ricini* Bhati (Panchaetothripinae: Terebrantia: Thysanoptera) - A new thrips pest of castor (*Ricinus communis*: Euphorbiaceae). *Proceedings: Animal Sciences* 92 (2): 87-94. <https://doi.org/10.1007/BF03186174>
- DARLING, H. S. 1946. Annual Report of the Agricultural Entomologist. Report. Department of Agriculture, Uganda, pt. 2, pp. 25-30.
- DE ONG, E. R. 1918. Insect pests of the castor bean. *Journal of Economic Entomology* 11 (6): 480. <https://doi.org/10.1093/jee/11.6.480>
- DE SIBIO, P. R.; ROSSI, M. N. 2016. Interaction effect between herbivory and plant fertilization on extrafloral nectar production and on seed traits: An experimental study with *Ricinus communis* (Euphorbiaceae). *Journal of Economic Entomology* 109 (4): 1612-1618. <https://doi.org/10.1093/jee/tow115>
- DELAYA, V. P.; RAJPUT, S. G.; AWATE, B. G.; MOHITE, P. B. 1985. Chemical control of castor semilooper *Achoea janata* Linnaeus. *Indian Journal of Plant Protection* 11: 136-137.
- DEVANAND, P.; USHA RANI, P. 2008. Biological potency of certain plant extracts in management of two lepidopteran pests of *Ricinus communis* L. *Journal of Biopesticides* 1: 170-176.
- DINESH, D. S.; KUMARI, S.; KUMAR, V.; DAS, P. 2014. The potentiality of botanicals and their products as an alternative to chemical insecticides to sandflies (Diptera: Psychodidae): A review. *Journal of Vector Borne Diseases* 51: 1-7.
- DOUKA, C.; TCHUENGUEM, F. N. 2014. Foraging and pollination activity of *Musca domestica* L. (Diptera: Muscidae) on flowers of *Ricinus communis* L. (Euphorbiaceae) at Maroua, Cameroon. *Journal of Biodiversity and Environmental Sciences* 4: 63-76.
- DURAIMURUGAN, P.; ALIVELU, K. 2018. Determination of an action threshold for tobacco caterpillar, *Spodoptera litura* (F.) based on pheromone trap catches in castor (*Ricinus communis* L.). *Journal of Entomological Research* 42 (2): 189-194. <https://doi.org/10.5958/0974-4576.2018.00032.4>
- DURAIMURUGAN, P.; LAKSHMINARAYANA, M.; VIMALA DEVI, P. S. 2015. Comparative efficacy of microbial, botanical and chemical insecticides against lepidopteran pests in castor. *The Ecoscan* 9: 7-10.
- DURAIMURUGAN, P.; SAMPATHKUMAR, M.; SRINIVAS, P. S. 2017. Field evaluation of synthetic kairomonal attractants against major lepidopteran pests of castor. *Journal of Environmental Biology* 38: 1421-1427. <https://doi.org/10.22438/jeb/38/6/MRN-629>
- DURÁN, J. M.; PARRA, N. R.; YUGUEROS, R. M.; DE PAULA QUEIROGA, V. 2010. El cultivo de ricino (*Ricinus communis* L.) en Andalucía: Una alternativa para producción de biodiésel. pp. 39-58. En: Recalde, E. R.; Durán Altisent Posso, J. M. (Eds.). *Cultivos Energéticos Alternativos*. Centro Iberoamericano de Investigación y Transferencia de Tecnología en Oleaginosas (CIITOL), Grupo Seritex, Ecuador. 177 p.
- EGONYU, J. P.; MUKASA, Y.; EKWARU, R.; OGARI, I.; AHUMUZA, G. 2017. Occurrence of *Euwallacea* sp. (Coleoptera: Scolytidae) and its ambrosia fungus *Cunninghamella echinulata* on *Ricinus communis* in *Coffea canephora* gardens in Uganda. *International Journal of Tropical Insect Science* 37 (3): 198-207. <https://doi.org/10.1017/S1742758417000121>
- EL-SHAARAWY, M. F.; GOMAA, A. A.; EL-GARHY, A. T. 1975. Chemical determination and utilization of dietary constituents of two-castor bean varieties by larvae of the Eri silk worm, *Attacus ricini* Boisd. *Zeitschrift für Angewandte Entomologie* 78 (1-4): 171-176. <https://doi.org/10.1111/j.1439-0418.1975.tb04166.x>
- FAOSTAT, Food and Agriculture Organization of the United Nations. 2015. Production—castor oil seeds. In: <http://faostat.fao.org/site/567/DesktopDefault.aspx?PageID=567>. [Review date: 19 February 2017].
- FREGADOLLI, A. M. V.; FREGADOLLI, F. L.; OLIVEIRA, M. S. DE; SILVA, M. J. C. DA; FERREIRA, P. V.; ENDRES, L.; SILVA, R. M.; MACUVELE, D. L. P.; GOULART, H. F.; SANT'ANA, A. E. G.; PIRES, E. V. 2012. Busca da composição feromonal de *Corythucha gossypii*, *Cyclocephala melanocephala* e *Pachycoris torridus*. *EnerBiomassa'12*, Maceió, Alagoas, Brasil, 21 a 23 de Novembro 2012 / Brazil, 21-23 November, 2012. A2.4 (EB 065). 2 p.
- FREITAS, B. M.; CRUZ, D. de O. 2010. As abelhas na polinização de culturas agrícolas. pp. 9-21. In: Cohelo de Lima, M. A. (Ed.). *II Semana dos Polinizadores: Palestras*, 28 e 29 de setembro de 2009, Petrolina, PE, EMBRAPA, Brasil. 82 p.
- FREITAS, B. M.; RIZZARDO, R.A.G.; MILFONT, M. de O. 2009. Pollination and honney production in castor bean (*Ricinus communis* L.) plantations for biofuel. *41<sup>a</sup> Apimondia Congress September 2009*, Montpellier, France. 29 p.
- GAHKAR, R. T. 2010. Bioefficacy of indigenous plant products against pests and diseases of Indian forest trees: A review. *Journal of Forestry Research* 21: 231-238. <https://doi.org/10.1007/s11676-010-0038-7>
- GAHKAR, R. T. 2018. Management of pests and diseases of castor (*Ricinus communis* L.) in India: current status and future perspective. *Archives of Phytopathology and Plant Protection* 51 (17-18): 956-978. <https://doi.org/10.1080/03235408.2018.1541655>
- GALANDE, S. M.; MOTE, U. N.; GHORPADE, S. A. 2005. Host range of serpentine leafminer (*Liriomyza trifolii* Burgess) and its activity in scarcity zone. *Journal of Maharashtra Agricultural Universities* 30: 307-310.
- GEETHA, B.; VENKATESAN, S.; SHANMUGAM, K. 2003. Occurrence of *Helicoverpa armigera* (Hubner) on castor capsules in Tamil Nadu. *Insect Environment* 9: 170-171.
- GHOSH, C. C. 1914. Life-Histories of Indian Insects. *Lepidoptera. Memoirs of the Department of Agriculture* 5: 1-72.
- GOLDEN, M.; FOLLETT, P. A. 2006. First report of *Nezara viridula* f. *aurantiaca* (Hemiptera: Pentatomidae) in Hawaii. *Proceedings of the Hawaiian Entomological Society* 38: 131-132.
- GOVAERTS, R.; FRODIN, D. G.; RADCLIFFE-SMITH, A. 2000. World checklist and bibliography of Euphorbiaceae (with Pandaceae). Redwood Books Limited, Trowbridge, Wiltshire. 1621 p.
- HEDGE, J. N.; CHAKRAVARTHY, A. K.; GANIGAR, P. C. 2009. Screening castor (*Ricinus communis* Linn.) germplasm against leafminer, *Liriomyza trifolii*, semilooper, *Achaea janata* and shoot and capsule borer, *Conogethes punctiferalis* in South Karnataka. *Current Biotica* 3: 386-396.
- HEIL, M. 2008. Indirect defence via tritrophic interactions. *New Phytologist* 178 (1): 41-61. <https://doi.org/10.1111/j.1469-8137.2007.02330.x>
- HEUNGENS, A.; DEGHEELE, D. 1986. Control of the broad mite, *Polyphagotarsoneurus latus* (Banks), with acaricides on *Psophocarpus tetragonolobus* and *Ricinus communis*. *Parasitica* 42: 3-10.
- HUA, Y. Q. 1984. Occurrence and control of *Achaea melicerta* (Drury). *Insect Knowledge (Kunchong Zhishi)* 18: 171-172. <https://doi.org/10.1002/j.1545-7249.1984.tb00512.x>
- HUERTAS DIONISIO, M. 2002. Lepidópteros de Huelva (I) especies detectadas en las márgenes del Río Guadiana. *Boletín de la SAE*, No. 4: 9-29.
- HUSSAIN, M.; KHAN, M. Q. 1966. Record of *Lasioderma serricorne* F. (Anobiidae: Coleoptera) on stored castor beans. *Science and Culture* 32: 212.
- IDRISS, M.; ABDALLAH, N.; AREF, N.; HARIDY, G.; MADKOUR, M. 1997. Biotypes of the castor bean whitefly *Trialeurodes ricini* (Misra) (Hom. Aleyrodidae) in Egypt.

- biochemical characterization and efficiency of geminivirus transmission. *Journal of Applied Entomology* 121 (1-5): 501-509. <https://doi.org/10.1111/j.1439-0418.1997.tb01440.x>
- ISLAM, W.; NARGIS, A.; JOARDER, O. I. 1988. Biology, seasonal occurrence, host range and damage potential of the castor hairy caterpillar, *Euproctis lunata* Walk. (Lymentriidae: Lepidoptera). *Crop Protection* 7 (5): 332-335. [https://doi.org/10.1016/0261-2194\(88\)90081-6](https://doi.org/10.1016/0261-2194(88)90081-6)
- JACOB, P. S.; RAMASUBBARAO, V.; PUNNAIAH, K. C. 2000. Leafhopper fauna associated with oilseed crops in Andhra Pradesh, India. *Pest Management and Economic Zoology* 8: 11-27.
- JANNONE, G. 1952. Studies and researches on agricultural entomology in Eritrea and Ethiopia. VIII. The phytopathological condition of crops on a farm in the Fadis territory (Harar, Ethiopia) with particular reference to an infestation of Aphids on safflower. *Rivista di Agricoltura Subtropicale e Tropicale* 46: 132-137.
- JAYARAJ, S. 1964. Influence of a phytotoxemia on the activities of catalase and free auxins of castor bean varieties in relation to their resistance to *Empoasca flavescens* (F.) (Homoptera, Jassidae). *Zeitschrift für Angewandte Entomologie* 63 (1-4): 32-39. <https://doi.org/10.1111/j.1439-0418.1969.tb04360.x>
- JENA, J.; GUPTA, A. K. 2012. *Ricinus communis* Linn: A phytopharmacological review. *International Journal of Pharmacy and Pharmaceutical Sciences* 4: 25-29.
- JOSEPH, B.; SUJATHA, S.; JEEVITHA, M. V. 2010. Screening of pesticidal activities of some marine sponge extracts against chosen pests. *Journal of Biopesticides* 3: 495-498.
- JUNIOR, E. M. de A.; FERNANDES, I. M. dos S.; SANTOS, C. S.; MESQUITA, L. X.; PEREIRA, R. A.; BORGES MARACAJÁ, P. B.; SOTO-BLANCO, B. 2011. Toxicity of castor bean (*Ricinus communis*) pollen to honeybees. *Agriculture, Ecosystems and Environment* 141 (1-2): 221-223. <https://doi.org/10.1016/j.agee.2011.02.010>
- JYOTHSNA, Y.; KAPIL, M.; USHA, R. 2009. Effects of herbivore feeding on biochemical and nutrient profile of castor bean, *Ricinus communis* L. plants. *Allelopathy Journal* 24: 131-132.
- KALRA, V. K. 1984. *Palexorista parachrysops* parasitizing castor capsule borer. *Plant Protection Bulletin, FAO* 32: 30.
- KATTEGOUDAR, N. F.; ALAGAWADI, A. R.; PAKALE, N. 1994. Efficacy of whole culture of *Bacillus cereus* and culture filtrate against the castor semilooper, *Achaea janata* L. *Journal of Entomological Research (New Delhi)* 18: 223-227.
- KHATTAK, S. U.; JABBAR, A.; HUSSAIN, N. 1991. Studies on host plants of hairy caterpillar *Diacrisia obliqua* Wlk in Peshawar. *Pakistan Journal of Zoology* 23: 297-300.
- KIRKMAN, W.; MOORE, S. 2007. A study of alternative hosts for the false codling moth, *Thaumatotibia* (=*Cryptophlebia*) *leucotreta* in the Eastern Cape. *South African Fruit Journal* 6: 33-38.
- KITTOCK, D. L.; WILLIAMS, J. H. 1970. Effects of plant population on castorbean yield. *Agronomy Journal* 62 (4): 527-529. <https://doi.org/10.2134/agronj1970.00021962006200040030x>
- KLEIN, H. Z. 1936. Contributions to the knowledge of the red spiders in Palestine. I. The oriental red spider, *Anychus orientalis* Zacher. *Bulletin Agricultural Experimental Station, Rehovoth* 21: 2-36.
- KOLTE, S. J. 1995. Castor diseases and crop improvement. Shipra publications, Delhi, India. 119 p.
- LAKSHMAMMA, P.; LAKSHMINARAYANA, M.; LAKSHMI, P.; ALIVELU, K.; LAVANYA, C. 2009. Effect of defoliation on seed yield of castor (*Ricinus communis*). *The Indian Journal of Agricultural Sciences* 79: 620-623.
- LAKSHMI, P.; LAKSHMAMMA, P.; LAKSHMINARAYANA, M. 2010. Contribution of upper leaves to seed yield of castor. *Journal of Oilseeds Research* 27: 209-212.
- LAKSHMI, P. V.; SATYANARAYANA, J.; SINGH, H.; RATNASUDHAKAR, T. 2005. Incidence of green leafhopper, *Empoasca flavescens* Fab., on castor, *Ricinus communis* L., in relation to morphological characters and date of sowing. *Journal of Oilseeds Research* 22: 93-99.
- LAMA, A. D.; VUORISALO, T.; NIEMELÄ, P. 2015. Global patterns of arthropod herbivory on an invasive plant, the physic nut (*Jatropha curcas* L.). *Journal of Applied Entomology* 139 (1-2): 1-10. <https://doi.org/10.1111/jen.12161>
- LAVANYA, C.; MURTHY, I. Y. L. N.; NAGARAJ, G.; MUKTA, N. 2012. Prospects of castor (*Ricinus communis* L.) genotypes for biodiesel production in India. *Biomass and Bioenergy* 39: 204-209. <https://doi.org/10.1016/j.biombioe.2012.01.008>
- LIMA, R. L. S.; SEVERINO, L. S.; SAMPAIO, L. R.; SOFIATTI, V.; GOMES, J. A.; BELTRÃO, N. E. M. 2011. Blends of castor meal and castor husks for optimized use as organic fertilizer. *Industrial Crops and Products* 33 (2): 364-368. <https://doi.org/10.1016/j.indcrop.2010.11.008>
- LINE, S.; HONG, K.; CARMONA-GALINDO, V. D. 2013. The relationship between castor bean stem diameter and extrafloral nectary gland size. *Bios* 84 (2): 98-100. <https://doi.org/10.1893/0005-3155-84.2.298>
- LIU, X.; LI, D. 2006. Biological activity of ricinine and outlook of its applied development. *Chinese Journal of Pharmacology and Toxicology* 20 (1): 76-78.
- LOHAR, M. K.; JUNO, G. M.; LANJAR, A. G.; RAHOO, G. M.; MAHAR, A. N. 1997. Seasonal population and host plants of *Spodoptera litura* (F.) in lower Sindh. *Pakistan Entomologist* 19: 53-57.
- LONDÓN-ZULUAGA, M. 2008. Plagas. pp. 15-17. In: Navas, A. A.; Córdoba, O.; Bran, A.; Macias, F. (Eds.). *Higuerilla: Alternativa Productiva, Energética y Agroindustrial para Colombia*. Corpocaja, Rionegro, Antioquia, Colombia. 34 p.
- LÓPEZ-GUILLÉN, G.; GÓMEZ RUIZ, J.; BARRERA-GAYTÁN, J. F.; HERNÁNDEZ-GÓMEZ, E.; ZAMARRIPA-COLMENERO, A. 2012. Artrópodos asociados a *Jatropha curcas* L. y *Ricinus communis* L. en Chiapas. *Entomología Mexicana* 11: 375-380.
- LOURENÇÃO, A. L.; DE CARVALHO, L. O.; DE CAMPOS LASCA, E. D. H. 1989. *Anacraga citrinopsis* Dyar (Lepidoptera: Dalceridae) em mamoneira no estado de São Paulo. *Bragantia Campinas* 48 (1): 109-112. <https://doi.org/10.1590/S0006-87051989000100010>
- LU, Y.; WU, K.; WYCKHUYS, K. A. G.; GUO, Y. 2010. Overwintering hosts of *Apolygus lucorum* (Hemiptera: Miridae) in northern China. *Crop Protection* 29 (9): 1026-1033. <https://doi.org/10.1016/j.cropro.2010.03.017>
- LUI, Z. G.; LUI, N. Z. 1986. A preliminary report on *Tetranychus piercei* McGregor. *Insect Knowledge* 23:18-19.
- MALATHI, B.; RAMESH, S.; RAO, K. V.; REDDY, V. D. 2006. Agrobacterium-mediated genetic transformation and production of semilooper resistant transgenic castor (*Ricinus communis* L.). *Euphytica* 147: 441-449. <https://doi.org/10.1007/s10681-005-9043-x>
- MANSON, E. E. D.; POLLOCK, M. R.; TRIDGELL, E. J. 1954. A comparison of the properties of penicillinase produced by *Bacillus subtilis* and *Bacillus cereus* with and without addition of penicillin. *Journal of General Microbiology* 11 (3): 493-505. <https://doi.org/10.1099/00221287-11-3-493>
- MARTÍN, C.; MOURE, A.; MARTÍN, G.; CARRILLO, E.; DOMÍNGUEZ, H.; PARAJÓ, J. C. 2010. Fractional characterisation of jatropha, neem, moringa, trisperma, castor and candlenut seeds as potential feedstocks for biodiesel production in Cuba. *Biomass and Bioenergy* 34 (4): 533-538. <https://doi.org/10.1016/j.biombioe.2009.12.019>
- MARTÍNEZ, M. A.; BLANCO, E.; SURÍS, M. 2005. Fauna de chinches harinosas (Hemiptera: Coccoidea) asociada a plantas de interés: I. plantas arbóreas. *Revista de Protección Vegetal* 20: 125-127.
- MATHUR, Y. K.; ALAM, M. A.; KUMAR, J. 1994. Effectiveness of different formulations of *Bacillus thuringiensis* Berliner against

- Pericallia ricini* Fabricius (Lepidoptera: Arctiidae). Journal of Entomological Research (New Delhi) 18: 95-104.
- MAU, R. F. L.; KESSING, J. L. M. 2007. *Achaea janata* (Linnaeus). Available in: <http://Achaea.janata.mht> [Review date: 05 October 2015].
- McGREGOR, E. A. 1914. Four new tetranychids. Annals of the Entomological Society of America 7 (4): 354-364. <https://doi.org/10.1093/aesa/7.4.354>
- McQUATE, G. T.; JAMESON, M. L. 2011. Control of Chinese rose beetle through the use of solar-powered nighttime illumination. Entomologia Experimentalis et Applicata 141 (3): 187-96. <https://doi.org/10.1111/j.1570-7458.2011.01186.x>
- MEAD, F. W. 1989. Cotton lace bug, *Corythucha gossypii* in Florida (Hemiptera: Tingidae). Fla. Dept. Agr. & Consumer Serv. Divison of Plant Industry. Entomology Circular No. 324: 4.
- MILLER, L. T.; NAGAMINE, W. T. 2005. First records of *Corythucha gossypii* (Hemiptera: Tingidae) in Hawaii, including notes on host plants. Proceedings of the Hawaiian Entomological Society 37: 85-88.
- MOLL, R. H.; SALHUANA, W. S.; ROBINSON, H. F. 1962. Heterosis and genetic diversity in variety crosses of maize. Crop Science 2 (3): 197-198. <https://doi.org/10.2135/cropsci1962.0011183X000200030005x>
- MONA, H. A.; MOHAMED, H. A.; HAFEZ, S. F. M. 2005. Biological and physiological effects of bioinsecticide Spinosad on the cutworm, *Agrostis ipsilon* (Hüfnagel). Egyptian Journal of Biological Pest Control 15: 139-145.
- MONTEIRO, A. H. R. R.; GOMES, S.; GOMES, I.; QUEIROZ, P. R.; LIMA, L. H. C.; OLIVEIRA, M. R. V. 2005. Current status of the whitefly *Aleurodicus dispersus* as an invasive pest in the Cape Verde Islands. pp. 261-262. In: Alford, D. V.; Backhaus, G. F. (Eds.). Plant protection and plant health in Europe: Introduction and spread of invasive species, held at Humboldt University, Berlin, Germany, 9-11 June 2005, 441 p.
- NAIK, M. I.; KUMAR, M. A. A.; MANJUNATHA, M.; SHIVANNA, B. K. 2010. Survey for the pests of castor and natural enemies of castor semilooper. Environment and Ecology 28: 558-563.
- NANDAGOPAL, V.; RATHOD, V. 2007. Development of efficient pheromone trap for field catch of *Spodoptera litura* (F.) in groundnut and castor ecosystem. Journal of Applied Zoological Researches 18: 29-32.
- NASS, L. L.; PEREIRA, P. A. A.; ELLIS, D. 2007. Biofuels in Brazil: An overview. Crop Science 47 (6): 2228-2237. <https://doi.org/10.2135/cropsci2007.03.0166>
- NAVATHA, L.; SREEDEVI, K. 2012. Insect pollinator diversity and abundance in Castor, *Ricinus communis* L. Current Biotica 6: 251-253.
- NAVEEN KUMAR, P.; PRASAD, Y. G.; PRABHAKAR, M.; PHANIDHARA, A.; VENKATESWARLU, B. 2013. Granulovirus of semilooper, *Achaea janata* L. (Lepidoptera: Noctuidae): its bioefficacy and safety in mammalian toxicity tests. Journal of Biological Control 27: 99-104.
- NEELANARAYANAN, P.; INDIRA, P. 2010. Effect of *Calotropis gigantean* leaves extract on the feeding activities of pest of castor woolly bear, *Pericallia ricini*. Insect Environment 16: 39-40.
- NEGREIROS, J.; VENDRAMIN, J. D.; RODRIGUES, M. E. V.; MENDONCA, G. A. 1998. Comparison of the biology of the eri silkworm *Philosamia ricini* (Drury) (Lepidoptera, Saturniidae) on four castor bean genotypes. Revista de Agricultura (Piracicaba) 73: 155-166.
- OSHAIBAH, A. A.; BADR, M. A.; HUSSEIN, H. R.; AL-GAMAL, M. M. 1986. Identification of *Sathrobrota rileyi* (Wals.) (Lep.-Cosmopterigidae as a new record in Egypt). Agricultural Research Review 61: 273-283.
- PARKMAN, P.; DUSKY, J. A.; WADDILL, V. H. 1989. Biological Studies of *Liriomyza sativae* (Diptera, Agromyzidae) on castor bean. Environmental Entomology 18 (5): 768-772. <https://doi.org/10.1093/ee/18.5.768>
- PARMAR, P. D.; VYAS, H. J.; RATHOD, R. R. 2006. Bionomics of leaf hopper, *Jacobiasca furcostylus* (Ramakrishnan and Menon), on castor. Indian Journal of Entomology 68: 107-112.
- PATEL, B. S.; PATEL, I. S. 2009. Management of shoot and capsule borer, *Conogathes punctiferalis* L in castor by intercropping. Trends in Biosciences 2 (2): 66-67.
- PATEL, P. V.; ACHARYA, M. F.; GEDIA, M. V.; VYAS, H. J. 2009. Study on growing degree days and population dynamics of thrips, *Scirtothrips dorsalis* Hood (Thysanoptera: Thripidae) on castor. Journal of Agrometeorology 11: 79-82.
- PAUL, S. K.; JHA, S.; GHOSH, M. R. 2000. Morphometric studies of *Pericallia ricini* F. (Arctiidae) and *Euproctis fraterna* (Moore) (Lymantriidae) infesting castor (*Ricinus communis* L.). Uttar Pradesh Journal of Zoology 20: 69-71.
- POMARI, A. F.; BUENO, A. F.; BUENO, R. C. O. F.; MENEZES-JR, A. O. 2013. *Telenomus remus* Nixon egg parasitization of three species of *Spodoptera* under different temperatures. Neotropical Entomology 42: 399-406. <https://doi.org/10.1007/s13744-013-0138-0>
- PONS, X.; LUMBIERRES, B.; GARCÍA, S.; MANETTI, P. L. 2002. *Metcalfa pruinosa* (Say) (Homoptera: Flatidae), ¿una plaga potencial de plantas ornamentales en espacios verdes urbanos de Cataluña? Boletín de Sanidad Vegetal Plagas 28: 217-222.
- PRABHAKAR, M.; SRINIVASA, R. M.; PRASAD, Y. G. 2003. Evaluation of bio-intensive integrated pest management modules against castor semilooper, *Achaea janata* Linn. Indian Journal of Plant Protection 31: 56-58.
- PRABHAKAR, M.; PRASAD, Y. G.; VENKATESWARLU, B. 2010. New record of *Hexameritis dactylocercus* Poinar and Linares (Nematoda: Mermithidae) parasitizing red hairy caterpillar, *Amsacta albistriga* (Walker) (Lepidoptera: Arctiidae) from India. Journal of Biological Control 14: 385-387.
- RAGHAVAIAH, G. 2011. Pests of crops and their management 2011-12. Department of Entomology Agricultural College, Bapatla, Acharya N G Ranga Agricultural University, Faculty of Agriculture, the India. 177 p.
- RAJASEKHAR, D. W.; RACHAPPA, V. H.; AWAKNAVAR, J. S. 1999. Role of *Chrysoperla carnea* Stephens and insecticides in suppression of castor mite. Insect Environment 4: 151.
- RAO, P. S. 1977. New records of predatory bugs on Castor semilooper, *Achaea janata* L. (Lepidoptera: Arctiidae) on soybean. Current Science 44: 481-482.
- RAOOOF, M. A.; LAKSHMINARAYANA, M.; CHANDER RAO, S. 2003. Crop Protection. pp. 50-73. In: Hegde, D. M.; Sujatha, M.; Singh, N. B. (Eds.). Castor in India. Directorate of Oilseeds Research, Rajendranagar, Hyderabad, India. 118 p.
- RAVEENDRANATH, S.; KRISHNAYYA, P. V.; RAO, P. A.; MURTHY, K. V.; HUSSAINI, S. S. 2008. Bioefficacy of entomopathogenic nematodes, *Steinernema carpocapsae* and *Heterorhabditis indica* against pupae of *Spodoptera litura*. Indian Journal of Plant Protection 36: 288-291.
- REF, J. C.; GUMPERT, F. M.; FISCHER, S.; MELCHINGER, A. E. 2007. Impact of interpopulation divergence on additive and dominance variance in hybrid populations. Genetics 176 (3): 1931-1934. <https://doi.org/10.1534/genetics.107.074146>
- RIBEIRO, L. do P.; COSTA, E. C. 2008. Occurrence of *Erinnyis ello* and *Spodoptera marima* in castor bean plantation in Rio Grande do Sul State, Brazil. Ciencia Rural 38 (8): 2351-2353. <https://doi.org/10.1590/S0103-84782008000800040>
- RIZZARDO, R. A. G.; MILFONT, M. O.; SILVA, E. M. S.; FREITAS, B. M. 2012. *Apis mellifera* pollination improves agronomic productivity of anemophilous castor bean (*Ricinus communis*). Anais da Academia Brasileira de Ciências 84 (4): 1137-1145, <https://doi.org/10.1590/S0001-37652012005000057>
- ROSSI, G. D.; SANTOS, C. D.; CARVALHO, G. A.; ALVES, D. S.; PEREIRA, L. L. S.; CARVALHO, G. A. 2012. Biochemical analysis of a castor bean leaf extract and its insecticidal effects

- against *Spodoptera frugiperda* (Smith) (Lepidoptera: Noctuidae). *Neotropical Entomology* 41: 503-509. <https://doi.org/10.1007/s13744-012-0078-0>
- SAILAJA, M.; TARAKESWARI, M.; SUJATHA, M. 2008. Stable genetic transformation of castor (*Ricinus communis* L.) via particle gun-mediated gene transfer using embryo axes from mature seeds. *Plant Cell Reports* 27: 1509-1519. <https://doi.org/10.1007/s00299-008-0580-3>
- SALDARRIAGA CARDONA, A.; LONDOÑO ZULUAGA, M. E.; CÓRDOBA GAONA, O. de J. 2011. Problemas fitosanitarios asociados al cultivo de higuerilla en Colombia. Corporación Colombiana de Investigación Agropecuaria, CORPOICA, Centro de Investigación La Selva, Rionegro, Antioquia, Colombia. Cartilla Divulgativa. 52 p. Available in: <http://bibliotecadigital.agronet.gov.co/bitstream/11348/6461/1/Revista%20Cultivo%20de%20Higuera.pdf> [Review date: December 2017]
- SALIHU, B. Z.; GANA, A. K.; APUYOR, B. O. 2014. Castor oil plant (*Ricinus communis* L.): Botany, ecology and uses. *International Journal of Science and Research* 3: 2319-7064.
- SARMA, A. K.; SINGH, M. P.; SINGH, K. I. 2005. Studies of insect-pests of castor in the agro-ecosistem of Manipur. *Journal of Applied Zoological Researches* 16: 164-165.
- SARMA, A. K.; SINGH, M. P.; SINGH, K. I. 2006. Resistance of local castor genotypes to *Achaea janata* Linn. and *Spodoptera litura* Fabr. *Journal of Applied Zoological Researches* 17: 179-181.
- SATYAGOPAL, K.; SUSHIL, S. N.; JEYAKUMAR, P.; SHANKAR, G.; SHARMA, O. P.; BOINA, D. R.; SAIN, S. K.; RAM, A.; KAPOOR, K. S.; SANJAY, A.; SUBHASH, K.; PATNI, C. S.; SURESH, D. E.; RAJASHEKARAPPA, K.; LAKSHMINARAYANA, M.; NARAYANASWAMY, H.; SHIVANNA, B. K. 2014. AESA based IPM package for castor. National Institute of Plant Health Management. Department of Agriculture and Cooperation, Ministry of Agriculture. Government of India. Rajendranagar, Hyderabad, Telangana, India. 384 p.
- SATYANARAYANA, J.; SING, T. V. K. 2003. Outbreak of *Helicoverpa armigera* (Hubner) on castor in Southern Telangana Region of Andhra Pradesh. *Insect Environment* 8: 176-177.
- SATYANARAYANA, J.; BALLAL, C. R.; RAO, N. S. 2005. Evaluation of egg parasitoid, *Telenomus remus* Nixon and larval parasitoid, *Campoplexis chlorideae* Uchida on *Spodoptera litura* (Fabricius) in castor. *Indian Journal of Plant Protection* 33: 26-29.
- SAVY, A. F. 2005. Mamona tecnologia agrícola. Campinas-SP, Brasil. EMOP. 105 p.
- SEVERINO, L. S.; MILANI, M.; DE ALMEIDA, M. C. R.; DE SOUZA, G. T. M.; CARDOSO, G. D. 2006. Avaliação da produtividade e teor de óleo de dez genótipos de mamoneira cultivados em altitude inferior a 300 metros. *Revista Ciência Agronómica* 37: 188-194.
- SEVERINO, L. S.; FREIRE, M. A. O.; LUCENA, A. M. A.; VALE, L. S. 2010. Sequential defoliations influencing the development and yield components of castor plants (*Ricinus communis* L.). *Industrial Crops and Products* 32 (3): 400-404. <https://doi.org/10.1016/j.indcrop.2010.06.007>
- SEVERINO, L. S.; AULD, D. L.; BALDANZI, M.; CÂNDIDO, M. J. D.; CHEN, G.; CROSBY, W.; TAN, D.; HE, X.; LAKSHMAMMA, P.; LAVANYA, C.; MACHADO, O. L. T.; MIELKE, T.; MILANI, M.; MILLER, T. D.; MORRIS, J. B.; MORSE, S. A.; NAVAS, A. A.; SOARES, D. J.; SOFIATTI, V.; WANG, M. L.; ZANOTTO M. D.; ZIELER, H. 2012. A review on the challenges for increased production of castor. *Agronomy Journal* 104 (4): 853-880. <https://doi.org/10.2134/agronj2011.0210>
- SHALABY, F. F.; IBRAHIM, A. A.; KARES, E. A. 1988. Effect of parasitism by *Microplitis rufiventris* Kok. on the susceptibility of *Spodoptera littoralis* (Boisd.) larvae to bolstar 702 E.C. *Bulletin of the Entomological Society of Egypt, Economic Series* 15: 165-172.
- SHARMA, A.; SINGH, R. 2002. Oviposition preference of cotton leafhopper in relation to leaf-vein morphology. *Journal of Applied Entomology* 126 (10): 538-544. <https://doi.org/10.1046/j.1439-0418.2002.00697.x>
- SHARMA, M. L.; SHUKLA, A.; REDDY, R. K. 1995. Field testing of castor cultivars for their yield potential and resistance against shoot and capsule borer, *Dichocrocis punctiferalis* Guenea. *Crop Research (Hisar)* 10: 54-58.
- SINGH, A. S.; KUMARI, S.; MODI, A. R.; GAJERA, B. B.; NARAYANAN, S.; KUMAR, N. 2015. Role of conventional and biotechnological approaches in genetic improvement of castor (*Ricinus communis* L.). *Industrial Crops and Products* 74: 55-62. <https://doi.org/10.1016/j.indcrop.2015.05.001>
- SINGH, G.; GREWAL, G. S. 1982. Chemical control of the Bihar hairy caterpillar *Diacrisia obliqua* Walker, on castor and sesame. *International Journal of Entomology, India* 1: 69-71.
- SINGH, S.; SINGH, S. K.; SUDHAKAR, R. 2006. Comparative performance of different pest management treatments for castor+pigeonpea intercropping system under rainfed conditions. *Research on Crops* 7: 548-551.
- SINGH, S.; SINGH, S. K.; SUDHAKAR, R. 2008. Validation of integrated pest management module for castor and pigeonpea intercropping system for rainfed agroecosystem. *Pesticide Research Journal* 20: 217-220.
- SINGH, T. V. K.; SINGH, K. M.; SINGH, R. N. 1991. Host range of groundnut jassid, *Empoasca kerri* Pruthi. *Indian Journal of Entomology* 53: 1-17.
- SINGH, U. D.; AGARWAL, D. P.; PANDEY, P. N. 1989. *Eclipta alba* and *Euphorbia hirta* as new food plants and foci of infestation of *Amsacta moorei* Butl. (Lepidoptera: Arctiidae). *Bulletin of Entomology (New Delhi)* 30: 238-239.
- SOARES, J. J.; ARAÚJO, L. H. A.; BATISTA, F. A. S. 2001. Pragas e seu controle. pp. 213-227. In: Azevedo, D. M. P.; Beltrão, N. E. M. (Eds.). *O agronegócio da mamona no Brasil*. Embrapa Algodão/Emrappa Informação Tecnológica, Campina Grande, Brasília, Brazil. 350 p.
- SOUZA-SILVA, M.; FONTENELLE, J. C. R.; MARTINS, R. P. 2001. Seasonal abundance and species composition of flower-visiting flies. *Neotropical Entomology* 30 (3): 351-359. <https://doi.org/10.1590/S1519-566X2001000300002>
- SPAIN, L. A. 1940. Some reactions of grasshoppers to castor bean plants. *Iowa State College Journal Science* 14: 353-355.
- SRINIVASA RAO, M.; RAMA RAO, C. A.; SRINIVAS, K.; PRATIBHA, G.; VIDYA SEKHAR, S. M.; SREE VANI, G.; VENKATESWARLU, B. 2012. Intercropping for management of insect pests of castor, *Ricinus communis*, in the semi-arid tropics of India. *Journal of Insect Science* 12 (1): 1-10. <https://doi.org/10.1673/031.012.1401>
- SUGANTHY, M. 2007. Survey and monitoring the incidence of pests of castor. *The Madras Agricultural Journal* 94: 133-135.
- SUGANTHY, M. 2010. Integrated pest management strategies against castor defoliators. *The Madras Agricultural Journal* 97 (7/9): 278-280.
- SUJATHA, M.; VIMALA DEVI, P. S.; REDDY, T. P. 2011. Insect pests of castor (*Ricinus communis* L.) and their management strategies. pp. 177-198. In: Dashavantha Reddy, V.; Nagaraja Rao, P.; Venkateswara Rao, K. (Eds.). *Pests and pathogens: Management Strategies*. BS Publications, CRC Press, India, 590 p.
- TURLINGS, T. C. J.; WÄCKERS, F. 2004. Recruitment of predators and parasitoids by herbivore-injured plants. pp. 21-75. In: Cardé, R. T.; Millar, J. G. (Eds.). *Advances in insect chemical ecology*. Cambridge University Press. Cambridge University Press. 352 p. <https://doi.org/10.1017/CBO9780511542664.003>
- USHA RANI, P. 2009. Sensillary morphology on the rostral apex and their possible role in prey location behaviour of the carnivorous stinkbug, *Eocanthecona furcellata* (Wolff) (Heteroptera: Pentatomidae). *Acta Zoologica (Stockholm)* 90 (3): 246-253. <https://doi.org/10.1111/j.1463-6395.2008.00346.x>

- USHA RANI, P.; PRATYUSHA, S. 2014. Role of castor plant phenolics on performance of its two herbivores and their impact on egg parasitoid behaviour. *BioControl* 59: 513-524. <https://doi.org/10.1007/s10526-014-9590-y>
- USHA RANI, P.; RAJASEKHARREDDY, P. 2009. Toxic and antifeedant activities of *Sterculia foetida* (L.) seed crude extract against *Spodoptera litura* (F.) and *Achaea janata* (L.). *Journal of Biopesticides* 2: 161-164.
- USHA RANI, P.; SUDHEER, S. D.; PADMINI, G.; LAVANYA, C. 2006. Nutrient-allelochemical interactions of castor, *Ricinus communis* (L.) plants, resistant and susceptible to infestation of green leaf hopper, *Empoasca flavescens* (Fabr.). *Journal of Applied Zoological Researches* 17: 1-8.
- VANDENBORRE, G.; SMAGGHE, G.; VAN DAMME, E. J. M. 2011. Plant lectins as defense proteins against phytophagous insects. *Phytochemistry* 72 (13): 1538-1550. <https://doi.org/10.1016/j.phytochem.2011.02.024>
- VARÓN, E. H.; MOREIRA, M. D.; CORREDOR, J. P. 2010. Efecto de *Corythucha gossypii* sobre las hojas de higuerilla: criterios para su muestreo y control con insecticidas. *Corpoica Ciencia y Tecnología Agropecuaria* 11 (1): 41-47. [https://doi.org/10.21930/rcta.vol11\\_num1\\_art:193](https://doi.org/10.21930/rcta.vol11_num1_art:193)
- VIMALA DEVI, P. S. V.; SUDHAKAR, R. 2006. Effectiveness of a local strain of *Bacillus thuringiensis* in the management of castor semilooper, *Achaea janata* on castor (*Ricinus communis*). *Indian Journal of Agricultural Sciences* 76: 447-449.
- WÄCKERS, F. L.; ZUBER, D.; WUNDERLIN, R.; KELLER, F. 2001. The Effect of herbivory on temporal and spatial dynamics of foliar nectar production in cotton and castor. *Annals of Botany* 87 (3): 365-370. <https://doi.org/10.1006/anbo.2000.1342>
- WANG, X. S.; CHEN, Q. Z.; ZHANG, S. Z.; LIU, T. X. 2016. Parasitism, host feeding and immature development of *Encarsia formosa* reared from *Trialeurodes vaporariorum* and *Bemisia tabaci* on *Trialeurodes ricini*. *Journal of Applied Entomology* 140 (5): 346-352. <https://doi.org/10.1111/jen.12271>
- WATERS, T.; CHIRIKIAN, D.; CARMONA-GALINDO, V. D. 2014. Insect visitation of peduncular and petiolar extrafloral nectar glands on castor bean (*Ricinus communis* L.) plants in southern California. *Journal of Evolutionary Biology Research* 6 (2): 5-8. <https://doi.org/10.5897/JEBR2014.0058>
- WENE, G. P. 1933. Injurious insects found on castor beans. *Journal of Economic Entomology* 48 (1): 110. <https://doi.org/10.1093/jee/48.1.110>
- WOLOOTT, G. N. 1917. Report of the entomologist. Fifth Rept. Bd. Comm. Agr. P. R. for the period from 1st July, 1915 to 30th June, 1916.
- YADAV, R.; YADAV, N.; YADAV, R.; KATIYAR, R. R. 2010. Natural parasitization by certain parasitoids on the pests of field crops. *International Journal of Plant Protection* 3: 408-409.
- YASUR, J.; MATHUR, K.; RANI, P. 2009. Effects of herbivore feeding on biochemical and nutrient profile of castor bean, *Ricinus communis* L. plants. *Allelopathy Journal* 24: 131-142.
- YLLA, J.; MACIÀ, R.; HUERTAS DIONISIO, M. 2008. Piráldos y Crámbidos detectados en Almería, España (Lepidoptera: Pyraloidea). *SHILAP Revista de Lepidopterología* 36: 191-204.
- YOUNIS, A. M. 1992. Effect of constant temperatures on development and survival of the immature stages of the black cutworm *Agrotis ipsilon* (Hüfnagel) Lepidoptera: Noctuidae. *Assiut Journal of Agricultural Sciences* 23: 291-301.
- ZHANG, H. J.; DUAN, G. Q.; ZHANG, Z. B.; LIANG, Z. J.; ZHIANG, D. M.; XU, Q.; WANG, X. M.; XU, A. L.; LIU, Z. 2006. Effect of leaf mining by *Liriomyza sativa* larvae Photosynthesis of some crops. *Acta Entomologica Sinica* 40: 100-105.

### Origin and funding

This work was part of a project on pests of *Ricinus communis* in Mexico and was supported by Secretaría de Agricultura, Ganadería, Desarrollo Rural, Pesca y Alimentación (SAGARPA, Mexico).

### Author contribution

Guillermo López-Guillén, Jaime Gómez Ruiz and Juan F. Barrera defined the content of the study, conducted the literature review and wrote the manuscript. All authors read and approved the final manuscript.