

Influence of temperature and duration of storage on the lipid reserves of entomopathogenic nematodes

Influencia de la temperatura y del tiempo de almacenaje sobre las reservas lipídicas de nemátodos entomopatógenos

VANESSA ANDALÓ¹, ALCIDES MOINO JR.², CLEBER MAXIMINIANO³, VICENTE P. CAMPOS⁴,
and LÚCIA A. MENDONÇA⁵

Abstract: Lipids represent the main source of energy in entomopathogenic nematodes. In the infective juvenile (IJ) phase, the level of such reserves can be influenced by storage, and this may affect their infectivity. The aim of this study was to evaluate the percentage of lipids and the associated infectivity in IJs of *Steinernema carpocapsae*, *S. riobrave*, *Heterorhabditis* sp. JPM4, *Heterorhabditis* sp. CCA and *Heterorhabditis* sp. PI that had been stored under different temperatures (8-28°C) for various times (0 to 180 days). The amounts of lipids present in IJs were evaluated histologically using a colorimetric method, while infectivity was assayed against *Galleria mellonella* larvae. Lipid levels diminished with increasing storage time for all nematodes, but the rates of decrease varied according to storage temperature and species. Lipid reserves were conserved for longer storage periods at 8, 16 and 20°C, while at 24 and 28°C the percentage of lipids decreased rapidly. The infectivities of IJs of *Heterorhabditis* spp. were less tolerant than those of *Steinernema* spp. to temperatures of 8, 24 and 28°C. Thus, while storage at 8°C was optimal for conserving lipid reserves, infectivity was best preserved at temperatures of 16 and 20°C gave rise to the least reduction in infectivities after 180 days of storage. In this way, lipids and infectivity are influenced by different storage temperatures for the species tested. These data are useful for greater success in using entomopathogenic nematodes as biocontrol agents.

Key words: Biological control. Heterorhabditidae. Infectivity. Steinernematidae

Resumen: La principal fuente de energía de los nemátodos entomopatógenos (NEP) en su fase infectiva son los lípidos, el nivel de esas reservas puede estar influenciado por el almacenamiento e interferir en su infectividad. El objetivo de este trabajo fue evaluar el porcentaje de lípidos en juveniles infectantes (JI) almacenados en diferentes temperaturas y asociar esto con su infectividad. Suspensiones de *Steinernema carpocapsae*, *S. riobrave*, *Heterorhabditis* sp. JPM4, *Heterorhabditis* sp. CCA y *Heterorhabditis* sp. PI fueron incubadas por cero, 15, 30, 60, 90, 120, 150 y 180 días a 8, 16, 20, 24 y 28°C. La cantidad de lípidos se evaluó a través del método colorimétrico. Para cada tratamiento se evaluó la infectividad de los JI en *Galleria mellonella*. Todos nemátodos presentaron reducción de lípidos a lo largo del tiempo, sin embargo, hubo variación de acuerdo con la temperatura. Los nemátodos mantuvieron sus lípidos por mayor tiempo en 8, 16 y 20°C, ya en 24 y 28°C el porcentaje disminuyó rápidamente. Los JI de *Heterorhabditis* se mostraron menos tolerantes a las temperaturas de 8, 24 y 28°C comparados con los del género *Steinernema* en relación a la infectividad. Las temperaturas de 16 y 20°C fueron las que menos afectaron la infectividad al final de los 180 días. De esta manera, los lípidos y la infectividad están influenciados por el almacenamiento a diferentes temperaturas, en las especies experimentadas. Estos datos son de gran utilidad para tener éxito en el uso de nemátodos entomopatógenos como agentes de control biológico.

Palabras clave: Control biológico. Heterorhabditidae. Infectividad. Steinernematidae

Introduction

Entomopathogenic nematodes (EPNs) in the families Steinernematidae and Heterorhabditidae are considered particularly useful in the biological control of insect pests. The ubiquitous distribution of EPNs reflects their ability to adapt and survive under stress conditions, including changes in osmotic tension, temperature or desiccation, and the presence of chemicals and predators (Finnegan *et al.* 1999; Glazer and Salame 2000). The infective juvenile (IJ) of EPNs are specifically adapted to search for insect hosts, and are able to survive in the absence of a host for weeks, or even months through a reliance on energy reserves (Hatab and Gaugler 1999; Qiu and Bedding 2000; Hass *et al.* 2002). Additionally, in order to conserve energy, the IJs of some EPN species become inactive when stored in water (Fitters and Griffin 2004).

The success of EPNs depends largely on their physiology and biochemistry that enables them to achieve efficient control of pests. In this context, the amount of lipid within the body of the IJ greatly influences the viability and infectivity of the nematode (Wright and Perry 2002). Saturated lipids are the most important energy reserves of aerobic nematodes, particularly in IJ, and represent between 11% and 67% of the dry weight of the organism (Barrett and Wright 1998). Neutral fats typically account for more than 70% of the total lipid content of nematodes, with triglycerides being the major storage form (Barrett and Wright 1998; Chitwood 1998). IJs rely on lipids as their sole energy source while searching for a host (Lee and Atkinson 1977; Van Gundy 1985), and a high lipid content therefore allows survival for extended periods. There is evidence that an extended period of storage and a reduction in total lipid content can lead to a decrease

¹ Ph. D, Instituto Federal de Minas Gerais, Department of Agricultural Sciences, C.P. 05, Bambuí, MG, Brazil. E-mail: vanessa.andalo@ifmg.edu.br. Corresponding author. ² Ph. D, Universidade Federal de Lavras, Department of Entomology, C.P. 3037, 37200-000, Lavras, MG, Brazil. E-mail: alcemoino@ufla.br. ³ Ms. Sc., Universidade Federal de Lavras, Nematology Laboratory, Department of Phytopatology, C.P. 3037, 37200-000, Lavras, MG, Brazil. E-mail: nema@ufla.br. ⁴ Ph. D, Universidade Federal de Lavras, Nematology Laboratory, Department of Phytopatology, C.P. 3037, 37200-000, Lavras, MG, Brazil. E-mail: nema@ufla.br. ⁵ Ph. D, Instituto Nacional da Propriedade Industrial, Rua Mayrink Veiga, 9, CEP 22221-070, Rio de Janeiro, RJ, Brazil. E-mail: luciaamendonca@yahoo.com.br

in the infectivity of IJs (Lewis *et al.* 1995; Hass *et al.* 2002; Fitters and Griffin 2004). Since the levels of storage lipids may be affected by alterations in environmental conditions, it is likely that the supply of different lipids can determine the survival and pathogenicity of IJs (Hatab and Gaugler 1999; Menti *et al.* 2003).

Triglycerides can be analyzed using various chromatographic techniques or by the application of histological methods based on staining with oil red O (Lee 1960; Croll 1972). Chromatographic analysis is time consuming and demands a large number of nematodes, whereas histological assay is straightforward and can be applied to individual specimens. Image analysis has been successfully employed to quantify the degree of oil red O staining, and allows the examination of many samples within a short period (Stamps and Linit 1995). Moreover, results acquired using the staining method has been validated against those obtained by conventional gas chromatography (Patel *et al.* 1997). The method can thus provide specific and conclusive results regarding the utilization of lipid reserves, and was hence selected for application in the present study.

Many of the market difficulties in introducing EPNs are related to low efficacy under unfavorable conditions, application conditions, refrigeration requirements and limited room temperature shelf life (product quality) (Georgis *et al.* 2006). Lipid reserves can affect each of these factors and are therefore an important consideration when developing EPN biological controls. The objective of the present paper was to determine the levels of IJ lipid reserves in different isolates of *Steinernema* and *Heterorhabditis* after storage and at different temperatures. Additionally, the association between lipid reserves and IJ infectivity towards larvae of *Galleria mellonella* L., 1758 (Lepidoptera: Pyralidae) was evaluated.

Materials and Methods

Multiplication of EPNs. Infective juveniles of *Steinernema carpocapsae* (Weiser, 1955) Wouts, Mráček, Gerdin & Bedding, 1982 (obtained from North Carolina, USA, soil sample), *Steinernema riobrave* Cabanillas, Poinar & Raulston, 1994 (obtained from Texas, USA, soil sample), *Heterorhabditis* sp. JPM4 (from Lavras, MG, Brazil), *Heterorhabditis* sp. CCA (from Araras, SP, Brazil) and *Heterorhabditis* sp. PI (from Piauí, Brazil) were maintained in the form of aqueous suspensions (500 IJs/ml) under biochemical oxygen demand at 16°C.

Multiplication of *G. mellonella* was performed according to the method of Dutky *et al.* (1964). Larvae of *G. mellonella* were bred on a modified artificial diet (Parra 1998). Ten larvae were placed into a Petri dish (9cm diameter) lined with filter paper. An aliquot (1ml) of IJ suspension (diluted in order to provide the equivalent of 20±5 IJ/larvae) was transferred to the filter paper, and the dishes were incubated under BOD at 24±1°C for 72h with a 24h scotophase until the larvae died. The dead larvae were transferred to fresh Petri dishes lined with dry filter paper and maintained for four days following Molina and López (2001) methodology. The larvae were subsequently placed on the inner plate of a modified White trap (White 1927), and the trap was incubated at 24±1°C for 3 to 7 days. Trap water, which contained IJs that had emerged from the host, was collected daily and transferred to a 1L measuring cylinder containing 800ml of distilled water. Adult nematodes and insect fat bodies were separated from the IJs by decanting. The concentration of IJs present in the final suspension was quantified using polystyrene plates normally employed for serological tests, applying 0.1ml of IJ suspension in each well, after ten wells it was obtained the amount of nematodes in 1ml suspension. This process was repeated three times and obtained an average.

Influence of temperature and storage on IJ lipid content and infectivity. The concentrations of neutral lipids present in IJs that had been stored at different temperatures and for various periods were determined using a colorimetric method based on staining with oil red O (Storey 1983). Stain solution was prepared by dissolving 0.5g of the dye in 100ml of absolute alcohol under constant agitation for 15min, followed by filtration through Whatman no. 1 filter paper and storage at 5°C in the dark. Suspensions of IJs (3000 IJs/ml; in 40ml) derived from the various isolates were placed in plastic cups (110 ml) containing distilled water (50 ml) and covered with lids, each perforated with a 2cm diameter hole in order to allow adequate aeration. The cups were then incubated at 8, 16, 20, 24 or 28°C, for 15, 30, 60, 90, 120, 150 or 180 days, under constant darkness. Each treatment was replicated four times. After incubation, the IJ suspension was thoroughly mixed, concentrated to 0.5ml by decantation and transferred to a test tube together with 3 ml of stain solution. The resulting mixture was incubated in a water bath at 60°C for 20min and allowed to cool at room temperature in order to permit the sedimentation of the IJs. The supernatant (*ca.* 2.5ml) was discarded, 3ml of water: glycerine (1:1) solution was added to the concentrated IJ suspension, and the mixture was stored at room temperature until required for histological analysis.

Table 1. Percentage of lipids present during storage of IJs from *Heterorhabditis* sp. CCA for 15 to 180 days at different temperatures.

Day	Temperature (°C)				
	8	16	20	24	28
15	87.8 ± 0.51 ABa	88.7 ± 1.27 Aa	83.7 ± 1.58 Ca	84.7 ± 1.35 BCa	73.5 ± 3.63 Da
30	89.8 ± 0.56 Aa	81.6 ± 1.06 Bb	81.7 ± 1.04 Ba	79.9 ± 1.78 Bb	65.8 ± 1.01 Cb
60	81.6 ± 0.40 Ab	76.6 ± 1.16 Bc	68.5 ± 2.34 Cb	61.3 ± 2.45 Dc	52.7 ± 1.56 Ec
90	87.0 ± 1.99 Aa	80.1 ± 1.61 Bbc	70.6 ± 1.93 Cb	28.6 ± 3.58 Dd	21.6 ± 1.67 Ed
120	80.3 ± 1.49 Ab	68.3 ± 1.30 Bd	59.3 ± 1.78 Cc	0.0 ± 0.0 De	0.0 ± 0.0 De
150	68.1 ± 2.54 Ac	58.6 ± 1.65 Be	50.8 ± 0.57 Cd	0.0 ± 0.0 De	0.0 ± 0.0 De
180	55.2 ± 1.55 Ad	51.3 ± 4.9 Bf	47.1 ± 1.67 Cd	0.0 ± 0.0 De	0.0 ± 0.0 De

Means followed by the same small letter within columns and same capital letter within lines are not significantly different by the Tukey test ($p > 0.01$).

Table 2. Percentage of lipids present during storage of IJs from *Steinernema carpocapsae* for 15 to 180 days at different temperatures.

Day	Temperature (°C)				
	8	16	20	24	28
15	85.9 ± 2.67 Aa	82.6 ± 5.56 Aab	82.8 ± 0.46 Aa	68.7 ± 2.04 Ba	70.2 ± 1.46 Ba
30	85.8 ± 3.98 Aab	86.2 ± 2.03 Aa	80.2 ± 1.41 Ba	71.8 ± 1.58 Ca	61.2 ± 3.27 Db
60	78.9 ± 3.00 Acd	68.5 ± 2.37 Bd	65.3 ± 1.62 Bb	49.0 ± 6.62 Cb	37.5 ± 3.26 Dc
90	80.3 ± 1.48 Abc	74.9 ± 2.27 Bc	70.2 ± 1.23 Bb	4.0 ± 0.48 Cc	1.4 ± 2.87 Cd
120	73.5 ± 3.5 Ad	66.7 ± 0.71 Bd	48.2 ± 1.43 Cc	1.7 ± 1.98 Dc	0.2 ± 0.42 Dd
150	63.1 ± 1.85 Be	68.7 ± 2.54 Ad	4.2 ± 7.36 Cd	0.0 ± 0.0 Cc	0.0 ± 0.0 Cd
180	61.3 ± 1.92 Ae	60.5 ± 1.17 Ae	1.2 ± 0.49 Bd	0.0 ± 0.0 Bc	0.0 ± 0.0 Bd

Means followed by the same small letter within columns and same capital letter within lines are not significantly different by the Tukey test ($p > 0.01$).

Table 3. Percentage of lipids present during storage of IJs from *Heterorhabditis* sp. JPM4 for 15 to 180 days at different temperatures.

Day	Temperature (°C)				
	8	16	20	24	28
15	80.9 ± 0.44 ABa	86.6 ± 2.79 Aa	85.5 ± 3.02 ABa	81.4 ± 7.01 ABa	78.4 ± 4.33 Ba
30	80.5 ± 0.94 ABa	84.0 ± 1.75 Aa	80.2 ± 0.43 ABab	76.2 ± 4.37 Ba	73.7 ± 3.11 Ba
60	77.4 ± 1.82 ABa	79.0 ± 1.36 Aab	71.4 ± 7.33 BCc	66.6 ± 3.84 CDb	63.2 ± 3.89 Db
90	61.9 ± 9.01 Ab	65.2 ± 4.37 Ac	69.0 ± 2.50 Ac	30.6 ± 4.48 Bc	5.7 ± 4.44 Cc
120	59.5 ± 2.12 Abc	55.2 ± 2.07 ABd	50.0 ± 1.40 Bd	7.5 ± 7.30 Cd	0.0 ± 0.0 Dc
150	50.8 ± 3.66 Ad	42.4 ± 3.22 Be	40.3 ± 0.84 Be	0.0 ± 0.0 Cd	0.0 ± 0.0 Cc
180	52.5 ± 2.61 Acd	40.1 ± 1.71 Be	21.2 ± 2.68 Cf	0.0 ± 0.0 Dd	0.0 ± 0.0 Dc

Means followed by the same small letter within columns and same capital letter within lines are not significantly different by the Tukey test ($p > 0.01$).

For each treatment repetition, three stained nematodes, taken randomly from the stored suspension, were placed onto a glass slide and observed and photographed using a trinocular microscope Olympus SZ40. The total IJ body area and the areas corresponding to the red-stained lipids were estimated from the digitized images with the aid of Image Tool for Windows version 3.0 (The University of Texas Health Science Center, San Antonio, Texas), from which it was possible to determine the percentage of triglycerides in relation to body area.

The effects of storage on the infectivity of IJs towards larvae of *G. mellonella* were evaluated by placing five larvae into a Petri dish (5cm diameter) lined with filter paper and adding a 0.5ml aliquot of IJ suspension that had been treated as described above. Dishes were incubated at $24 \pm 1^\circ\text{C}$ for 3 days, after which the mortality rates of the larvae were

determined. Each treatment was replicated four times. *Galleria mellonella* larvae were tested as control applying water, without IJ.

The percentage data corresponding to lipid concentration and infectivity of the stored IJs were submitted to variance analysis and to the Tukey test ($p > 0.01$) for comparisons between the means.

Results

Concentration of lipids in IJs after storage at different temperatures. The concentrations of neutral lipids, expressed as percentages of the total body areas of the IJs, varied between the species tested. Maximum values were detected in newly emerged IJs. The average of neutral lipids for newly emerged IJ just before storage were 97.4% for *S.*

Table 4. Percentage of lipids present during storage of IJs from *Heterorhabditis* sp. PI for 15 to 180 days at different temperatures.

Day	Temperature (°C)				
	8	16	20	24	28
15	81.2 ± 1.3 Aa	85.7 ± 2.9 Aa	81.3 ± 1.88 Aa	78.6 ± 2.87 Aa	68.1 ± 2.63 Ba
30	79.8 ± 2.08 ABa	81.9 ± 2.93 Aab	77.6 ± 2.54 ABa	73.9 ± 3.74 BCa	68.5 ± 2.06 Ca
60	76.5 ± 2.35 Aa	77.5 ± 1.87 Abc	74.5 ± 3.63 Aab	59.6 ± 1.92 Bb	54.0 ± 2.17 Bb
90	75.7 ± 7.76 Aab	70.4 ± 5.83 Ac	68.8 ± 6.47 Ab	36.3 ± 9.98 Bc	34.3 ± 5.12 Bc
120	68.0 ± 4.54 Abc	44.9 ± 2.9 Bd	39.4 ± 1.46 Bc	4.6 ± 9.27 Cd	0.0 ± 0.0 Cd
150	60.7 ± 3.84 Ac	42.5 ± 2.51 Bde	38.0 ± 1.94 Bc	0.0 ± 0.0 Cd	0.0 ± 0.0 Cd
180	52.0 ± 4.5 Ad	35.0 ± 2.76 Be	13.5 ± 2.66 Cd	0.0 ± 0.0 Dd	0.0 ± 0.0 Dd

Means followed by the same small letter within columns and same capital letter within lines are not significantly different by the Tukey test ($p > 0.01$).

Table 5. Percentage of lipids present during storage of IJs from *Steinernema riobrave* for 15 to 180 days at different temperatures.

Day	Temperature (°C)				
	8	16	20	24	28
15	84.7 ± 1.27 Aa	89.6 ± 2.83 Aa	88.1 ± 3.14 Aa	85.3 ± 4.60 Aa	71.3 ± 8.20 Ba
30	83.4 ± 1.36 Aa	87.6 ± 2.53 Aab	86.3 ± 2.52 Aa	73.4 ± 4.05 Bb	71.4 ± 2.07 Ba
60	83.0 ± 3.90 Aa	81.5 ± 2.31 Ab	78.8 ± 1.32 Ab	69.0 ± 1.19 Bb	58.5 ± 1.61 Cb
90	82.5 ± 0.96 Aa	81.9 ± 0.93 Ab	81.7 ± 2.19 Aab	57.6 ± 2.65 Bc	32.9 ± 2.87 Cc
120	80.5 ± 2.71 Aab	61.5 ± 1.31 Bc	43.1 ± 10.6 Cc	29.3 ± 3.54 Dd	26.8 ± 1.91 Dc
150	78.4 ± 2.96 Aab	53.6 ± 2.38 Bd	35.2 ± 3.06 Cd	8.7 ± 1.79 De	0.0 ± 0.0 Ed
180	75.0 ± 1.41 Ab	38.4 ± 4.47 Be	32.3 ± 3.25 Cd	9.8 ± 0.52 De	0.0 ± 0.0 Ed

Means followed by the same small letter within columns and same capital letter within lines are not significantly different by the Tukey test ($p > 0.01$).

riobrave, 90.3% for *S. carpocapsae*, 89.2% for *Heterorhabditis* sp. PI, 88.6% for *Heterorhabditis* sp. CCA and 90.7% for *Heterorhabditis* sp. JPM4.

The intensity of the lipid stain was stronger in the area of the intestine and lighter in the anterior portion, with irregular spots distributed throughout the body (Fig. 1). For all of the nematodes studied, the percentage of lipids generally diminished with time, although there were variations in the rate of decrease depending on the temperature of storage on the EPN. At temperatures between 8 and 20°C, the IJs conserved lipid reserves longer, whereas between 24 and 28°C, the percentage of lipids diminished rapidly. After 180 days at 24 and 28°C the reserves of lipids were reduced to < 50% in 90 days, and were almost completely depleted after 120 days (Tables 1-5). Between 8 and 16°C, the IJs were motionless, and for *S. carpocapsae* it was possible to observe that the typical “J” form. In contrast, between 20 and 28°C, the IJs were in constant motion. Hence, the maintenance of lipid reserves was more efficient when IJs were stored at 8 or 16°C, independent of species or strain, even when the IJs were dead.

Infectivity of IJs after storage at different temperatures.

For all of the nematodes studied, the IJs stored for 0 days showed the highest mortality rates towards larvae of *G. mellonella*. For *Heterorhabditis* sp. PI, *S. carpocapsae* and *S. riobrave* these mortality rates were 100%, for *Heterorhabditis* sp. CCA was 95% and for *Heterorhabditis* sp. JPM4

was 97.5%. The percentage infectivity of IJs diminished upon storage, but the rate of reduction varied according to storage temperature and EPN. At temperatures of 24 and 28°C, the ability of all nematodes to infect insect larvae typically diminished after just 15 days storage. On the other hand, the infectivities of IJs from *S. carpocapsae* and *S. riobrave* were unaffected by storage at 8°C for up to 90 days, although those of IJs from the *Heterorhabditis* species were greatly reduced after 15 days of incubation under such conditions. Indeed, IJs from *Heterorhabditis* sp. CCA were no longer able to infect *G. mellonella* larvae after 60 days of storage at 8°C. Thus, the IJs from *Heterorhabditis* isolates were less tolerant to both the lowest (8°C) and the highest (24 and 28°C) temperatures tested in comparison with those from *Steinernema* species.

The most favorable temperature for the maintenance of EPN infectivity was estimated to be between 16 and 20°C, since the percentage infectivities after an incubation period of 180 days were 72.5% for *S. carpocapsae*, 70.0% for *S. riobrave* and 62.5% for *Heterorhabditis* sp. CCA with storage at 16°C, and 40% for *Heterorhabditis* sp. PI, 60.0% for *Heterorhabditis* sp. CCA and 67.5% for *Heterorhabditis* sp. JPM4 with storage at 20°C (Tables 6-10).

Correlation between lipid content and infectivity. In the case of IJs from the two *Steinernema* species, a direct correlation could be established at all storage temperatures between




Figure 1. Variations in the concentration of lipids present in infective juveniles of *Heterorhabditis* sp. JPM4 I and *Steinernema riobrave* II, as determined by oil red O staining, after storage at A. 15 days at 8°C; B. 180 days at 8°C; C. 15 days at 28°C; and D. 180 days at 28°C.

Table 6. Percentage infectivity to larvae of *Galleria mellonella* during storage of IJs from *Heterorhabditis* sp. CCA for 15 to 180 days at different temperatures.

Day	Temperature (°C)				
	8	16	20	24	28
15	92.5 ± 5.0 Aa	97.5 ± 5.0 Aab	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa
30	0.0 ± 0.0 Bb	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa
60	0.0 ± 0.0 Bb	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa
90	0.0 ± 0.0 Cb	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	70.0 ± 8.16 Bb	0.0 ± 0.0 Cb
120	0.0 ± 0.0 Bb	87.5 ± 12.5 Ab	82.5 ± 12.5 Ab	0.0 ± 0.0 Bc	0.0 ± 0.0 Bb
150	0.0 ± 0.0 Bb	67.5 ± 9.57 Ac	70.0 ± 11.5 Ac	0.0 ± 0.0 Bc	0.0 ± 0.0 Bb
180	0.0 ± 0.0 Bb	62.5 ± 9.57 Ac	60.0 ± 8.16 Ac	0.0 ± 0.0 Bc	0.0 ± 0.0 Bb

Means followed by the same small letter within columns and same capital letter within lines are not significantly different by the Tukey test ($p > 0.01$).

lipid content and infectivity, i.e. as the percentage of lipids in the nematode body diminished, the infectivity to *G. mellonella* larvae tended to decrease. With respect to the *Heterorhabditis* isolates, however, there was an inverse correlation at 8°C, thus, despite a high lipid content infectivity was lost after 60 days of storage.

After 180 days of incubation at 16°C all IJs, except for those derived from *Heterorhabditis* sp. PI, induced > 50% mortality in *G. mellonella* larvae. For *S. carpocapsae* and *S. riobrave*, the percentage infectivities were < 50% when the lipid reserves fell below 20% and 32%, respectively. After incubation at 8°C, all *Heterorhabditis* isolates presented low infectivities even when the lipid content was high. Thus, IJs from *Heterorhabditis* spp. PI, CCA and JPM4 that had been stored for 180 days at 8°C could not induce mortality in *G. mellonella* larvae although their lipid contents were 70%, 80% and 77%, respectively. At storage temperatures > 8°C, infectivities of around 50% were attained by IJs from *Heterorhabditis* spp. PI, CCA and JPM4 that exhibited lipid contents of < 46%, 15% and 15%, respectively. Hence, although the lipid reserves of *Heterorhabditis* nematodes were adequately preserved at 8°C, the infectivities of the IJs were not.

Discussion

According to Grewal (2000), *S. carpocapsae* is able to survive longer in water at 5°C than other nematodes because it adopts a “J” form, thus becoming quiescent and conserving energy. The differential thermo-adaptation of some nematodes is associated with the accumulation of trehalose lipid,

a non-reducing disaccharide that plays a role in the maintenance of virulence during acclimatization to low (5°C) or high (35°C) temperatures (Jagdale and Grewal 2003). The amount of trehalose accumulated by different nematodes varies with temperature, and this may confer specific advantages to certain species, as revealed in the present study for *Steinernema* sp. in which lipid reserves were preserved for longer periods at all temperatures tested.

Some IJs become inactive in water in order to conserve energy, and this ability also varies with species. Several studies have shown that the percentage of inactive IJs in water is greater for *S. carpocapsae* than for *S. glaseri* or *H. bacteriophora* (Lewis *et al.* 1995; Fitters and Griffin 2004). Additionally, a significant increase in infectivity towards *G. mellonella* larvae was reported for *H. megidis* IJs in the first two weeks of storage in water, after which infectivity declined (Fitters and Griffin 2004).

Molecular studies of genetic modulation have revealed that IJ stress response is associated with proteins involved in desiccation that are differentially regulated by more than 30 genes, thus determining a differential capacity for adaptation to extreme environments such as excessive salt concentrations, excessive temperature oscillations and low humidity. The response to stress selects for IJ populations that are genetically adapted to survive under severe conditions (Serwer-Rodriguez *et al.* 2004; Bornstein-Forst *et al.* 2005), as is the case for some *Steinernema* species that are well adapted to resist large oscillations of temperature.

The levels lipids in *Heterorhabditis* isolates varied less than in *Steinernema* isolates, and some of the lipids found in

Table 7. Percentage infectivity to larvae of *Galleria mellonella* during storage of IJs from *Steinernema carpocapsae* for 15 to 180 days at different temperatures.

Day	Temperature (°C)				
	8	16	20	24	28
15	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa
30	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa
60	95.0 ± 5.77 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa
90	90.0 ± 8.16 ABab	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	82.5 ± 5.0 Bb	27.5 ± 9.57 Cb
120	80.0 ± 8.16 Bb	95.0 ± 5.77 Aab	90.0 ± 8.16 ABa	0.0 ± 0.0 Cc	0.0 ± 0.0 Cc
150	65.0 ± 12.9 Bc	82.5 ± 17.0 Abc	50.0 ± 8.16 Cb	0.0 ± 0.0 Dc	0.0 ± 0.0 Dc
180	52.5 ± 17.0 Bc	72.5 ± 17.0 Ac	47.5 ± 5.0 Bb	0.0 ± 0.0 Cc	0.0 ± 0.0 Cc

Means followed by the same small letter within columns and same capital letter within lines are not significantly different by the Tukey test ($p > 0.01$).

Table 8. Percentage infectivity to larvae of *Galleria mellonella* during storage of IJs from *Heterorhabditis* sp. JPM4 for 15 to 180 days at different temperatures.

Day	Temperature (°C)				
	8	16	20	24	28
15	77.0 ± 8.71 Ba	95.0 ± 10.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa
30	7.5 ± 9.57 Bb	100.0 ± 0.0 Aa	95.0 ± 10.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa
60	0.0 ± 0.0 Bb	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa
90	0.0 ± 0.0 Cb	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	87.5 ± 9.57 Bb	0.0 ± 0.0 Cb
120	0.0 ± 0.0 Bb	65.0 ± 5.77 Ab	70.0 ± 8.16 Ab	0.0 ± 0.0 Bc	0.0 ± 0.0 Bb
150	0.0 ± 0.0 Bb	60.0 ± 8.16 Ab	67.5 ± 15.0 Ab	0.0 ± 0.0 Bc	0.0 ± 0.0 Bb
180	0.0 ± 0.0 Cb	55.0 ± 12.9 Bb	67.5 ± 9.57 Ab	0.0 ± 0.0 Cc	0.0 ± 0.0 Cb

Means followed by the same small letter within columns and same capital letter within lines are not significantly different by the Tukey test ($p > 0.01$).

the latter were typically either not present or present only in small amounts in the former. Additionally, *Steinernema* nematodes survived longer than *Heterorhabditis* species from which the lipids were depleted more rapidly. In all species of *Steinernema*, a reduction in lipid reserves was observed from the fourth week onwards with storage at 23°C, and from the fifth week onwards with storage at 20°C (Lewis *et al.* 1995; Patel *et al.* 1997; Fitters *et al.* 1999). The levels of lipids reserves observed in the newly-emerged IJ populations were similar (i.e. between 80% and 96%) to those reported for *S. feltiae* and *H. megidis* in which IJs of the former survived for 16 to 20 weeks whilst those of the latter survived for only 12 weeks and with declining viability (Patel and Wright 1997). The present study thus corroborates these previous reports since the percentage of lipids in *Heterorhabditis* isolates stored at 24°C diminished from the 30th day onwards at a rate that was much higher than that observed for *Steinernema* species. It has been reported that *S. carpocapsae* can regulate the water content of its membranes depending on the temperature (Fodor *et al.* 1994), whilst *S. riobrave* accumulates large proportions of unsaturated fatty acids at high temperatures (Hatab and Gaugler 1999), characteristics that contribute to the thermo-tolerance of these species.

According to Patel *et al.* (1997), the survival of *S. carpocapsae* and *S. riobrave* varies between 120 and 135 days, the infectivities of *S. riobrave*, *S. feltiae* and *S. glaseri*, decline as the lipid reserves are depleted. Moreover, the sizes of the IJs and the initial amounts of lipid present in their bodies appear to be related to survival and the manner in which the organisms use energy reserves. Thus smaller nematodes, like those

S. carpocapsae and *S. riobrave*, utilize their lipids more rapidly than larger IJs like those of *S. feltiae* and *S. glaseri* (Patel *et al.* 1997). In the present study, however, the infectivity of *S. carpocapsae* IJs was maintained even when lipid reserves were low, suggesting that nematode populations are quite heterogeneous and are able to use such reserves for different purposes. Furthermore, the present findings demonstrated the importance of locomotive activity of IJs during storage in water and the significance of the quiescent form for minimizing the use of energy.

Steinernema carpocapsae IJs exhibited a rapid decline in infectivity after 90 days of storage, which is similar to the findings of Wright *et al.* (1997). These authors emphasized that the use of glycogen as energy reserve ceases earlier on whilst the lipid reserves continue for a much longer period, perhaps explaining the initial rapid decrease in the infectivity of *S. carpocapsae*. In *Heterorhabditis* IJs, a direct relationship between infectivity and the percentage of lipids was also observed, as previously reported by Hass *et al.* (2002).

As stated by Wright and Perry (2002) the efficacy of the selection of EPN isolates that are already adapted for commercial use depends on a comprehensive knowledge of their physiological and biochemical attributes. Clearly the quality and quantity of lipids present in the IJs are of utmost importance, since such energy reserves critically influence the viability and infectivity of EPNs.

Conclusions

Thus, this paper presents the importance of lipid reserves for IJs, showing that temperature directly influences its variation

Table 9. Percentage infectivity to larvae of *Galleria mellonella* during storage of IJs from *Heterorhabditis* sp. PI for 15 to 180 days at different temperatures.

Day	Temperature (°C)				
	8	16	20	24	28
15	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	87.5 ± 15.0 Ba
30	70.0 ± 11.5 Bb	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	90.0 ± 14.1 Aa
60	0.0 ± 0.0 Cc	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	92.5 ± 9.57 ABa	85.0 ± 12.6 Bab
90	0.0 ± 0.0 Bc	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	87.5 ± 12.6 Aa	0.0 ± 0.0 Bc
120	0.0 ± 0.0 Bc	67.5 ± 9.57 Ab	70.0 ± 8.16 Ab	0.0 ± 0.0 Bb	0.0 ± 0.0 Bc
150	0.0 ± 0.0 Cc	47.5 ± 5.0 Bc	62.5 ± 9.57 Ab	0.0 ± 0.0 Cb	0.0 ± 0.0 Cc
180	0.0 ± 0.0 Bc	30.0 ± 8.16 Ad	39.2 ± 12.9 Ac	0.0 ± 0.0 Bb	0.0 ± 0.0 Bc

Means followed by the same small letter within columns and same capital letter within lines are not significantly different by the Tukey test ($p > 0.01$).

Table 10. Percentage infectivity to larvae of *Galleria mellonella* during storage of IJs from *Steinernema riobrave* for 15 to 180 days at different temperatures.

Day	Temperature (°C)				
	8	16	20	24	28
15	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa
30	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	97.5 ± 5.0 Aab	95.0 ± 10.0 Aa
60	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	92.5 ± 9.57 Aab	90.0 ± 14.1 Aa
90	97.5 ± 5.0 ABa	100.0 ± 0.0 Aa	100.0 ± 0.0 Aa	82.5 ± 9.57 Bb	40.0 ± 8.16 Cb
120	87.5 ± 9.57 Aa	90.0 ± 8.16 Aab	80.0 ± 8.16 Ab	60.0 ± 8.16 Bc	47.5 ± 15.0 Bb
150	67.5 ± 12.5 Ab	80.0 ± 14.1 Abc	70.0 ± 18.2 Abc	32.5 ± 12.5 Bd	0.0 ± 0.0 Cc
180	62.5 ± 12.5 Ab	70.0 ± 14.1 Ac	62.5 ± 9.57 Ac	0.0 ± 0.0 Be	0.0 ± 0.0 Bc

Means followed by the same small letter within columns and same capital letter within lines are not significantly different by the Tukey test ($p > 0.01$).

and correlates the decline of the reserve to a decrease in infectivity for all species tested. By using this information it is possible to handle properly the situations for greater success in using EPNs.

Cited Literature

- BARRETT, J.; WRIGHT, D. J. 1998. Intermediary metabolism. p. 331-353. En: Perry, R. N.; Wright, D. J. (eds.). The physiology and biochemistry of free-living and plant-parasitic nematodes. CAB International. Wallingford. 448 p.
- BORNSTEIN-FORST, S.; KIGER, H.; RECTOR, A. 2005. Impacts of fluctuating temperature on the development and infectivity of entomopathogenic nematode *Steinernema carpocapsae* A10. *Journal of Invertebrate Pathology* 88 (2): 147-153.
- CHITWOOD, D. J. 1998. Biosynthesis. p. 303-330. En: Perry, R. N.; Wright, D. J. (eds.). The physiology and biochemistry of free-living and plant-parasitic nematodes. CAB International. Wallingford. 448 p.
- CROLL, N. A. 1972. Energy utilization of infective *Ancylostoma tubaeforme* larvae. *Parasitology* 64 (3): 355-368.
- DUTKY, S. R.; THOMPSON, J. V.; CANTWE, G. E. 1964. A technique for the mass propagation of the DD-136 nematode. *Journal of Insect Pathology* 6 (4): 417-422.
- FINNEGAN, M.; DOWNES, M. J.; O'REGAN, M.; GRIFFIN, C. T. 1999. Effect of salt and temperature stresses on survival and infectivity of *Heterorhabditis* spp. infective juveniles. *Nematology* 1 (1): 69-78.
- FITTERS, P. F. L.; GRIFFIN, C. T. 2004. Spontaneous and induced activity of *Heterorhabditis megidis* infective juveniles during storage. *Nematology* 6 (6): 911-917.
- FITTERS, P.; PATEL, M. N.; GRIFFIN, C. T.; WRIGHT, D. J. 1999. Fatty acid composition of *Heterorhabditis* sp. during storage. *Comparative Biochemistry and Physiology* 124 (1): 81-88.
- FODOR, A.; DEY, I.; FARKAS, T.; CHITWOOD, D. J. 1994. Effects of temperature and dietary lipids on phospholipid fatty acids and membrane fluidity in *Steinernema carpocapsae*. *Journal of Nematology* 26 (3): 278-285.
- GEORGIS, R.; KOPPENHÖFER, A. M.; LACEY, L. A.; BÉLAIR, G.; DUNCAN, L. W.; GREWAL, P. S.; SAMISH, M.; TAN, L.; TORR, P.; VAN TOL, R. W. H. M. 2006. Successes and failures in the use of parasitic nematodes for pest control. *Biological Control* 38 (1): 103-123.
- GLAZER, I.; SALAME, L. 2000. Osmotic survival of the entomopathogenic nematode *Steinernema carpocapsae*. *Biological Control* 18 (3): 251-257.
- GREWAL, P. S. 2000. Anhydrobiotic potential and long-term storage of entomopathogenic nematodes (Rhabditida: Steinernematidae). *International Journal for Parasitology* 30 (14): 995-1000.
- HASS, B.; DOWNES, M. J.; GRIFFIN, C. T. 2002. Persistence of four *Heterorhabditis* spp. isolates in soil: role of lipid reserves. *Journal of Nematology* 34 (2): 151-158.
- HATAB, M. A.; GAUGLER, R. 1999. Lipids of in vitro cultured *Heterorhabditis bacteriophora*. *Biological Control* 15 (3): 113-118.
- JAGDALE, G. B.; GREWAL, P. S. 2003. Acclimation of entomopathogenic nematodes to novel temperatures: trehalose accumulation and the acquisition of thermotolerance. *International Journal for Parasitology* 33 (2): 145-152.
- LEE, D. L. 1960. The distribution of glycogen and fat in *Thelastoma bulhõesi* (Magalhães, 1900), a nematode parasitic in cockroaches. *Parasitology* 50: 247-259.
- LEE, D. L.; ATKINSON, H. J. 1977. Physiology of nematodes. Columbia University, New York. 215 p.
- LEWIS, E. E.; SELVAN, S.; CAMPBELL, J. F.; GAUGLER, R. 1995. Changes in foraging behaviour during the infective stages of entomopathogenic nematodes. *Parasitology* 110: 583-590.
- MENTI, H.; PATEL, M. N.; WRIGHT, D. J.; PERRY, R. N. 2003. Lipid utilisation during storage of the entomopathogenic nematodes *Steinernema feltiae* and *Heterorhabditis megidis* from Greece and the UK. *Nematology* 5 (1): 31-37.
- MOLINA, J. P.; LÓPEZ, N. J. C. 2001. Producción in vivo de tres entomopatodados con dos sistemas de infección en dos hospedantes. *Revista Colombiana de Entomología* 27 (1-2): 73-78.
- PARRA, J. R. P. 1998. Criação de insetos para estudos com patógenos. p. 1015-1037. En: Alves, S. B. (ed.). Controle microbiano de insetos. FEALQ. Piracicaba. 1163 p.
- PATEL, M. N.; WRIGHT, D. J. 1997. Phospholipid fatty acid composition in steinernematid entomopathogenic nematodes. *Comparative Biochemistry and Physiology* 118 (3): 649-657.
- PATEL, M. N.; STOLINSKI, M.; WRIGHT, D. J. 1997. Neutral lipids and the assessment of infectivity in entomopathogenic nematodes: observations on four *Steinernema* species. *Parasitology* 114: 489-496.
- QIU, L.; BEDDING, R. 2000. Energy metabolism and its relation to survival and infectivity of infective juveniles of *Steinernema carpocapsae* under aerobic conditions. *Nematology* 2: 551-559.
- SERWE-RODRIGUEZ, J.; SONNENBERG, K.; APPLEMAN, B.; BORNSTEIN-FORST, S. 2004. Effects of in host desiccation on development, survival, and infectivity of entomopathogenic nematode *Steinernema carpocapsae*. *Journal of Invertebrate Pathology* 85 (3): 175-181.
- STAMPS, T. W.; LINIT, M. J. 1995. A rapid and simple method for staining lipid in fixed nematodes. *Journal of Nematology* 27 (2): 244-247.
- STOREY, R. M. J. 1983. The initial neutral lipid reserves of juveniles of *Globodera* spp. *Nematologica* 29 (2): 144-150.
- VAN GUNDY, S. D. 1985. Ecology of *Meloidogyne* spp.- emphases on environmental factors affecting survival and pathogenicity. p. 177-182. En: Sasser, J. N.; Carter, C. C. (eds.). An advanced treatise on *Meloidogyne*. North Carolina State University Graphics. Raleigh. 111 p.
- WHITE, G. F. 1927. A method for obtaining infective nematode larvae from cultures. *Science* 66 (1709): 302-303.
- WRIGHT, D. J.; PERRY, R. N. 2002. Physiology and biochemistry. p. 145-168. En: Gaugler R. (ed.). Entomopathogenic nematology. CAB International. Wallingford. 388 p.
- WRIGHT, D. J.; GREWAL, P. S.; STOLINSKI, M. 1997. Relative importance of neutral lipids and glycogen as energy stores in dauer larvae of two entomopathogenic nematodes, *Steinernema carpocapsae* and *Steinernema feltiae*. *Comparative Biochemistry and Physiology* 118 (2): 269-273.