

Artículo de Revisión

Phylogenetic studies and modern classification of the Pyraloidea (Lepidoptera)

Estudios filogenéticos y clasificación actual de los Pyraloidea (Lepidoptera)

M. ALMA SOLIS¹

Abstract. Pyraloidea, the third largest superfamily of the Lepidoptera, is comprised of two families - Pyralidae and Crambidae. The history of families previously placed in the Pyraloidea is discussed. The group now includes about 16,000 species worldwide. Morphologically, the superfamily is defined by a basally scaled proboscis and the presence of abdominal tympanal organs. The larvae of many species are economically important pests of crops (e. g.: sugarcane, corn, rice), and stored products such as seeds and grains. Currently 22 subfamilies comprise the Pyraloidea; only the 19 subfamilies that occur in the Western Hemisphere are discussed. There is a paucity of recent research using cladistic methods and phylogenetic analyses across all taxa.

Key words. Pyralidae. Crambidae. Morphology. Cladistic analyses. Searchable on-line catalogs.

Resumen. Pyraloidea, la tercera superfamilia más grande de Lepidoptera, está compuesta por dos familias Pyralidae y Crambidae. Se discute la historia de las familias previamente ubicadas en Pyraloidea. Actualmente el grupo incluye cerca de 16.000 especies en el mundo. Morfológicamente, la superfamilia se define por una proboscis con escamas basales y por la presencia de órganos timpánicos abdominales. Las larvas de muchas especies son plagas de cultivos económicamente importantes (p. ej.: caña de azúcar, maíz, arroz) y de productos almacenados tales como granos y semillas. Actualmente los Pyraloidea comprenden 22 subfamilias, solo se discuten las 19 subfamilias que se encuentran en el hemisferio occidental. Hay pocas investigaciones recientes que usen métodos cladísticos y análisis filogenéticos en todos los taxones.

Palabras clave. Pyralidae. Crambidae. Morfología. Análisis cladístico. Consulta de catálogos en línea.

Pyraloidea, the third largest superfamily of the Lepidoptera following Noctuoidea and Geometroidea, are comprised of two families - Pyralidae and Crambidae. The group includes about 16,000 species worldwide, with greatest richness in the tropics (Fig. 1). Morphologically, the superfamily is defined by a basally scaled proboscis and the presence of abdominal tympanal organs. The larvae of many species are economically important pests of crops (e. g.: sugarcane, corn, rice), and stored products such as seeds and grains (Solis 1997).

The familial composition of the Pyraloidea (Table 1) has changed markedly over the past 30 years, with a general decrease in the number of families. Pterophoridae, Alucitidae, Thyrididae, Hyblaeidae, Oxychirotidae and Tineodidae formerly included in the Pyraloidea, are now in their own superfamilies with unresolved affinities (Fletcher and Nye 1984; Minet 1986; Nielsen 1989; Common 1990). Dudgeoneidae is now assigned to Cossidae, and Lathrotelidae, proposed by Clarke (1971) for a monotypic genus from the Pacific islands, is now considered to belong to the Acentropinae in the Crambidae (Minet 1991).

Pyraloidea moths are ditrysian moths characterized by the following morphological features: paired tympanal chambers on sternite 2, each with a tympanum and a conjunctiva (Fig.

2), a maxillary palpus that is usually present, a basally scaled proboscis, veins R_3 and R_4 of the forewing stalked or fused, and $Sc+R_1$ and R_s of the hindwing anastomosed distad of the discal cell.

Although published catalogs on subfamilies are listed below, there are two searchable on-line catalogs of the Pyraloidea. The LepIndex, at The Natural History Museum, London, United Kingdom, is based on their card catalog (Beccaloni *et al.* 2003). The other, GLOBIZ, at the State Museum of Zoology, Dresden, Germany, is compiled and updated by current researchers in the Pyraloidea (Nuss 2006).

Pyraloidea currently is divided into two families based primarily on two distinct tympanal organ types of the adult abdomen, but characters of the larvae also confirm this division. The Pyralidae have a tympanal case that is almost closed, the conjunctiva and tympanum are in the same plane, and the praecinctorium is absent. The Crambidae have a tympanal case that is open with a wide anteromedial aperture, the conjunctiva and tympanum are in a different plane and meet at a distinct angle, and the praecinctorium is present (Minet 1981; Maes 1995). Börner (1925) was the first to recognize the difference between the two groups in the Pyraloidea, and Munroe (1972, 1973, 1976) proposed the informal groups, Pyraliformes and Crambiformes based on the major differences between the two types of tympanal organs. Minet (1983) subsequently elevated Munroe's groups to the Pyralidae and Crambidae based on an extensive study of tympanal organs in Lepidoptera.

Although some contemporary lepidopterists do not recognize the two families as distinct because they "cannot be separated by external diagnostic characters" (Landry 1995), characters of the tympanal organs and larvae are certainly external. From a cladistic perspective, most of the structures of the crambid type of tympanal organ represent synapomorphies. The most

Table 1. Higher-level Pyraloidea taxa.

Pyraloidea, 1980	Other superfamilies	Pyraloidea Today
Pyralidae		Pyralidae + Crambidae
Pterophoridae	Pterophoroidea	
Thyrididae	Thyridoidea	
Hyblaeidae	Hyblaeoidea	
Alucitidae	Alucitoidea	
Tineodidae	Alucitoidea	
Dudgeoneidae	Cossoidea	
Lathrotelidae		<i>Lathroteles</i> , crambid genus

¹ Systematic Entomology Laboratory, USDA, Smithsonian Institution, P.O. Box 37012, National Museum Natural History, E-517, MRC 168, Washington, D.C. 20013-7012. alma.solis@ars.usda.gov



Figure 1. Crambidae: Acentropinae: **A.** *Parapoynx obscuralis* (Grote, 1881). **B.** *Petrophila bifascialis* (Robinson, 1869). **C.** *Synclita oblitalis* (Walker, 1859). Crambidae: Crambinae: **D.** *Crambus praefectellus* (Zincken, 1821). **E.** *Fissicrambus mutabilis* (Clemens, 1860). **F.** *Microcrambus elegans* (Clemens, 1860). **G.** *Urola nivalis* (Drury, 1773). Crambidae: Evergestinae: **H.** *Evergestis rimosalis* (Guenée, 1854). Crambidae: Glaphyriinae: **I.** *Chalcoela pegasalis* (Walker, [1866]). **J.** *Eustixia pupula* (Hübner, 1823). **K.** *Glaphyria glaphyralis* (Guenée, 1854). Crambidae: Pyraustinae: **L.** *Achyra rantalis* (Guenée, 1854). **M.** *Hahncappsia marculenta* (Grote & Robinson, 1867). **N.** *Ostrinia nubilalis* (Hübner, 1796). **O.** *Saucrobotys futilalis futilalis* (Lederer, 1863). Crambidae: Schoenobiinae: **P.** *Donacaula aquilella* (Clemens, 1860). Crambidae: Scopariinae: **Q.** *Eudonia heterosalis* (McDunnough, 1961). **R.** *Scoparia biplagiata* Walker, 1866. Crambidae: Spilomelinae: **S.** *Apogeshna stenialis* (Guenée, 1854). **T.** *Choristostigma roseopennalis* (Hulst, 1886). **U.** *Conchylodes ovulalis* (Guenée, 1854). **V.** *Desmia funeralis* (Hübner, 1796). **W.** *Hymenia perspectalis* (Hübner, 1796). **X.** *Nomophila nearctica* Munroe, 1973. **Y.** *Pantographa limata* (Grote & Robinson, 1867). **Z.** *Pleuroptya silicalis* (Guenée, 1854). **AA.** *Udea rubigalis* (Guenée, 1854). Pyralidae: Chrysauginae: **BB.** *Tosale oviplagalis* (Walker, 1866). Pyralidae: Epipaschiinae: **CC.** *Epipaschia superatalis* Clemens, 1860. Pyralidae: Galleriinae: **DD.** *Omphalocera cariosa* Lederer, 1863. Pyralidae: Phycitinae: **EE.** *Ephestia kuehniella* Zeller, 1879. **FF.** *Laetilia coccidivora* (Comstock, 1879). **GG.** *Peoria roseointictella* (Ragonot, 1887). Pyralidae: Pyralinae: **HH.** *Aglossa caprealis* (Hübner [1809]). **II.** *Herculia olinalis* (Guenée, 1854).

Table 2. Classification of Crambidae subfamilies (synonyms in italics).

Minet (1981[1982])	Munroe and Solis (1999)	Solis and Maes (2002)
Cathariinae*	Cathariinae*	Cathariinae*
Crambinae	Crambinae	Crambinae
Cybalomiinae	Cybalomiinae	Cybalomiinae
Evergestinae	Evergestinae	Evergestinae
Glaphyriinae	Glaphyriinae	Glaphyriinae
Dichogaminae	<i>Alatuncusiinae</i> <i>Dichogaminae</i>	<i>Alatuncusiinae</i> Dichogaminae
Linostinae	Linostinae	Linostinae
Midilinae	Midilinae	Midilinae
Musotiminae	Musotiminae	Musotiminae
Noordinae*	Noordinae*	Noordinae*
Nymphulinae	Nymphulinae	Acentropinae** <i>Nymphulinae</i>
Odontiinae	Odontiinae	Odontiinae
Pyraustinae	Pyraustinae	Pyraustinae
Spilomelinae	<i>Spilomelinae</i>	Spilomelinae
Schoenobiinae	Schoenobiinae	Schoenobiinae
Scopariinae	Scopariinae	Scopariinae
Heliiothelinae	<i>Heliiothelinae</i> Wurthiinae*	<i>Heliiothelinae</i> Wurthiinae*

*does not occur in the Western Hemisphere.

**International Commission for Zoological Nomenclature sustained Acentropinae as the senior valid name (Solis 1999b).

convincing synapomorphy for Pyralidae is the sclerotized ring around the base of seta SD1 on segment A8 of the larva (Fig. 3). In a few genera in both Crambidae and Pyralidae, tympanal organs are absent or reduced (e. g.: *Lathroteles* (Minet 1991); *Michaelshaffera* (Solis 1998)), and a few pyralid species lack the sclerotized ring on A8, although a shiny white translucent ring is present if it is not sclerotized (*Etiella zinckenella* (Solis 1999a); laboratory reared *Galleria mellonella*, personal observation). A molecular study testing the utility of the *period*




Figure 2. A. Pyralidae, praecinctorium absent, tympanum and conjunctiva in the same plane (line horizontal only). B. Pyralidae, praecinctorium absent, tympanal case closed medially and with a small, anterior aperture (bold dotted lines). C. Crambidae, praecinctorium present, tympanum and conjunctiva meeting at a distinct angle (lines horizontal and vertical). D. Crambidae, praecinctorium present, tympanal case open anteromedially (bold dotted lines). From Munroe and Solis (1999).

Table 3. Classification of Pyralidae subfamilies (synonyms in italics).

Minet (1981[1982])	Solis and Mitter (1992)	Munroe and Solis (1999)
Galleriinae	Galleriinae	Galleriinae
<i>Macrothecinae</i>	<i>Macrothecinae</i>	<i>Cacotherapinae</i>
Chrysauginae	Chrysauginae	Chrysauginae
Pyralinae	Pyralinae	Pyralinae
Hypotiinae	<i>Hypotiinae</i>	<i>Hypotiinae</i>
Endotrichinae	<i>Endotrichinae</i>	<i>Endotrichinae</i>
Epipaschiinae	Epipaschiinae	Epipaschiinae
Phycitinae	Phycitinae	Phycitinae
<i>Anerastiinae</i>		<i>Anerastiinae</i>
<i>Peoriinae</i>	<i>Peoriinae</i>	<i>Peoriinae</i>

gene on three species of Pyraloidea showed two crambid species as sister group to one pyralid species (Regier *et al.* 1998)

The larvae of Pyraloidea exhibit diverse behavioral characteristics. Some are concealed feeders that fold, roll, web, or tie leaves together to form a shelter. Others are borers of monocots and can be found in the stems, roots, shoots, buds, fruits, or galls. A few mine leaves, make cases, or live in the nests of ants, bees, and wasps (selected references in Munroe and Solis 1999; Solis 1997). Some groups of pyraloids have adapted to aquatic environments (Lange 1956; Munroe 1972; Solis in press) where they mostly are plant feeders, but some have been observed as predators on other insects. The larvae of pyraloid species are scavengers, feeding on stored products or the fecal matter of sloths (Waage and Montgomery 1976) and bats (Solis and Mitter 1992). Pyraloids feed on a wide diversity of plants (Solis 1999a; Robinson *et al.* 2002), including monocots, dicots, and algae (Solis in press); one subfamily specializes on ferns (Solis *et al.* 2004b, 2005 *et al.*; Yen *et al.* 2004). Pyraloid larvae can be distinguished from other Lepidoptera by the following combination of characters: two prespiracular setae on the prothorax, crochets in a complete circle or penellipse (with the exception of aquatic immatures where they are usually in two rows), and three subventral setae on abdominal segments 3 to 6. Noctuids and carposinids also have two prespiracular setae on the prothorax, but noctuids have crochets in a mesoserie, and carposinids have four or




Figure 3. A. Pyralidae, sclerotized ring at base of SD1 of A8, Chrysaugine, sclerotized ring at base of SD1 of metathoracic segment 3. B. Pyralidae, sclerotized ring at base of SD1 of A8, Galleriinae, sclerotized ring at base of SD1 of abdominal segment 1.

introduced into the U.S. and has become a major pest on corn and other crops (Munroe and Solis 1999). There has been no higher level phylogenetic work on genera.

Subfamily Acentropinae. Acentropinae, formerly called Nymphulinae (Solis 1999b), has many species that are truly aquatic as immatures either with gills protruding from the integument or using a plastron for oxygen exchange. The larvae are mostly herbivorous, but a few have been shown to be predators of Simuliidae in Brazil (Yen 2004b; Solis in press). Most of the species in *Petrophila* are scrapers, feeding on algae and diatoms on the surfaces of rocks or other submerged objects, but other species feed on submerged plants, such as *P. drumalis* (Dyar, 1906), that feeds on the rootlets of water lettuce (*Pistia*) (Habeck and Solis 1994). Yoshiyasu (1985) defined the Acentropinae based on the protruding spiracles of the pupae on the second to fourth segments, but these characters have since been found to occur in members of the Musotiminae (Yen *et al.* 2004). Minet (1985) defined the subfamily by the presence of swollen scoloparia in the tympanal organs. Munroe (1972) reviewed the Nearctic Acentropinae, but his higher-level nomenclature is now outdated. There are no confirmed synapomorphies for this subfamily. There are several catalogs from China (Speidel and Mey 1999; You *et al.* 2002) and Chen has conducted several generic revisions from China (e. g.: 2006). Shen-Horn Yen (personal communication) is conducting a worldwide study of the genera of the Acentropinae.

Subfamilies Odontiinae, Evergestinae, Schoenobiinae, Scopariinae, Cybalomiinae, Musotiminae. Other smaller subfamilies of the Crambidae, such as Odontiinae, Evergestinae, Schoenobiinae, Scopariinae, Cybalomiinae, and Musotiminae, occur worldwide but are not very speciose in the Neotropics. Odontiinae was defined by Leraut and Luquet (1982) by the scalelike sclerotizations at the base of the vinculum of the male genitalia and two sets of non-deciduous scales on sternite VIII. The Evergestinae is even less well known; Minet (1981) defined the subfamily by derived characters of the tympanal organs. Schoenobiinae usually have long, narrow brown wings, but *Rupela*, a pest of rice, have brilliant white wings. Schoenobiinae are defined by a membranous area on sternite VIII covered by stiff scales from the posterior margin of sternite VI (Common 1990); the presence of platelike, scale-bearing coremata flanking the vinculum (Lewvanich 1981); a process on the lateral margin of the gnathos (Maes 1998); and a highly bilobed praecinctorium with a pair of circular formations called tubercula by Minet (1981, 1985). Larvae bore into grasses found in marshes and *Scirpophaga* in the Old World, and *Rupela* are pests of rice in the New World. Scopariinae moths are usually very small with black and white scales (Munroe 1972; Nuss 1999). Minet (1981) defined the subfamily by its longitudinal praecinctorium, reduced saccus tympani, and a well-developed gnathos in the male genitalia. Heliothelinae, currently is considered a synonym of Scopariinae, but this has been questioned by Nuss (1998) and Nuss *et al.* (1998). The most speciose genera, *Eudonia* and *Scoparia*, reach their greatest diversity on oceanic islands (Nuss 1999). The larvae tunnel in roots and stems of mosses and lycopods or feed on roots of vascular plants (Munroe and Solis 1999). Cybalomiinae is not well defined; nine species have been described in the Neotropics, but all but one belong in other subfamilies. Their placement in this subfamily is not tenable and is under investigation by Solis. Some larvae are known to feed on Cruciferae (Luquet and

Minet 1982). Musotiminae is a subfamily that was extracted from Acentropinae by Munroe (1972) based on the fact that most known species feed on ferns and do not have aquatic immatures. They have not been studied on worldwide basis, but Solis *et al.* (2004b) and Yen *et al.* (2004) have been transferring taxa to the Musotiminae and describing new taxa that feed on the Old World Climbing fern in southeast Asia and other ferns in the New World. Musotimines are defined by antennae that are laterally compressed, R_2 stalked with R_{3+4} , aedeagus with a reduced coecum, and tympanal cases enlarged (Minet 1981; Yoshiyasu 1985; Yen *et al.* 2004), but the relationship of Musotiminae with other subfamilies in Crambidae is unknown. Phillips and Solis (1996) studied *Neurophyseta* in Costa Rica; and Solis *et al.* (2005) described a new genus from Costa Rica that are leaf miners on a fern.

Subfamilies Linostinae, Midilinae, Glaphyriinae. Some subfamilies occur primarily in the Neotropics: Linostinae, Midilinae, and Glaphyriinae. The Linostinae is comprised of three species of *Linosta*. Although the characters that define this group are mainly absence of characters or secondary losses (reduced labial palpi and lack of maxillary palpi, proboscis, ocelli and chaetosemata (Munroe and Solis 1999)), they are easily recognizable by white forewings with distinct black lines. The immature stages are unknown. The Midilinae are spectacular moths with hyaline discal spots and sinuate terminal margins on red, green or yellow forewings; there are 47 described species (Munroe 1970). Minet (1981, 1985) and Munroe and Solis (1999) defined the subfamily on the basis of the reduced tympanal organs, a short praecinctorium, chaetosemata on the head often well delineated from the surrounding area, and an ostium bursae lacking a signum but with diffuse patches of spinules. The larvae are known to bore into stems and roots of Araceae. Solis and Adamski (1998), in their study of the Costa Rican Glaphyriinae, defined the subfamily by the absence of chaetosemata on the head, gnathos of the male genitalia, and the presence of a sclerotized antrum in the female genitalia. That study revealed that a generic revision of all 200 species in the Western Hemisphere is greatly needed. A few species of *Hellula* are pests of crucifers and occur worldwide. There has been no higher-level phylogenetic analysis of this subfamily.

Family Pyralidae

The Pyralidae is the smaller of the two families with a little less than 5,000 described species. It includes five subfamilies: Phycitinae, Epipaschiinae, Pyralinae, Chrysauginae, and Galleriinae. The monophyly of the Pyralidae is supported by the closed bullae tympani (Minet 1981, 1985), forewing veins R_3 and R_4 completely fused or stalked at their base, uncus arms in the male genitalia (Solis and Mitter 1992), and larvae with a sclerotized ring at the base of SD1 on A8 (Hasenfuss 1960). A phylogeny of the pyralid subfamilies proposed by Solis and Mitter (1992) (Fig. 5) shows support for the sister group relationship of the Epipaschiinae and Phycitinae, but the galleriines and chrysaugines were found to be unresolved basally.

Subfamily Galleriinae. Galleriinae occur worldwide and are defined by a lack of a gnathos (Whalley 1964; Munroe 1972). Although two genera of Chrysauginae also lack a gnathos (Cashatt 1968) it presumably has been lost independently in Galleriinae and Chrysauginae. Recently research on the

musculature (Solis unpublished) indicates that a remnant of a gnathos may be present but extremely difficult to see. Galleriine pupae have a prominent median ridge on the dorsum of the thorax and abdomen (Mosher 1916), and the larvae of most species have a sclerotized ring around SD1 of A1 (Hasenfuss 1960), except for *Omphalocera* and *Thyridopyralis* (Solis 1989; Solis and Mitter 1992). A world catalog of this subfamily was compiled by Whalley (1964), but no phylogenetic analyses have been conducted. This subfamily includes the bee or wax moth, *Galleria mellonella* (L., 1758), that feeds on honeycombs, and the rice moth, *Corcyra cephalonica* (Stainton, 1866). Other genera, such as *Aphomia*, have symbiotic relationships with other hymenopterans, and *Paralipsa* species are stored product pests worldwide (Munroe and Solis 1999).

Subfamily Chrysauginae. Chrysauginae are primarily neotropical (Cashett 1968). The monophyly of the subfamily is supported by a sclerotized ring at the base of SD1 on the metathorax (Hasenfuss 1960), but this feature is absent in *Azamora* (Solis 1989). A checklist of the Chrysauginae in the Western Hemisphere was compiled by Solis *et al.* (1995), but no phylogenetic analyses have been conducted. Four Australian genera were placed tentatively in this subfamily by Shaffer *et al.* (1996), but study of the adults did not confirm this, and the larvae are unknown. In the Western Hemisphere, this subfamily is home to the sloth moths. Adults of these species live in the fur, and the larvae feed on the sloth's dung (Waage and Montgomery 1976). Sloth moths are placed in several genera (*Cryptoses*, *Bradypodicola*, *Bradypophila*) that are

paraphyletic with respect to each other (being worked on Roe, pers. comm.).

Subfamily Pyralinae. Pyralinae is a worldwide group, but not very speciose in the New World. The subfamily is defined by a corpus bursae in the female genitalia that barely extends cephalad beyond abdominal segment 7 (Solis and Mitter 1992). A checklist of the Western Hemisphere and a morphological review of the Pyralinae was presented by Solis and Shaffer (1995, 1999), but no phylogenetic analyses have been conducted. The subfamily includes the meal moth, *Pyralis farinalis* L., 1758, and other species that are pests of stored product, leaf feeders, and one species, *P. manihotalis* Guenée, 1854, that has been reared from bat guano (Solis and Mitter 1992).

Subfamily Epipaschiinae. Epipaschiinae were examined by Solis and Mitter (1992) and Solis (1993), and three synapomorphies were discovered to define it. The third segment of the labial palpus is upturned and pointed, the phallobase of the aedeagus extends beyond the ductus ejaculatorius and is curved ventrally, and there is weakness in the sclerotization at the base of the tegumen that results in the appearance of a separate sclerite. Larvae are known to be leaf rollers, leaf tiers, and leafminers, but their morphology is not well known. The group includes minor pests of avocado (e. g.: *Accinctapubes*), mahoganies (e. g.: *Macalla*), and corn (e. g.: *Phidotricha*). Solis (1993) provided a phylogenetic analysis of 20 genera of the *Pococera* complex, which consists of 300 species in the Western Hemisphere, representing the first cladistic analysis of a pyraloid group. A similar study of about 500 Old World species is needed. Checklists of Old World and New World epipaschiines have been published (Solis 1992; Solis 1995).

Subfamily Phycitinae. The Phycitinae is the largest subfamily, consisting mostly of small moths worldwide. It is defined by larvae with a sclerotized circle at the base of seta SD1 on the mesothorax (Hasenfuss 1960). The female *frenulum* has one bristle as in the male (females have two bristles in pyralines and epipaschiines, three in galleriines and chrysaugines), and the ductus seminalis originates in the corpus bursae (in other pyralid subfamilies it originates in the ductus bursae) (Solis and Mitter 1992). Shaffer (1968, 1976) has revised many groups of the Peoriini. The Phycitinae includes many stored product pests, including *Plodia*, *Cadra*, *Ectomyelois*, and *Etiella* (Solis 1999a). The larvae are leaf rollers and borers in various parts of plants, although a few are known to be predators of Homoptera (e. g.: *Laetilia*). *Cactoblastis cactorum* (Berg, 1885), the cactus moth is the classical biological control agent of *Opuntia* in various countries, but in recent years has become an introduced invasive in the United States (Solis *et al.* 2004a). The New World Phycitinae were revised by Heinrich (1956), the Nearctic members by Neunzig (e. g.: 1986, 1990, 1997, 2003), and the Old World *Homeosoma* species by Roesler (1965). Whalley (1970) produced a world catalog of the Phycitinae. Recently, Neunzig and Solis (e.g. 2002, 2004, 2005a,b, 2006) have published a number of papers regarding Costa Rican phycitines, Horak (e. g.: 1998) has published on Australian Phycitinae, and Li and coauthors have published on Chinese genera (e. g.: Li and Ren 2006). Simonsen (pers. comm.) is conducting research on the cactus feeding phycitines. Recent studies on *Dioryctria*, which are pests of conifers in the Holarctic region, indicates that combining morphological and molecular characters increased nodal support in hypothesized phylogenetic trees (Du *et al.* 2005; Roe *et al.* 2006).




Figure 5. Two most parsimonious cladograms for the Pyralidae. Numbers outside parentheses indicated character number and numbers inside parentheses described character state transformations. From Solis and Mitter (1992).

Literature Cited

- ALLYSON, S. 1981. Last instar larve of Pyraustini of America north of Mexico (Lepidoptera: Pyralidae). *Canadian Entomologist* 113: 463-518.
- ALLYSON, S. 1984. Description of last-instar larvae of 22 species of North American Spilomelini (Lepidoptera: Pyralidae: Pyraustinae) with a key to species. *Canadian Entomologist* 116: 1301-1334.
- BECCALONI, G. W.; SCOBLE, M. J.; ROBINSON, G. S.; PITKIN, B. (eds.). 2003. The Global Lepidoptera Names Index (LepIndex). World Wide Web electronic publication. <http://www.nhm.ac.uk/entomology/lepindex>. Last revision: 1 February 2005. Last access: 15 May 2007.
- BLESZYNSKI, S. 1969. The taxonomy of the crambine moth borers of sugar cane, pp. 11-59. In: Williams, J. R.; Metcalf, J. R.; Mungomery, R. W.; Mather, R. (eds.). *Pests of Sugar Cane*. New York: Elsevier Publishing Co.
- BLESZYNSKI, S. 1970. A revision of the world species of the genus *Chilo* Zincken. (Lepidoptera: Pyralidae). *Bulletin of the British Museum (Natural History) Entomology* 25 (4): 101-195.
- BÖRNER, C. 1925. Lepidoptera, Schmetterlinge. pp. 382-421. In: Bröhm, P. (ed.). *Fauna von Deutschland*: Berlin: Leipzig, Quelle & Meyer: 561 p.
- BOX, H. E. 1931. The crambine genera *Diatraea* and *Xanthopherne* (Lep. Pyralidae). *Bulletin Entomological Research* 22: 1-50.
- CASHATT, E. D. 1968. Revision of the Chrysauginae of North America. Dissertation, Catholic University of America, Washington, D.C. 179 p.
- CHEN, F.; SONG, S.; WU, C. 2006. A review of the genus *Eristena* Warren in China (Lepidoptera: Crambidae: Acentropinae). *Aquatic Insects* 28 (3): 229-241.
- CLARKE, J. F. G. 1971. The Lepidoptera of Rapa Island. *Smithsonian Contributions Zoology* 56: 1-282.
- CLAVIJO A., J. A. 1990. Systematics of black and white species of the genus *Diaphania* Hübner (1818) (Lepidoptera: Pyralidae: Pyraustinae). Dissertation: Montreal, Canada: McGill University. 215 p.
- COMMON, I. F. B. 1990. Pyraloidea, pp. 343-358. In: *Moths of Australia*. Carlton, Victoria: Melbourne University Press.
- DU, Y.; A. D. ROE; F.A.H. SPERLING. 2005. Phylogenetic framework for *Dioryctia* (Lepidoptera: Pyralidae: Phycitinae) based on combined analysis of mitochondrial DNA and morphology. *Canadian Entomologist* 137: 685-711.
- FLETCHER, D. S.; I. W. B. NYE. 1984. The Generic Names of the Moths of the World. London: Trustees of the British Museum (Natural History): Vol. 5, 185 p.
- GASKIN, D. E. 1975. Revision of the New Zealand Crambini (Lepidoptera: Pyralidae: Crambinae). *New Zealand Journal Zoology* 2 (3): 265-363.
- GASKIN, D. E. 1985. Morphology and reclassification of the Australasian, Melanesian and Polynesian *Glauchocharis* Meyrick (Lepidoptera: Crambinae: Diptychophorini). *Australian Journal Zoology Supplement Series* 115: 1-75.
- GENTILI, P.; SOLIS, M. A. 1998. Checklist and key of New World species of *Omiodes* Guenée with descriptions of four new Costa Rican species (Lepidoptera: Crambidae). *Entomologica Scandinavica* 28: 471-492.
- HABECK, D.; SOLIS M. A. 1994. Transfer of *Petrophila drumalis* (Dyar) to *Argyractis* based on immature and adult characters with a larval description of *Argyractis subornata* (Hampson) (Lepidoptera: Crambidae: Nymphulinae). *Proceedings Entomological Society Washington* 96 (4): 726-734.
- HAMPSON, G. F. 1895. On the classification of the Schoenobiinae and Crambinae, two subfamilies of moths, of the family Pyralidae. *Proceedings of the Zoological Society of London* 1895: 897-974.
- HASENFUSS, I. 1960. *Die Larvalsystematik der Zünsler*. Berlin: Akademie Verlag: 263 p.
- HEINRICH, C. 1956. American moths of the subfamily Phycitinae. *Bulletin of the United States National Museum* 207: 1-581.
- HORAK, M. 1998. A reassessment and review of the Australian genus *Cternomeristis* Meyrick (Lepidoptera: Pyralidae: Phycitinae). *Entomologica Scandinavica* 28: 445-470.
- LANDRY, B. 1995. A phylogenetic analysis of the major lineages of the Crambinae and of the genera of the Crambini of North America (Lepidoptera: Pyralidae). *Memoirs of the American Entomological Institute* 1:1-242.
- LANGE, H. 1956. Aquatic Lepidoptera, pp. 271-288. In: Usinger, R. L. (ed.). *Aquatic Insects of California*. University of California Press. Berkeley. 508 p.
- LERAUT, P.; LUQUET, G. C. 1982. Statut de quelques genres et espèces d'Odontiinae paléarctiques et description de quatre nouveaux taxa (Lep. Crambidae). *Linneana Belgica* 8 (12): 527-555.
- LEWVANICH, A. 1981. A revision of the Old World species of Scirpophaga (Lepidoptera: Pyralidae). *Bulletin of the British Museum (Natural History) Entomology* 42 (4): 185-298.
- LI, H.-H.; REN, Y.-D. 2006. The genus *Thiallela* Walker, 1863 in China (Lepidoptera: Pyralidae: Phycitinae). *Entomological News* 117 (3): 323-331.
- LUQUET, G. C.; MINET, J. 1982. Découverte dans le Vaucluse d'une nouvelle espèce de Pyrale et designation de l'espèce-type du genre *Krombia* Chrétien (Lepidoptera, Crambidae, Cybalomiinae). *Alexandria* 12 (7): 317-326.
- MAES, K. V. N. 1987. Revisionary notes on the genus *Achyra* Guenée with a new synonym and the description of *Achyra takowensis* sp. n. (Lepidoptera: Pyralidae, Pyraustinae) (Studies on Pyralidae I). *Nota Lepidopterologica* 10 (3): 175-182.
- MAES, K. V. N. 1994. Some notes on the taxonomic status of the Pyraustinae (sensu Minet 1981 [1982]) and a check list of the Palearctic Pyraustinae (Lepidoptera, Pyraloidea, Crambidae). *Annales de la Société Royale Entomologique de Belgique* 130: 159-168.
- MAES, K. V. N. 1995. A comparative morphological study of the adult Crambidae (Lepidoptera, Pyraloidea). *Annales de la Société Royale Entomologique de Belgique* 131: 383-434.
- MAES, K. V. N. 1998. On the morphology of the gnathos in the Pyraloidea (Lepidoptera). *Entomologica Scandinavica* 28 (4): 381-390.
- MAES, K. V. N. 2002. On the systematic position of some Palearctic Pyraustinae (Pyraloidea: Crambidae). *Nota Lepidopterologica* 24 (4): 51-57.
- MINET, J. 1981 (1982). Les Pyraloidea et leurs principales divisions systématiques. *Bulletin de la Société Entomologique de France* 86: 262-280.
- MINET, J. 1983. Étude morphologique et phylogénétique des organes tympaniques des Pyraloidea. 1. Généralités et homologies (Lepidoptera Glossata). *Annales de la Société Entomologique de France* 19 (2): 175-207.
- MINET, J. 1985. Étude morphologique et phylogénétique

- des organes tympaniques des Pyraloidea. 2 - Pyralidae, Crambidae, première partie (Lepidoptera Glossata). Annales de la Société Entomologique de France 21 (1): 69-86.
- MINET, J. 1986. Ébauche d'une classification moderne de l'ordre des Lépidoptères. *Alexanor* 14 (7): 291-313.
- MINET, J. 1991. Tentative reconstruction of the ditrysian phylogeny (Lepidoptera: Glossata). *Entomological Scandinavica* 22: 69-95.
- MOSHER, E. 1916. A classification of the Lepidoptera based on characters of the pupa. *Bulletin of the Illinois State Laboratory of Natural History* 12: 15-159.
- MUNROE, E. 1970. Revision of the subfamily Midilinae (Lepidoptera: Pyralidae). *Memoirs of the Entomological Society of Canada* 74: 1-94.
- MUNROE, E. 1972. (Fasc. 13.1A-1B), 1973 (Fasc. 13.1C), 1976 (Fasc. 13.2A-B). Pyraloidea. Pyralidae (in part). In: Dominick, R. B.; Ferguson, D. C.; Franclemont, J. G.; Hodges, R. W.; Munroe, E. G. (eds.). *The Moths of America North of Mexico*. London: E.W. Classey, Ltd. and The Wedge Entomological Research Foundation: 1- 304, 1-150.
- MUNROE, E.; SOLIS, M. A. 1999. Pyraloidea, pp. 233-256. In: Kristensen, N. (ed.). *Lepidoptera, Moths and Butterflies, Vol. 1, Arthropoda, Insect, Vol.4, Part 35. Handbook of Zoology*. Walter de Gruyter & Co. Berlin. 491 p.
- MUTUURA, A.; MUNROE, E. M. 1970. Taxonomy and distribution of the European corn borer and allied species: genus *Ostrinia* Hübner (Lepidoptera: Pyralidae). *Memoirs of the Entomological Society of Canada* 71: 1-112.
- NEUNZIG, H. H. 1986. (Fasc. 15.2), 1990 (Fasc. 15.3), 1997 (Fasc. 15.4), 2003 (Fasc. 15.5). Pyraloidea, Pyralidae (in part). In: Dominick, R. B.; Ferguson, D. C.; Franclemont, J. G.; Hodges, R. W.; Munroe, E. G. (eds.). *The Moths of America North of Mexico*. Lawrence, Kansas: Allen Press, Inc.: 1-112, 1-165, 1-157, 1-338.
- NEUNZIG, H. H. 1987. Pyralidae (Pyraloidea), pp. 462-494. In: Stehr, F. W. *Immature Insects*. Dubuque, Iowa: Kendall Hunt Publishing Co.
- NEUNZIG, H. H.; SOLIS, M. A. 2002. The *Ceracanthia* complex (Lepidoptera: Pyralidae: Phycitinae) in Costa Rica. I. *Ceracanthia*. *Proceedings Entomological Society of Washington* 104 (4): 837-855.
- NEUNZIG, H. H.; SOLIS, M. A. 2004. *Exguiana*, a new genus of Neotropical phycitines (Lepidoptera: Pyralidae). *Proceedings of the Entomological Society of Washington* 106 (3): 544-563.
- NEUNZIG, H. H.; SOLIS, M. A. 2005a. *Tumoriala*, a new neotropical phycitine genus (Lepidoptera: Pyralidae). *Proceedings of the Entomological Society of Washington* 107 (1): 84-89.
- NEUNZIG, H. H.; SOLIS, M. A. 2005b. A review of the genus *Difundella* (Lepidoptera: Pyralidae: Phycitinae). *Proceedings of the Entomological Society of Washington* 107 (2): 303-314.
- NEUNZIG, H. H.; SOLIS, M. A. 2006. Four new species of Costa Rican Phycitinae (Lepidoptera: Pyralidae). *Proceedings of the Entomological Society of Washington* 108 (4): 953-963.
- NIELSEN, E. S. 1989. Phylogeny of major lepidopteran groups. In: Fernholm, B.; Bremer, K.; Jörnvall, H. (eds.). *The Hierarchy of Life*. Copenhagen: Elsevier Science Publishers: 281-294.
- NUSS, M. 1998. The Scopariinae and Heliothelinae stat. Rev. (Lepidoptera: Pyraloidea: Crambidae) of the Oriental Region: a revisoinal synopsis with descriptions of new species from the Phillippines and Sumatra. *Nachrichten des entomologischen Vereins Apollo Supplement* 17: 475-528.
- NUSS, M. 1999. Revision der Gattungen der Scopariinae, Lepidoptera: Pyraloidea, Crambidae. *Nova supplementa Entomologica* 13: 1-151.
- NUSS, M. 2006. GLOBIZ, State Museum of Zoology, Dresden, Germany. http://globiz.sachsen.de/pyrdat_prototype/GlobIZ/globiz_start/index_start.html. Last revision: 16 May 2006. Last access: 1 May 2007.
- NUSS, M.; KARSHOLT, O.; MEYER, M. 1998. A taxonomic revision of the Scopariinae from the Macaronesian Region (Lepidoptera: Pyraloidea: Crambidae). *Entomological Scandinavica* 28: 509-551.
- PATOCKA, J.; TURCANI, M. 2005. *Lepidoptera Pupae, Central European Species*. Apollo Books. Denmark. 2 volumes 542 p., Plates 321 p.
- PHILLIPS, E.; SOLIS, M. A. 1996. *Neurophyseta* (Lepidoptera: Crambidae) de Costa Rica. *Revista de Biología Tropical* 44 (2): 693-717.
- REGIER, J. C.; FANG, Q. Q.; MITTER, C.; PIEGLER, R. S.; FREIDLANDER, T. P.; SOLIS, M. A. 1998. Evolution and Phylogenetic Utility of the *period* gene in Lepidoptera. *Molecular Biology and Evolution* 15 (9): 1172-1182.
- ROBINSON, G. S.; ACKERY, P.R.; KITCHING, I. J.; BECCALONI, G. W.; HERNANDEZ, L. M. 2002. Hostplants of the moth and butterfly caterpillars of America North of Mexico. *Memoirs of the American Entomological Institute* 69: 1-824. <http://www.nhm.ac.uk/entomology/hostplants>. Last revision: 25 March 2002. Last access: 25 April 2007.
- ROESLER, R. U. 1965. Untersuchungen über die Systematik und Chorologie des Homoeosoma-Ephestia-K[o]mplexes (Lepidoptera: Pyralidae). Saarbrücken: Dissertation Saarbrücken (1964) 265 p.
- ROE, A. D.; STEIN, J.D.; GILLETTE, N. E.; SPERLING, F.A.H. 2006. Identification of *Dioryctria* (Lepidoptera: Pyralidae) in a Seed Orchard at Chico, California. *Annals of the Entomological Society of America* 99 (3): 433-448.
- SCHOUTEN, R. T. A. 1992. Revision of the genera *Euchromius* Guenée and *Miyakea* Marumo (Lepidoptera: Crambidae: Crambinae). *Tijdschrift voor Entomologie* 135: 191-274.
- SHAFFER, J. C. 1968. A revision of the Peoriinae and Anerastiinae (auctorum) of America north of Mexico (Lepidoptera:Pyralidae). *Bulletin of the United States National Museum* 280: 1-124.
- SHAFFER, J. C. 1976. A revision of the neotropical Peoriinae (Lepidoptera: Pyralidae). *Systematic Entomology* 1: 281-331.
- SHAFFER, M.; NIELSEN, E. S.; M. HORAK. 1996. Pyraloidea, pp. 164-199 In: Nielsen, E.S.; Edwards, E. D.; Rangsi, T. V. (eds.). *Checklist of the Lepidoptera of Australia*. Monographs on Australian Lepidoptera 4: 1-529.
- SOLIS, M. A. 1989. A Phylogenetic analysis and reclassification of the *Pococera*-complex Genera (Lepidoptera: Pyralidae: Epipaschiinae) and a Preliminary Cladistic Analysis of the Pyralidae. Dissertation. University of Maryland, College Park. 428 p.
- SOLIS, M. A. 1992. Check list of the Old World Epipaschiinae and the related New World genera *Macalla* and *Epipaschia* (Pyralidae). *Journal of the Lepidopterists' Society* 46 (4): 280-297.
- SOLIS, M. A. 1993. A phylogenetic analysis and reclassification of the genera of the *Pococera*-Complex (Lepidoptera: Pyralidae: Epipaschiinae). *Journal of the New York*

- Entomological Society 101: 1-83.
- SOLIS, M. A. 1995. Epipaschiinae, pp. 89-93. In: Heppner, J. B. (ed.). Check List Part 2: Atlas of Neotropical Lepidoptera, Brill/Flora and Fauna Books, Gainesville, FL. 243 p.
- SOLIS, M. A. 1997. Snout moths: Unraveling the taxonomic diversity of a speciose group in the Neotropics, pp. 231-242. In: Reaka-Kudla, M. L.; Wilson, D.; Wilson, E. O. (eds.). Biodiversity II: Understanding and Protecting our Biological Resources. Joseph Henry Press, Washington, D. C. 551 p.
- SOLIS, M. A. 1998. *Michaelshaffera* gen. n. - a pyraloid taxon lacking an abdominal tympanal organ (Lepidoptera: Pyralidae). Entomological Scandinavica 28: 391-402.
- SOLIS, M. A. 1999a. Key to selected Pyraloidea (Lepidoptera) larvae intercepted at U.S. Ports of entry: Revision of Pyraloidea in "Keys to some frequently intercepted lepidopterous larvae" by D. M. Weisman, 1986. Proceedings of the Entomological Society of Washington 101 (3): 645-686.
- SOLIS, M. A. 1999b. Nymphulinae Duponchel [1845] (Insecta: Lepidoptera): proposed precedence over Acentropinae Stephens, 1836. Case 3048. Bulletin of Zoological Nomenclature 56 (1): 31-33.
- SOLIS, M. A. 2004. Systematics of Mexican stalkboring crambine Pyraloidea, pp. 6- 22. In: Rodríguez del Bosque, L.A.; Vejar Cota, G.; Cortez Mondaca, E. (eds.). Taller Internacional sobre Barrenadores del Tallo de Caña de Azúcar, Los Mochis, Sinaloa, México. Sociedad Mexicana de Control Biológico. 104 p.
- SOLIS, M. A. In press. Aquatic Lepidoptera. In: Merritt, R. W.; Cummins, K. W. (eds.). Aquatic Insects of North America, Kendall/Hunt Publishing Company.
- SOLIS, M. A.; ADAMSKI, D. 1998. Review of the Costa Rican Glaphyriinae (Lepidoptera: Pyraloidea: Crambidae). Journal of the New York Entomological Society 106 (1): 1-55.
- SOLIS, M. A.; MAES, K. V. N. 2002. Preliminary phylogenetic analysis of the subfamilies of Crambidae (Pyraloidea: Lepidoptera). Belgian Journal of Entomology 4: 53-95.
- SOLIS, M. A.; MITTER, C. 1992. Review and preliminary phylogenetic analysis of the subfamilies of the Pyralidae (sensu stricto) (Lepidoptera: Pyraloidea). Systematic Entomology 17: 79-90.
- SOLIS, M. A.; SHAFFER, M. 1995. Pyralinae, pp. 80-81. In: J.B. Heppner, ed. Check List Part 2: Atlas of Neotropical Lepidoptera, Brill/Flora and Fauna Books, Gainesville, FL. 243 p.
- SOLIS, M. A.; SHAFFER, M. 1999. Contribution towards the study of the Pyralinae (Pyralidae): historical review, morphology, and nomenclature. Journal of the Lepidopterists' Society 53 (1): 1-10.
- SOLIS, M. A.; MUNROE, E.; BECKER, V. O. 1995. Chrysauginae, pp. 81-88. In: Heppner, J. B. (ed.). Check List Part 2: Atlas of Neotropical Lepidoptera, Brill/Flora and Fauna Books, Gainesville, FL. 243 p.
- SOLIS, M.A.; HIGHT, S. D.; GORDON, D. R. 2004a. Alert: Tracking the cactus moth as it flies and eats its way westward in the U.S. News of the Lepidopterists' Society 46 (1): 3-4.
- SOLIS, M. A.; YEN, S.-H.; GOOLSBY, J. R. 2004b. Description and life history of *Lygomusotima* New Genus, and *Neomusotima conspurcatalis* (Lepidoptera: Crambidae) from Australia and Southeast Asia feeding on *Lygodium microphyllum*. Annals entomological Society of America 97 (1): 64-76.
- SOLIS, M. A.; DAVIS, D. R.; NISHIDA, K. 2005. Biology and systematics of *Albusambia elaphaglossumae*, a new genus and species of Crambidae (Lepidoptera: Pyraloidea: Crambidae) mining the fronds of *Elaphoglossum conspersum* (Pteridophyta: Lomariopsidaceae) in Costa Rica. International Journal of Tropical Biology 53 (3-4): 487-501.
- SPEIDEL, W.; MEY, W. 1999. Catalog of the Oriental Acentropinae (Lepidoptera, Crambidae). Tijdschrift voor Entomologie 142: 125-142.
- SUTRISNO, H. 2002a. Cladistic analysis of the Australian *Glyphodes* Guenée and allied genera (Lepidoptera: Crambidae: Spilomelinae). Entomological Science 5 (4): 457-467.
- SUTRISNO, H. 2002b. A preliminary study on relationships among selected Australian members of the Tribe Spilomelini (Lepidoptera: Crambidae: Pyraustinae). Zoological Science 19: 915-929.
- SUTRISNO, H.; AZUMA, N.; HIGASHI, S. 2006. Molecular phylogeny of the Indo-Australia *Glyphodes* and its allied genera (Insecta: Lepidoptera: Crambidae: Spilomelinae) inferred from mitochondrial COI and COII and nuclear EF-1 α gene sequences. Species Diversity 11: 57-69.
- YEN, S. H. 2004. Insecta: Lepidoptera, Crambidae, Acentropinae, pp. 545-554. In: Yule, C. M.; Sen, Y. H. (eds.). Freshwater invertebrates of the Malaysian Region. Academy of Sciences Malaysia. Kuala Lumpur, Malaysia. 861 p.
- YEN, S. H.; SOLIS, M. A.; GOOLSBY, J. A. 2004. *Austromusotima*, a new musotimine genus (Lepidoptera: Crambidae) feeding on Old World climbing fern. Annals of the Entomological Society of America 97 (3): 397-410.
- YOSHIYASU, Y. 1985. A systematic study of the Nymphulinae and the Musotiminae of Japan (Lepidoptera: Pyralidae). Scientific Report of Kyoto Prefectural University 37: 1-162.
- YOU, P.; LI, H.-H.; CHEN, K.-L. 2002. Catalogue of the Nymphulinae and Musotiminae of China (Lepidoptera: Crambidae). Fauna of China 4: 37-64.
- WAAGE, J. K.; MONTGOMERY, G. G. 1976. *Cryptoses choloepi*: A coprophagous moth that lives on a sloth. Science 193: 157-158.
- WHALLEY, P. E. S. 1964. Catalogue of the Galleriinae (Lepidoptera, Pyralidae) with descriptions of new genera and species. Acta Zoologica Cracoviensia 9 (10): 561-617.
- WHALLEY, P. E. S. 1970. A synonymic catalogue of the genera of the Phycitinae of the world. Bulletin of the British Museum (Natural History) Entomology 25: 33-72.